# THE RATE AND PREDICTORS OF CONTINUOUS POSITIVE AIRWAY PRESSURE ADHRENCE IN CANADIAN PATIENTS WITH OBSTRUCTIVE SLEEP APNEA

by

# Morvarid Mehrtash

BSFN, The University of British Columbia, 2016

# A THESIS SUBMITTED IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR THE DEGREE OF

# MASTER OF SCIENCE

in

The Faculty of Graduate and Postdoctoral Studies

(Experimental Medicine)

THE UNIVERSITY OF BRITISH COLUMBIA (Vancouver)

December 2018

© Morvarid Mehrtash, 2018

and Postdoctoral Studies for acceptance, the thesis entitled:
The rate and predictors of continuous positive airway pressure adherence in Canadian patients
with obstructive sleep apnea
submitted by Morvarid Mehrtash in partial fulfillment of the requirement for
the degree of Master of Science
in
Experimental Medicine
Examining Committee:
Najib T. Ayas
Supervisor
Ismail Laher
Supervisory Committee Member
Glen Foster
Additional Examiner

The following individuals certify that they have read, and recommend to the Faculty of Graduate

## **Abstract**

mortality. Continuous positive airway pressure (CPAP) is the first-line treatment for patients with moderate to severe OSA. CPAP ameliorates respiratory disturbances, leading to improvements in daytime sleepiness, quality of life, blood pressure, and cognition. However, despite its high efficacy, CPAP adherence is often sub-optimal. There are many factors that are proposed to affect CPAP adherence including: socio-demographic/economic characteristics, disease severity, psychological factors, side-effects, and initial experience with CPAP. Intervention studies have suggested that augmented support/education, cognitive behavioral therapy, and telemedicine may improve CPAP adherence. However, no studies regarding the rate and predictors of CPAP have been done within a Canadian context. Consequently, we sought to determine the rate and predictors correlated with CPAP adherence amongst Canadians. **Methods:** Patients referred for suspected OSA to the University of British Columbia Hospital Sleep Clinic between 2003 and 2009 were recruited. All patients underwent full polysomnography and completed a health questionnaire. Patient charts were reviewed, and CPAP adherence was defined as using the device for  $\geq 4$  hours for  $\geq 70\%$  of the time being used. CPAP predictors including severity of OSA, daytime sleepiness, age, BMI, depression, etc. were analysed. A P value of < 0.05 was considered to be statistically significant. **Results:** There were 1,250 patients in total; 844 were prescribed CPAP and 406 were not. Of those who were prescribed CPAP, 421 (49.9%) adhered to CPAP and 423 (50.1%) did not. In the multivariate model, the severity of OSA measured by AHI was the only predictor of CPAP adherence (p < 0.0001). The severity of OSA in adherers and non-adherers was 34 vs. 27

events/hr respectively. Patients in both groups were on average 51 years old, mostly males,

mostly non-smokers, had an Epworth sleeping scale (ESS) of 11, depression score of 16 and

Rationale: Obstructive sleep apnea (OSA) impacts mood, cardiovascular morbidity, and

were mostly involved in car crashes. Those with a higher education ( $\geq$  college) were more adherent (63% vs. 57%) (p = 0.14). But this was not significant.

**Conclusions:** In our data, CPAP adherence was low (49.9%) and OSA severity, as measured by AHI, was its only significant predictor.

# Lay summary

Obstructive sleep apnea (OSA) is a common sleep disorder affecting 5-10% of the general adult population. Continuous positive airway pressure (CPAP) is its main treatment and is effective in improving the harmful side effects of OSA. Yet, adherence rate to CPAP treatment ranges from 30-60% despite various improvements to the mask and the device itself. The published studies regarding the predictors of CPAP adherence are inconsistent in their findings and have not been investigated within a Canadian context. Consequently, the goal of this study was to find predictors of CPAP adherence amongst Canadians. By doing so, patient care and quality of life may improve amongst Canadians. Patients referred to University of British Columbia's Sleep Clinic were recruited between 2003-2009. Overall, 844 patients were prescribed CPAP but only 421 (49.9%) were adherent and the only predictor associated with CPAP adherence was OSA severity as measured by AHI.

# **Preface**

The work presented herein was part of an ongoing research study, which started in 2003 at the sleep clinic at the University of British Columbia. This study was designed by the principle investigator Dr. Najib Ayas and approved by the UBC Clinical Research Ethics board H13-00346.

A version of chapter 1 has been sent for publication to the Journal of Lung and the manuscript is being published. Mehrtash, M., Bakker, J. P., & Ayas, N. (2018). Predictors of Continuous Positive Airway Pressure Adherence in Patients with Obstructive Sleep Apnea. I conducted a broad research of the literature regarding OSA, predictors of CPAP adherence and interventions that may improve adherence. I wrote most of the manuscript.

This thesis is the original intellectual product of the author, M. Mehrtash, who was responsible for collecting and analysis of data. Dr. N. Ayas was involved in the early stages of developing the concept and the design of the study. He also supported concept development throughout the duration of the research and contributed thesis edits. N. Fox was in charge of ordering the medical charts from Iron Mountain Inc. B. Peres and A. Hirsch were in charge of helping clean the data from the medical charts and the health questionnaire and merging the datasets.

# **Table of Contents**

Abstract	iii
Lay summary	v
Preface	vi
List of abbreviations	xi
Acknowledgements	xiii
Dedication	xiv
Chapter 1: Introduction	
1.1 Purpose and objective	
1.2 Methods	
1.3 Significance	5
Chapter 2: Background and Literature Review	6
2.1 Socio-demographic characteristics	
2.2 Disease severity	9
2.3 Psychosocial factors	12
2.4 Side-effects	14
2.5 Interventions to improve CPAP adherence	17
2.6 Rationale	
Chapter 3: Methods	
3.1 Overall study design	25
3.2 Patient recruitment	27
3.3 Inclusion criteria	27
3.4 Risks and confidentiality	28
3.4 Polysomnography	28
3.5 Obtaining patient charts and the health questionnaire	29
3.6 Data sets and statistical analyses	30
3.7 Potential predictors of CPAP adherence	32
Chapter 4: Results	
4.1 The study population	33
4.2 The correlation between different predictors of CPAP adhe	
did not adhere to CPAP	
4.2.2 Age and adherence	
4.2.3 ESS and adherence	37
4.2.4 BMI and adherence	

4.2.6 Menopausal status and adherence	39
4.2.7 Antidepressant medication and adherence	40
4.2.8 Gender and adherence	
4.2.9 Smoking status and adherence	
4.2.10 Restless leg syndrome and adherence	41
4.2.11 Marital status and adherence	41
4.2.12 Bed partner and adherence	
4.2.13 Snoring severity and adherence	42
4.2.14 Occupational injury and adherence	
4.2.15 Neck circumference and adherence	
4.2.16 Mood disorder and adherence	
4.2.17 Anxiety disorder and adherence	
4.2.18 Periodic limb movement and adherence	
4.2.19 Hypertension and adherence	
4.2.20 Myocardial infraction and adherence	
4.2.21 Cardiac arrhythmia and adherence	
4.2.22 Stroke and adherence	
4.2.24 Alcohol consumption and adherence	
4.2.25 Education and adherence	
4.2.26 Car crashes and adherence	
4.2.27 Sleeping aid medication and adherence	46
Chapter 5: Discussion	47
5.1 Summary of the results	47
5.2 CPAP adherence	47
5.3 OSA severity	48
5.4 Strengths	48
5.5 Limitations	49
5.6 Conclusions	
5.7 Future studies	
Bibliography	52
Appendices	62
Annandiy A. Haalth quastiannaira	63

# **List of Tables**

Table 1.2. Potential predictors of CPAP adherence.	4
Table 2.0. Factors associated with CPAP adherence.	6
Table 2.1. Overview of socio-demographic characteristics variables influencing CPA	P
adherence.	8
Table 2.2.1. Overview of severity of symptoms influencing CPAP adherence	10
Table 2.2.2. Overview of severity of OSA (AHI) influencing CPAP adherence	11
Table 2.3. Overview of psychosocial variables influencing CPAP adherence	13
Table 2.4. Overview of side-effect variables influencing CPAP adherence	16
Table 2.5. Interventional studies to improve CPAP adherence	
Table 4.2.1. The correlation between continuous variables and CPAP adherence in	
adherent and non-adherent group using t-test	34
Table 4.2.2. Frequency of categorical variables by adherent and non-adherent using	
squared test.	
Table 4.2.3. OSA severity categories and CPAP adherence.	
Table 4.2.4. BMI categories and CPAP adherence.	

# **List of Figures**

Figure 3.1. Overall study design.	26
Figure 4.1. Cohort sample size	33
Figure 4.2.1. Association between AHI and CPAP adherence.	

# List of abbreviations

**AHI** Apnea Hypopnea Index

**APE** Active patient engagement tool

**BMI** Body Mass Index

**CBT** Cognitive behavioural therapy

**CPAP** Continuous Positive Airway Pressure

**ECG** Electrocardiographic

**EDS** Excessive daytime sleepiness

**EEG** Electroencephalographic

**EMG** Electromyographic

**EOG** Electrooculographic

**ESS** Epworth Sleepiness Scale

HR Hypopneas recorded

NA Not available

**ODI** Oxygen Desaturation Index

**OSA** Obstructive Sleep Apnea

**PHQ-9** Patient Health Questionnaire

**PLM** Periodic limb movement

**PSG** Polysomnography

**RCT** Randomized controlled trial

SaO2 oxygen saturation

SAS Statistical Analysis System

**SD** Standard deviation

**Tel-Ed** Web-based OSA education

**Tel-TM** CPAP telemonitoring with automated patient feedback

TLC Telephone-linked communications

**UBC** University of British Columbia

UCM Usual care monitoring

# **Acknowledgements**

Special thanks are owed to my parents, sisters and husband who have supported me throughout my years of education, both morally, and emotionally. I have come a long way to be where I am today, and I could not have done it without them and their support. My husband, Ali Rafie, showed me how to fight for my goals and how to achieve them one by one. He never asked me to stop working and he always pushed me towards achieving my goals. My parents, Hashem and Mitra Mehrtash, have done so much for me that I cannot simply list them. I owe them a big thank you for bringing me to Canada so that I could have a brighter future. I would not be writing this thesis if it were not for them. My success in life is due to their bravery and encouragement. My sisters, Sara and Sadaf, gave positive beliefs in my success that kept me going regardless of the challenges I was facing.

I owe a particular thanks to Dr. Najib Ayas, whose penetrating questions taught me to question more deeply and for enlarging my vision of science and providing coherent answers to my endless questions. His support, guidance and knowledge have pushed me closer to my goals in life. I hope that Dr. Ayas is as proud to be my mentor as I am to be his mentee. I offer my enduring gratitude to the faculty, staff and my fellow students at the University of British Columbia, who have inspired me to continue my work in this field.

# **Dedication**

To my loving family, who have supported me endlessly no matter the challenges I faced.

# **Chapter 1: Introduction**

## 1.1 Purpose and objective

The most prevalent respiratory sleep disorder is obstructive sleep apnea (OSA), which is characterized by repetitive episodes of upper airway obstruction while asleep that leads to sleep fragmentation and hypoxemia. Common symptoms of this disorder include loud snoring, excessive daytime sleepiness, and frequent nocturnal awakening.<sup>2</sup> Patients with untreated OSA are at increased risk of car crashes, acute cardiovascular events, and impaired neurocognitive performance. <sup>3</sup> Similarly, cardiovascular (CV) co-morbidity is very common among OSA patients, with observational studies suggesting an independent relationship between OSA and the development of CV sequelae including hypertension, heart failure, cardiac arrhythmia, stroke, resting heart rate, and pulmonary artery pressure. 4 OSA occurs in both genders and all age groups and affects 10-20% of the population. OSA treatment can relieve a huge economic burden (billions of dollars per year).<sup>5</sup> Direct expense of OSA burden in Canada alone in 2008 was around 145 million dollars. <sup>5</sup> According to a recent study, the treatment of moderate to severe OSA is cost effective and depending on the patients' choice of therapy and the severity of their OSA, cost utility ratios can range anywhere from \$7420 to \$43889. <sup>6</sup> This finding matches the findings of other countries. <sup>7</sup> According to a recent study by Hillman and colleagues, the direct health costs of sleep disorders were \$160 million and \$1.08 billion for associated conditions in Australia alone. 8

The first description of continuous positive airway pressure (CPAP) as a potential treatment of OSA was published by Dr. Colin Sullivan in 1981. <sup>9</sup> Since then, CPAP has become the first-line therapy for OSA. CPAP is efficacious in reducing and in many cases eliminating snoring, hypoxemia, and the resultant arousals in patients with OSA, resulting in significant

improvements in health-related quality of life, alertness, and neurocognitive function. <sup>10</sup> CPAP is associated with reduced daytime and nighttime blood pressure increases in OSA patients, and patients with greater use of CPAP have more robust blood pressure reductions. 11 However, despite the high efficacy of this device, adherence rates range from 30-60% and of those that continue using it, 20-30% of them apply it insufficiently. <sup>10</sup> CPAP adherence is commonly defined as using the device for  $\geq 4$  hours for  $\geq 70\%$  of the nights monitored. <sup>12</sup> According to a recent study that only looked at sixty-six randomized control trial (RCT) or longitudinal cohort studies that were published between 1994-2015, the rate of non-adherence was 34.1%.  $^{12}$ However, this number may not be accurate and generalizable to the whole population as all the studies were RCT studies. This finding suggests that despite numerous interventions (i.e. educational, telemedicine, behavioural, etc.) that were designed to improve CPAP adherence over the long term, trends do not seem to show clinically impactful changes. According to different studies, various predictors such as socio-demographic characteristics, disease severity, psychosocial factors and side effects may affect how adherent OSA patients are with CPAP. 12 However, their results are inconsistent, which may be due to small sample sizes. Moreover, none of these studies have been done in Canada and data is thus lacking in a Canadian context. The overarching aim of this retrospective study was to identify the rate and predictors of CPAP adherence in Canadian OSA patients. The results of this study may eventually lead to the overall improvement of patient care and the patients' quality of life by improving their CPAP adherence. The main objective of this project was to review the medical charts and health surveys of patients that were referred to the University of British Columbia (UBC) Sleep Disorders Clinic between 2003-2009 and to determine the rate and predictors that may have affected CPAP adherence within this cohort among those who could adhere and not adhere to CPAP.

# 1.2 Methods

In this retrospective cohort study, patients who were referred to the UBC sleep clinic for possible OSA diagnostic were recruited between 2003-2009. At the time of recruitment, the severity of the patients' OSA as measured by apnea hypopnea index (AHI), which was based on their results of their polysomnography (PSG) test had to be  $\geq 5$  events/hours in order for each patient to be considered in the study. The patients that were included also had to complete a health survey. Furthermore, of those that were considered to have OSA (AHI  $\geq 5$  events/hours), patients were further divided into those that were prescribed CPAP and those that were not prescribed CPAP. Of those that were prescribed CPAP, patients' charts were later reviewed for CPAP adherence. After analysis was completed, CPAP adherence was defined as using the device for  $\geq 4$  hours for  $\geq 70\%$  of the time being used. According to this definition, patients were further divided into two groups including: 1) adhered to CPAP, 2) did not adhere to CPAP. We then looked at a broad group of different predictors that may be have been correlated with CPAP adherence of the recruited patients (Table 1.2). Multivariate logistic regression analysis was used to determine the relationship between numerous variables and CPAP adherence by using SAS 9.4.

Table 1.2. Potential predictors of CPAP adherence.

Patient characteristics	Age		
	Gender		
	Body mass index (BMI)		
	Education		
	Income		
	Menopausal status		
	Smoking status		
	Marital status		
	Bed partner status		
	Neck circumference		
	Alcohol intake (i.e. wine, beer, liquor)		
Sleep Related	OSA severity measured by apnea hypopnea index (AHI)		
•	Daytime sleepiness measured by Epworth Sleepiness Scale (ESS)		
	Number of car crashes		
	Restless leg syndrome		
	Periodic limb movement (PLM)		
	Sleeping aid medication intake		
	Snoring severity		
Cardiovascular related	Stroke status		
	Cardiac arrhythmia status		
	Myocardial infraction status		
Mental health related	Depression scale measure by patient health questionnaire (PHQ-9)		
	Mood disorder		
	Anxiety disorder		
	Antidepressant medication		
Overall health	Hypertension		
	Occupational injury		

Definition of abbreviations: AHI = apnea-hypopnea index; BMI = body mass index; ESS = Epworth Sleepiness Scale; PLM = periodic limb movement; PHQ-9 = patient health questionnaire;

#### 1.3 Significance

OSA is the most common respiratory disorder of sleep. However, the majority of the patients with OSA are undiagnosed. OSA is associated with many adverse health outcomes some of which include increased rate of motor vehicle crashes, hypertension, stroke, daytime sleepiness, reduced quality of life, and cardiovascular diseases. CPAP is the primary treatment for OSA and is effective in treating it. CPAP includes a mask on the nose which pushes air into the throat to support the upper airway from collapsing at night. Those who use CPAP have less daytime sleepiness, cardiovascular diseases, motor vehicle crashes, and a better quality of life. However, many of the patients that are diagnosed with this disorder do not adhere to CPAP. Various factors and predictors including bed partner, age, gender, race, and etc. may affect whether a patient will or will not adhere to CPAP. However, studies are inconsistent about the rate and predictors and the factors that may influence CPAP adherence mostly because they had a small sample size. More importantly, none of the studies have been done within a Canadian context. As a result, this study, which to the best of our knowledge is the largest study done in Canada, addresses the rate and predictors of CPAP adherence within a Canadian population. Also, by identifying the predictors of CPAP adherence and improving adherence to CPAP, patient care and the overall quality of life of these patients can be improved.

# **Chapter 2: Background and Literature Review**

There are a number of factors that may affect CPAP adherence (Table 2.0).

Table 2.0. Factors associated with CPAP adherence.

Socio-demographic	Age <sup>13, 14</sup>
characteristics	Gender 14, 15
	Race/Ethnicity <sup>13, 16</sup>
	Socioeconomic status (e.g., income, and education) <sup>17, 18</sup>
Disease severity	Severity of OSA <sup>19, 20, 21</sup>
	Severity of symptoms (e.g., daytime sleepiness) <sup>22</sup>
Psychosocial factors	Psychological variables <sup>23</sup>
	Social support <sup>24, 25</sup>
Side-effects	Dry nose <sup>26, 27, 28</sup>
	Dry mouth <sup>26, 27, 28</sup>
	Mouth leak <sup>28</sup>
	Mask leak <sup>23</sup>
	Nasal congestion <sup>29</sup>

# 2.1 Socio-demographic characteristics

Patient characteristics of relevance include age, gender, race and socioeconomic status. In general, the studies have been fairly inconsistent in associating these factors with adherence. For example, one study concluded that African American patients use the CPAP machine less often than Caucasians. This finding was supported by results from another study. <sup>16</sup> However, in a more recent study, sleep duration differed by race and those that slept shorter hours used CPAP for fewer hours and consequently, this may have skewed their results. <sup>30</sup>
Studies have been inconsistent with their results in regard to the correlation between gender and CPAP adherence. In a study by Budhiraja and colleagues, 57 Caucasian patients were compared to 39 African American patients who had electronic adherence information to a laboratory-based computer relayed by the CPAP machine through a telephone modem. <sup>13</sup> They concluded that only age (p = 0.018) and race were associated with CPAP adherence, more specifically, Caucasians had higher daily CPAP use compared to the African Americans (p = 0.005). <sup>13</sup> The

authors also concluded that gender and severity of OSA showed no association with CPAP adherence. <sup>13</sup> However, the number of patients in their study was not sufficient to draw firm conclusions. Another study that included 507 OSA patients concluded that women were less adherent to CPAP than men. 31 They also concluded that African American patients (i.e. both males and females) were 5.02 times more likely to be nonadherent to CPAP than Caucasians; however, this finding may be biased because there were only 18 Caucasians in this study compared to 196 African Americans. <sup>31</sup> In another study by Collen and colleagues, 400 patients were included after presenting for initial follow up between 4-6 weeks of CPAP therapy. Most of their patients were men (78%) with a mean age of 47±7.7 years.<sup>32</sup> Among this cohort, CPAP was used 78.1% of the time while asleep with an average of 3.13 hours/night. Only age ( $48.2 \pm 8.0$ years vs.  $45.8 \pm 7.3$  years; p=0.02) was associated with good adherence in this cohort and patients >50 years of age were more adherence with CPAP.<sup>32</sup> No other variable such as gender, degree of sleepiness or disease severity predicted adherence in their study. Another study concluded that age (18-79 years, P=0.156, n=605) and gender (P=0.553) were not associated with long term CPAP adherence. <sup>20</sup> However, in this study, many of their medical records were unattainable as they were misfiled, lost and moved.

Socio-economic factors such as income and education level may play a role in CPAP adherence. In one report of 162 OSA patients, patients who made >120% of the average income of the study sample had an increased odds of CPAP adherence (defined as  $\geq$  4 hours/night) compared to those who made less than 80% of average income. <sup>17</sup> Years of education among low, average and high-income level patients in this study were  $9.9 \pm 3.4$ ,  $13.1 \pm 3.0$ , and  $15.2 \pm 3.4$  respectively (p < 0.001). <sup>17</sup> Moreover, many patients cite cost as a reason for not adhering to CPAP therapy. For example, out of 265 OSA patients in one study, only 21% of the patients with low income accepted and purchased CPAP compared to 51.4% of those with average income and 75.6% of

those with higher income. <sup>18</sup> However, patients' income may vary depending on the country. Table 2.1 shows the findings of various studies regarding socio-demographic variables and whether or not they effect patients' adherence to CPAP. Top five studies with the largest sample size where chosen.

Table 2.1. Overview of socio-demographic characteristics variables influencing CPAP adherence.

Variables	Impact on CPAP adherence	Sample size (n)	Reference
Age (Increased)	Increase	16,783	Palm 2017 33
,		4,281	Woehrle 2011 34
		1,772	Means 2010 35
		1,155	Mcardle 1999 <sup>22</sup>
	No difference	1,141	Gagnadoux 2011 <sup>36</sup>
Gender (male)	Increase	4,281	Woehrle 2011 34
		16,783	Palm 2017 <sup>33</sup>
	No difference	1,141	Gagnadoux 2011 36
		750	Stuck 2012 <sup>37</sup>
		639	Kohler 2010 <sup>20</sup>
Race (African	Decrease	1,772	Means 2010 35
American, non-		507	Joo 2007 <sup>31</sup>
white, Maori, non-		403	Pamidi 2012 <sup>38</sup>
European)		214	Campbell 2012 <sup>39</sup>
	No difference	330	Platt 2009 40
Socio-economic	Increase	330	Platt 2009 40
status (higher socio-	No difference	1,141	Gagnadoux 2011 36
economic status)		265	Gulati 2017 <sup>18</sup>
(may include both		124	Wallace 2013 41
or only one of these		303	Galetke 2011 <sup>42</sup>
variables: income, education)			

*Definition of abbreviations*: CPAP = continuous positive airway pressure.

# 2.2 Disease severity

Disease severity, as measured by the apnea-hypopnea index (AHI) and nocturnal hypoxemia may improve CPAP adherence. In general, the more severe the disease, the higher the chance of adhering with CPAP therapy. However, the results are once again inconsistent. While many studies suggest that patients are more adherent to CPAP when their OSA is more severe, <sup>22, 43, 44,</sup> <sup>45</sup> others suggest no correlation between the two variables. <sup>46, 47</sup>A large study of 639 patients concluded that severity of OSA determines adherence to the CPAP therapy. <sup>20</sup> They further explained that the hypopneas recorded (HR) of 0.97 for oxygen desaturation index (ODI) indicated an increase of 10 in ODI, which is linked to a risk reduction of 26% for not adhering to CPAP. <sup>20</sup> They concluded that the rate of CPAP adherence was 81% and 70% after 5 years and 10 years respectively and that patients used it for 6.2 hours/night on average. <sup>20</sup> However, many of their medical records were missing due to misfiling or moving within the hospital, which can potentially affect the records of people adhering with CPAP therapy. On the other hand, another prospective study containing 112 newly diagnosed OSA patients concluded that no correlation existed between the severity of OSA and CPAP adherence. <sup>48</sup> The authors concluded that frequent apneas were associated with better adherence at 1 month after initiating CPAP (p=0.039) but not at 3 months (p=0.22). <sup>48</sup>

Excessive daytime sleepiness is one of the most important determinants of CPAP adherence and is usually measured by the Epworth Sleepiness Scale (ESS). The degree of excessive daytime sleepiness (EDS) along with the severity of OSA is associated with the long-term use of CPAP therapy. There have been considerable investigations on the effects of CPAP on EDS and many investigators have confirmed that EDS is reduced by continuous use of CPAP. <sup>49, 50, 51</sup> One small study of only 14 patients concluded that 85% of their patients who had severe OSA (AHI>80) and used a pressure of 12 cm H<sub>2</sub>O for a treated mean AHI of <10 showed significant

improvement in EDS.<sup>23</sup> However, they did not indicate how many of their patients required repeated titration to achieve the most effective CPAP level but the investigators did have a follow up with the patients after 1 week of CPAP use at home. Moreover, most of the results from the studies show a strong relationship between EDS reduction and CPAP adherence. In contrast to other studies, one large study with 639 patients concluded that EDS was not associated (P=0.185) with CPAP adherence and that other important determinants of long-term adherence with CPAP may affect the quality of life in OSA patients. <sup>20</sup> However, they had limited data on other daytime symptoms, which could potentially change their results regarding CPAP adherence. Table 2.2.1 and Table 2.2.2 show the findings of various studies regarding disease severity variables and whether or not they effect patients' adherence to CPAP. Top five studies with the largest sample size where chosen.

Table 2.2.1. Overview of severity of symptoms influencing CPAP adherence.

Variable(s)	Impact on CPAP adherence	Sample size (n)	Reference
Severity of	Increase	16,783	Palm 2017 <sup>33</sup>
symptoms daytime		1,155	Mcardle 1999 <sup>22</sup>
sleepiness)		740	Rosenthal 2000 52
(improved	No difference	1,141	Gagnadoux 2011 36
symptoms)		1,105	Budhiraja 2017 <sup>53</sup>

*Definition of abbreviations*: CPAP = continuous positive airway pressure.

Table 2.2.2. Overview of severity of OSA (AHI) influencing CPAP adherence.

Impact on	OSA (AHI) Sample	Male	Mean	Mean	Mean AHI	р-	Reference
CPAP	size (n)	(%)	age	BMI	(apneic	value	
adherence			(years)	$(kg/m^2)$	events/h)		
			(SD)	(SD)	(SD)		
Increase	16,783			Adherent			Palm 2017
		NA	58.6	NA	38.1 (22.1)	< 0.001	33
			(13.6)				-
				n-adherent			-
		NA	56.5	NA	30.3 (20.7)		
	1.000		(13.6)				
	1,889			Adherent	11.0 (10.0)	0.001	Marin 2012
		84	49.5	31.4 (3.9)	41.2 (19.9)	< 0.001	34
			(9.7)				-
				n-adherent	24.2 (42.4)		-
		82	49.9	30.8 (3.9)	31.3 (13.4)		
	1 155		(11.0)	A 33			3.6 11
	1,155			Adherent 20 (27	21 (10.52)	0.07	Mcardle
		86	50 (43-	30 (27-	31 (18-53)	0.07	1999 <sup>22</sup>
			58)	35)			-
				n-adherent	22 (12 52)		-
		69	44 (32-	30 (27-	22 (13-53)		
	1 1 / 1		57)	36)			C 1
	1,141	NI A		Adherent	46.4 (22.1)	0.0002	Gagnadoux 2011 <sup>36</sup>
		NA_	NA	32.9 (6.7)	46.4 (22.1)	0.0002	2011 30
		NT A		on-adherent	41.5 (21.2)		-
	926	NA	NA	32.1 (6.9)	41.5 (21.3)		Tamaniyalı
	826			Adherent	41.5 (10.1)	0.001	Tarasiuk 2012 <sup>55</sup>
		77	54.4	33.5 (5.4)	41.5 (19.1)	0.001	
			(9.9)	n adharar4			-
		81		on-adherent	26.1 (20.5)		-
		91	47.1 (10.1)	31.2 (5.3)	36.1 (20.5)		

*Definition of abbreviations*: AHI = apnea-hypopnea index; BMI = body mass index; CPAP = continuous positive airway pressure; SD = standard deviation; NA = not available.

Adherent patients: continuous positive airway pressure (CPAP) use ≥4hours/night. Non-adherent patients: CPAP use <4 hours/night.

## 2.3 Psychosocial factors

Psychological variables including health value, health locus of control, and self-efficacy were investigated by Wild and colleagues, using Wallston's model of social cognition. <sup>56</sup> The locus of control scale measures beliefs in regard to health along three dimensions including internality (patient's perception as to the extent to which their health is within their own control), chance (external control of their health outcome) and powerful others (belief that others have control). <sup>56</sup> Self-efficacy measures the person's belief in their own ability to accomplish a task. <sup>56</sup> The authors found that higher adherence was associated with higher internal locus of control and lower powerful others. This may be because those who believe they have control over their own circumstances and environment are more likely to internalise the advice that is given to them by their sleep specialist; they may then adhere more to CPAP despite its inconveniences. <sup>56</sup> Of note, these traits are modifiable and could be influenced by positive experiences. <sup>56</sup> Social factors including social support in general, and more specifically the involvement of a bed partner, are important factors associated with a patient's adherence to CPAP. If the idea to seek medical attention was not the patient's but of the bed partner's, adherence to CPAP is negatively influenced as the patient may not be ready to commit to therapy. <sup>57</sup> Of note, when patients use CPAP, bed partners often have improved subjective sleep quality, quality of life, alertness, mood and marital relationships. <sup>57</sup> Furthermore, sleeping with a bed partner who provides feedback on symptomatic improvements such as snoring may promote CPAP adherence. <sup>57</sup> How an individual manages and copes with anxiety and difficult situations in their life can also influence adherence to CPAP. <sup>58, 59</sup> Patients who are motivated to resolve their health problems tend to adhere better to CPAP compared to those who are less motivated and worry less about their health. <sup>58</sup> Moreover, self-efficacy in addition to social support and knowledge predict CPAP adherence during the first week of CPAP use and one month after starting CPAP. <sup>60</sup> There are

inconclusive findings in regard to the association between anxiety, depression and adherence. Some studies have found a relationship between these variables and adherence, <sup>35, 61, 62</sup> whereas others did not. <sup>63,64, 65</sup> Studies regarding depression and anxiety are generally small in sample size and are at risk of being underpowered. Table 2.3 shows the findings of various studies regarding psychosocial variables and whether or not they effect patients' adherence to CPAP. Top five studies with the largest sample size where chosen.

Table 2.3. Overview of psychosocial variables influencing CPAP adherence.

Variables	Impact on CPAP adherence	Sample size (n)	Reference
Psychological			
Self-efficacy	Increase	404	Philip 2018 66
(Increased)		288	Micoulaud-Franchi 2018 <sup>67</sup>
	No difference	126	Bakker 2011 <sup>68</sup>
		124	Wallace 2013 41
		119	Wild 2004 <sup>56</sup>
Anxiety (mental	Decrease	1,772	Means 2010 35
disorder, depression		242	Kjelsberga 2005 <sup>61</sup>
symptoms)		240	Law 2014 <sup>69</sup>
(presence of anxiety	No difference	1,141	Gagnadoux 2011 36
symptoms/disorder)		124	Wallace 2013 41
Social support			
Marital status	Decrease	1,141	Gagnadoux 2011 36
(living alone)	No difference	330	Platt 2009 40
,		124	Wallace 2013 41
		110	Kaplan 2007 70
		122	Poulet 2009 71

*Definition of abbreviations*: CPAP = continuous positive airway pressure.

#### 2.4 Side-effects

There have been many technological advances to make the use of the CPAP machine easier. These advancements evolved due to the patients' difficulty complying and using the device. Many patients refused to use CPAP because of its side effects, which include nasal congestion, mask discomfort, claustrophobia and dry nose and mouth. <sup>72, 73</sup> Two thirds of OSA patients will experience a side effect from using the CPAP device. <sup>65</sup> However, these side effects are not predictive of CPAP adherence, and those who have experienced one or all of these side effects have used CPAP more often than those who do not experience these side effects. <sup>74,75</sup> That is, patients who do not use CPAP obviously will not suffer the side effects.

CPAP has a variety of interfaces such as nasal, orinasal and oral, which include nasal mask, nasal pillow, face mask and oral mask. <sup>72</sup> It is critical for the patients to choose the optimal interface because it will influence their acceptance of CPAP therapy and their long-term adherence with the device. There is not a universally accepted mask interface that will fit every patient and as a result, a properly fitted and comfortable mask interface is crucial when it comes to adhering with CPAP. <sup>76</sup> Using the mask for some time often irritates the skin in a particular spot or causes a feeling of claustrophobia. <sup>77</sup> In a retrospective study, those who used CPAP less often felt claustrophobic more frequently than those who used CPAP on a regular basis. <sup>27</sup> Similarly patients who use CPAP on a regular basis experience mask side effects and discomfort more often compared to those who do not use their device as often. <sup>27</sup> Nasal congestion is a very common side effect when using CPAP and as much as 65% of the patients using the device report nasal congestion, dry nose or throat and discomfort. <sup>78</sup> Patients that encounter nasal congestion are typically mouth breathers and consequently while asleep, the temporomandibular joint retracts and further reduces the retroglossal airway. <sup>9</sup>

The early CPAP machines included a mask attachment using silastic glue and/or rudimentary

straps. Over time, there have been significant improvements in mask technology with a variety of interfaces available; however, it is unclear whether changes in mask type improves adherence to CPAP or if these reflect a marketing strategy. There is a lack of research that compares these interfaces. One of the most common side-effects of using CPAP is mask leaks, which results in the patient not receiving the proper air pressure to treat their sleep apnea and consequently reduces CPAP effectiveness. <sup>79</sup> Mask leak can cause discomfort and dryness in the nasal passage. <sup>79</sup> However, its measurement and reporting terminology are inconsistent since CPAP downloads do not typically report the amount of leak (i.e. average leak or large leak) and also do not report total leak minus expected leak or even artifact-free leak. <sup>79</sup>

Furthermore, patients that sleep with their mouth open have a higher chance of upper air-way collapsibility. <sup>80</sup> Mouth leak occurs in 10-15% of patients <sup>80</sup> and is undesirable for many different reasons including the fact that they reduce the effect of pressure which drives air into the lungs or splint open the upper airway. <sup>28</sup> In order for this treatment to be effective, higher pressure must be applied at the nose mask to compensate for the leak. Moreover, mouth leaks may lead to drying of the mucosal surfaces in the presence of a one-way flow (i.e. air enters through the nose and passes out of the mouth). <sup>28</sup> This condition may then lead to nasal obstruction (i.e. dry and stuffy nose and mouth) which may be intolerable for many patients. <sup>28</sup> As a result, those that are mouth breathers have low adherence to CPAP therapy. <sup>80</sup> Also, mouth leak may compromise CPAP therapy and the first period of CPAP use does in fact predict long-term adherence. <sup>81</sup> Humidification is another technological advancement designed to reduce CPAP side effects such as dry mouth, throat, or nose. <sup>78</sup> Two types of humidity are commonly used: cold passover humidity and heated humidity. <sup>78</sup> Heated humidity is used more often as it improves the patient's comfort by preventing the increase in mucosal blood flux, which means that the drying effect of the airflow is responsible for this change. <sup>78</sup> However, it is unclear whether heated humidity

increases CPAP adherence. In a couple of studies, there was no difference in side effects between patients who used heated humidification and those who did not.<sup>82, 83</sup> In contrast, another study found that patients exposed to heated humidity compared to no humidity adhered more to CPAP (5.52 hrs/night vs. 4.93 hrs/night; p = 0.008). <sup>84</sup> This particular study noted that heated humidity reduced dry mouth, throat, and nose and increased refreshed feeling upon awakening. <sup>84</sup> Table 2.4 shows the findings of various studies regarding side-effect variables and whether or not they effect patients' adherence to CPAP. Top five studies with the largest sample size where chosen.

Table 2.4. Overview of side-effect variables influencing CPAP adherence.

Variables	Impact on CPAP adherence	Sample size (n)	Reference
Adverse effects (may include one or	Decrease	2,311	Borel 2013 85
all of mask leak, nasal congestion,		385	Kreivi 2010 86
dry nose and mouth) (presence of		300	Kalan 1999 87
side-effects)		204	Engleman 1996 45
	No difference	193	Pepin 1995 88

*Definition of abbreviations*: CPAP = continuous positive airway pressure.

#### 2.5 Interventions to improve CPAP adherence

Although a wealth of research demonstrating the efficacy of many of these interventions exist, they can be difficult to implement clinically due to issues around feasibility and cost. Some of the most effective interventions regarding CPAP adherence include supportive, educational, telemedicine, cognitive behavioural, and technological (Table 2.5). Supportive interventions are described as reinforcement, support and/or enhanced access to sleep specific and healthcare resources. 82 Educational interventions mainly focus on patient education and highlight the potential benefits of using CPAP, function and care of the device, and ways to eliminate the side effects and problems that patients may experience. <sup>12</sup> Telemedicine is defined as "the use of information and communication technology to deliver health services, expertise and information over distance, geographic, time, social and cultural barriers". 89 Moreover, telemedicine interventions have been used to reinforce CPAP adherence by sending digital images via email, fax machines, telephone and videoconferencing. 90 Cognitive behavioural interventions are simply described as how the patients make their decisions. This intervention focuses on enhancing patients' new learning experiences and work on correcting their irrational beliefs through exposure to positive stimuli. 82 Finally, technological interventions include the improvements of the CPAP machine and its mask (i.e. nasal cushions (sealing around the outside of the nose), nasal pillows (sealing around the base of each nostril), oronasal (also known as a 'full face mask', sealing around the outside of the nose and either below the lower lip or under the chin) and oral (used rarely)). 12

Table 2.5. Interventional studies to improve CPAP adherence.

<b>Supportive Interventions</b>					
Study	Design	Sample size (n)	Intervention	CPAP adher ence metric	CPAP adherence outcome change (Y=yes; N=No)
Stepnowsk y 2007 <sup>91</sup>	RCT	45 (Experime ntal group=20; Control group=20)	Patients in the experimental group were assigned to telemonitored clinical care (i.e. joint decision making and partnership between provider and patient). These patients had objective flow generator data monitored as frequently as needed per specified clinical pathways throughout the active 2-month treatment period compared to the control group that had their CPAP adherence data downloaded at 1mo.	Mask- on time	Y at 2mo; Telemonitored group used CPAP for 4.1 hours compared to 2.8 hours in the control group (p=0.07).
Smith 2006 59	RCT	19 (Experime ntal group=10; Control group=9)	Telehealth (i.e. telephone delivered) intervention. The experimental group had a conversation in their home and with their research nurse for a period of 12 weeks. The phone calls were to reinforce nightly, use of CPAP by emphasizing the improvement of general health and how to manage barriers of CPAP use and its related problems. The control group received phone calls regarding the importance of using vitamins every day. Both groups received	Machi ne-on time	Y at 12wk; 9 out of 10 in the experimental group compared to 4 out of 9 in the control group adhered to CPAP after the intervention (p=0.03).

			telehealth contact 3X during the first week of CPAP initiation and weekly for the remaining 11 weeks of the study.		
Educational	Interventi	ons			
Study	Design	Sample	Intervention	CPAP adher ence metric	CPAP adherence outcome change (Y=yes; N=No)
Wiese 2005 92	RCT	100 (Experime ntal group=51; Control group=49)	During the first visit, both groups received explanations about the importance of using CPAP. The experimental group also received a 15-minute videotape about what CPAP was, what CPAP felt like on a sensory basis, and how it improved the patients' conditions.	Undefine d	Y; Those in the experimental group had a return rate of 72.9% compared to 48.9% (p=0.017) at 1month follow-up.
Golay 2006 93	One group pre-test; post test	35	Patients were enrolled in an educational program including what CPAP serves for, workshops, hospital CPAP titration sleep study, Ear Nose and Throat consultation, round table with the spouses, and treatment plan discussion groups.	Machi ne-on time	Y; Baseline CPAP readings averaged 4.4 h while the readings raised to 5.1 h after 3 months (no statistical significance was reported).
Telemedicin	e Intervent	ions			
Study	Design	Sample	Intervention	CPAP adher ence metric	CPAP adherence outcome change (Y=yes; N=No)
<b>Sparrow 2010</b> <sup>94</sup>	RCT	250 (telephone -linked communic ations for CPAP (TLC- CPAP)	The experimental group received TLC-CPAP (i.e. motivational enhancement therapy for CPAP). A Sleep Symptoms Checklist, the Functional Outcomes of Sleep Questionnaire, the	Mask- on time	Y at 6mo and 12mo; CPAP use was higher in the intervention group compared to the control group at 6mo (1hr/night higher;

		=124; Placebo control=12 6)	Center for Epidemiological Studies Depression Scale and the Psychomotor Vigilance Task were administered at study entry and at 6-mo and 12-mo follow-up.		p=0.006) and 12mo (2hr/night higher; p=0.004).
Hwang 2018 95	Four arms, randomiz ed, factorial design clinical trial	1,455 (home sleep apnea testing=95 6; Prescribed CPAP=55 6)	Two telemedicine interventions were implemented with primary endpoint of 90-day of CPAP usage: 1) web-based OSA education (Tel-Ed) and 2) CPAP telemonitoring with automated patient feedback (Tel-TM). Patients were randomized to 1) usual care, 2) Tel-Ed added, 3) Tel-TM added, or 4) Tel-Ed and Tel-TM added (Tel-both).	Mask- on time	Y; CPAP average daily use at 90 days was 3.8, 4.0, 4.4, and 4.8 hours in usual care, Tel-Ed, Tel-TM, and Tel-both groups. Usage was significantly higher in the Tel-TM and Tel-both groups versus usual care $(P = 0.0002 \text{ for both})$ but not for Tel-Ed $(P = 0.10)$ .
Cognitive I	Behavioural <b>T</b>	Therapy Inte	rventions		

Study	Design	Sample	Intervention	CPAP adher ence metric	CPAP adherence outcome change (Y=yes; N=No)
<b>Richards 2007</b> <sup>96</sup>	RCT	100 (Interventi on group=50; Control group=50)	The intervention group received Two 1-hour sessions along with their partners regarding distorted beliefs and promoting a positive outlook for CPAP.	Mask- on time	Y at both 7-days (p<0.0001) and 28-days (p<0.0001); A Higher adherence to CPAP therapy was found in the intervention group (2.9 hours difference).
<b>Aloia 2007</b> 97	RCT	142 (Motivatio nal enhancem ent therapy group=54;	Motivational enhancement therapy group and traditional education group received two, 45-min, face-to-face therapy sessions	Mask- on time	N at 13 weeks; The three groups' adherence did not differ $(p > 0.05)$ . 61% of the individuals in the control group,

		Traditiona 1 education =47; Control group=41)	beginning after 1 week with CPAP treatment compared to the control group, which only received brochures regarding OSA and the importance of CPAP adherence.		68% of those in the traditional education group and 67% of those in the motivational enhancement group adhered to CPAP.
Technologic	cal Interven	tions			
Study	Design	Sample	Intervention	CPAP adher ence metric	CPAP adherence outcome change (Y=yes; N=No)
<b>Malhotra 2018</b> <sup>98</sup>	Retrospe ctive study	128,037 (active patient engageme nt tool (APE)=42, 679; usual care monitorin g (UCM)=8 5,358)	AirView, which was used in the UCM group, is a cloud-based technology. CPAP data were transferred automatically to this cloud daily and clinicians managed adherence and therapy for patients with OSA. MyAir, which was used in the APE group, provided real-time feedback and coaching to patients based on their data within AirView. Activation date in the APE group was within 7 days of therapy initiation. Patients in both groups were followed for 10months.	Mask- on time	Y; The addition of APE increased CPAP adherence (5.9hr; p=<0.0001) compared to those with UCM (4.9hr).
<b>Mador 2005</b> <sup>99</sup>	RCT	98 (treatment group=49, control group=49)	Patients received heated humidification at CPAP initiation in the treatment group. Patients were followed for 1months, 3months and 12 months.	Mask- on time	N; heated humidification did not improve CPAP adherence with 4.3 hr/night at 1month, 4.8 hr/night at 3 months, and 4.8 hr/night at 12 months in the control group and

4.3 hr/night at 1
month and 3
months, and 4.2
hr/night at 12
months in the
treatment group.

Definition of abbreviations: RCT = randomized control trial; CPAP = continuous positive airway pressure; mo=month; TLC= telephone-linked communications; OSA=obstructive sleep apnea; Tel-Ed = web-based OSA education; Tel-TM = CPAP telemonitoring with automated patient feedback; APE = active patient engagement tool; UCM = usual care monitoring; CBT = cognitive behavioural therapy.

#### 2.6 Rationale

OSA is a common sleep disorder and it is commonly undiagnosed. Symptoms of this disorder include loud snoring, excessive daytime sleepiness, frequent nocturnal awakening, neurocognitive impairments, increased rate of car crashes and cardiovascular events including heart attacks, strokes, and hypertension. 100 Patients with untreated OSA are at increased risk of car crashes, acute cardiovascular events, and impaired neurocognitive performance. <sup>101</sup> OSA occurs in both genders and all age groups and affects 10-20% of the population. OSA treatment can cause a huge economic burden in Canada. Adherence rates to CPAP, which is the most effective treatment for OSA, are low (30%-60%) and over the past twenty years or so, the adherence rates have not increased despite numerous interventions (i.e. patient education, supportive, technological, telemedicine and cognitive behavioral), which are aimed to improve adherence. <sup>10</sup> CPAP adherence is affected by various factors including socio-demographic characteristics (i.e. age, <sup>13-14</sup> gender, <sup>14-15</sup> race, <sup>13, 16</sup> and socio-economic status <sup>17-18</sup>), disease severity (i.e. severity of OSA <sup>19-21</sup> and its symptoms <sup>22</sup>), psychosocial factors (i.e. psychological variables <sup>23</sup> and social support <sup>24-25</sup>), side-effects (i.e. dry nose <sup>26-28</sup>, dry mouth <sup>26-28</sup>, mouth leak <sup>28</sup>, mask leak <sup>28</sup>, and nasal congestion <sup>29</sup>) and is a complex behavioral problem. However, to the best of our knowledge, none of the studies were done within a Canadian context. As a result, the aim of this thesis was to identify the rate and predictors of CPAP adherence in Canadian OSA patients.

#### 2.7 Hypothesis and objectives

# 2.7.1 Objective

The objective was to assess adherence rates with CPAP in patients with OSA. Various factors including age, gender, apnea hypopnea index (AHI), Epworth sleepiness scale (ESS), body mass index (BMI), patient health questionnaire (PHQ-9), education, income, menopausal status, rate of car crashes, smoking status, marital status, bed partner status, snoring severity, occupational injury, restless leg syndrome, neck circumference, mood disorder, anxiety disorder, alcohol intake, stroke status, cardiac arrhythmia status, myocardial infraction status, periodic limb movement (PLM), hypertension, sleeping aid medication intake, and antidepressant medication were determined.

# 2.7.2 Hypothesis

We hypothesized that predictors including age, gender, apnea hypopnea index (AHI), Epworth sleepiness scale (ESS), body mass index (BMI), depression, education, income, menopausal status, rate of car crashes, smoking status, marital status, bed partner status, snoring severity, occupational injury, restless leg syndrome, neck circumference, mood disorder, anxiety disorder, alcohol intake (i.e. wine, beer, liquor), stroke status, cardiac arrhythmia status, myocardial infraction status, periodic limb movement (PLM), hypertension, sleeping aid medication intake, and antidepressant medication correlate with CPAP adherence; however, given the inconsistency within the literature, it is likely that some of these factors will not be predictors of CPAP adherence.

# **Chapter 3: Methods**

# 3.1 Overall study design

Patients that were referred to UBC sleep clinic between 2003-2009 for a possible OSA diagnosis were recruited (Figure 3.1). Patients were divided into those that had done an overnight PSG test and those that had not done an overnight PSG test. Of these patients, only those with an AHI  $\geq$  5 apneic events/hour were included in the study. Furthermore, patients with an AHI  $\geq$  5 apneic events/hour were divided into those that were prescribed CPAP and those that were not prescribed CPAP. Moreover, patients that were prescribed CPAP were further divided into those that adhered to CPAP and those that did not adhere to CPAP. CPAP adherence was in accordance to other studies and was defined as using the device for  $\geq$  4 hours for  $\geq$ 70% of the nights monitored. Lastly, in order to find the factors that could potentially predict CPAP adherence, we looked into those that adhered to CPAP.




Figure 3.1. Overall study design. As shown in the figure, patients were recruited from UBC sleep clinic from 2003-2009. These patients were further categorized into those that had done a PSG test and those that had not done a PSG test. Of those that had done a PSG test, only patients with an AHI of  $\geq$  5 apneic events/hour were included. Of the patients with an AHI of  $\geq$  5 apneic events/hour, patients were further categorized into those that were prescribed CPAP and those that were not prescribed CPAP. In order to find adherence in this cohort, patients that were prescribed CPAP were once more divided into those that were to CPAP and those that did not

adhere to CPAP. Lastly, in order to find the predictors of CPAP adherence, adherent patients were considered.

#### 3.2 Patient recruitment

The protocol of this single center study was approved by the Clinical Research Ethics Board (H13-00346) at the University of British Columbia. Patients who were referred to UBC sleep clinic for a possible OSA diagnosis were recruited from 2003-2009. Moreover, PSG is recommended by both the American Thoracic Society and the American Academy of Sleep Medicine and was performed by using conventional instrumentation and analysis according to the recommendations on the definition of syndrome and measurement techniques, which was published by the American Academy of Sleep Medicine. <sup>102</sup> The overnight PSG test was used to objectively identify OSA and its severity according to AHI. Consenting patients were also given an extensive questionnaire (Appendix A) regarding their demographics, sleep habits, and subjective sleepiness as measured by Epworth Sleepiness Scale (ESS), which needed to be completed before the patients left the clinic the next day. For this project, patients with AHI of ≥ 5 events/hr as assessed by their overnight PSG were considered to have OSA.

#### 3.3 Inclusion criteria

Study subjects were recruited at the University of British Columbia's sleep clinic between 2003-2009 in accordance to the ethics approval from UBC's research ethics board. They had to be referred to UBC sleep clinic for a suspected OSA diagnosis. The inclusion criteria included: 1) being able to read and understand English, 2) being able to provide informed consent, 3) not used CPAP within the past 3 months prior to their PSG test, and 4) being able to fill out the health survey. Patients referred to the UBC sleep clinic for a PSG by sleep psychiatrists instead of respiratory physician were excluded from the study. This was because these patients mainly have non-respiratory issues such as insomnia, depression or narcolepsy and as a result, as such, they

have other disorders leading to EDS. Lastly, those with serious medical conditions including mental disability or dementia, active psychiatric disease and those unable to provide an informed consent form were also excluded from the study.

## 3.4 Risks and confidentiality

The health survey and the procedures that were used in this project had no physical risks to the patients. Initially, when patients arrived at the sleep clinic to see a sleep specialist regarding their condition, sleep specialists or a research coordinator explained the study to potential subjects and gave them consent forms. Subjects were also told they could withdraw from the study anytime they wished and could have their data removed at any point during the study. The original confidential databases and datasheets were stored in a locked cabinet in an office at Vancouver General Hospital (VGH).

# 3.4 Polysomnography

Polysomnography is the main procedure used in clinical practice in order to diagnose and quantify OSA severity. Different sleep stages were documented by using electroencephalographic (EEG), electrooculographic (EOG) and electromyographic (EMG) criteria. Airflow was measured by recording the nasal pressure. Moreover, snoring was measured by using a microphone (Model SL120, Pacer Industries, Toronto, Canada), which was attached to the respitrace band at the level of third intercostal space. An electrocardiographic (ECG) was monitored to detect cardiac arrhythmias. Arterial oxygen saturation (SaO2) was monitored with a pulse oximeter (Model N-100, Nellcor Inc., Hayward, CA), which was attached to the index figure. Moreover, chest wall movement was monitored using a respiratory plethysmograph (Respitrace, Ambulatory Monitoring Equipment, Ardsley, NY). The entire record was manually

scored for sleep stage, type of apnea and the duration. The recorded data was eventually merged with the remainder of the data gathered from the questionnaires.

Hypopnea was associated with a reduction in flow with either a 3% or more drop in arterial saturation or an arousal. Similarly, apnea was defined as a pause in respiration for more than 10 seconds. <sup>103</sup> The severity of OSA was defined in terms of the number of apnea or hypopnea events per hour while asleep and is measured by the apnea hypopnea index (AHI). Patients were further divided into two groups depending on their PSG results. If they had an AHI of 5 events/hour or more, they were considered to have OSA and were put into the "yes" group. Those with an AHI of less than 5 were considered to not have OSA and were put into "no OSA" group. The severity of OSA was further categorized into no OSA (<5), mild OSA (5-14), moderate OSA (15-29), severe OSA (≥30).

#### 3.5 Obtaining patient charts and the health questionnaire

Initially, the health surveys were self-completed when the patients were at the sleep laboratory. The questions in the survey (Appendix A) mostly concentrated on demographic variables and the events the patients experienced within two years prior to their PSG. These variables included self-reported occupational injury, occupational history, sleep habits, witnessed snoring and apneas, past medical history, medication intake, health habits (i.e. alcohol, smoking, caffeine intake), type of industry work, height, weight and lastly, subjective daytime sleepiness as measured by ESS. Body mass index (BMI) was calculated from using the patient's height and weight.

Patients' medical history were collected from Iron Mountain Inc. from 2016 to 2018. The charts were then extensively reviewed and when possible, both subjective and objective CPAP adherence rates were extracted. Objective adherence was obtained from the charts by looking at patients' CPAP's downloaded data. Moreover, subjective data was obtained by looking at the

sleep physician's notes from the patient's last visit and was further categorized into either adhering or not-adhering depending on the doctors' notes. However, at the time of patient recruitment, most of the patient files lacked objective data because at the time the data was collected, objective measurements of CPAP treatment were less ubiquitous than they are today, as a result, the adherence data in this study was taken from both subjective data and objective data that was retrieved from the patient charts at the UBC sleep clinic. Also, when objective data was lacking, subjective data was used on its own. Objective measures were used for threshold. However, about 60% of the collected data was in the form of subjective data as we were missing a large number of objective data. For the objective data, we used the most commonly defined definition of CPAP adherence (i.e. using the device for  $\geq 4$  hours for  $\geq 70\%$  of the nights monitored), however, for the subjective data, we used the physician's notes from the patient's last visit to that doctor. The doctor's notes commonly identified the patients as either adherent or not adherent, however, when the notes were vague in terms of whether a patient was adherent or not adherent to CPAP, we used our judgement to categorise those patients as either adherent or non-adherent. Two experts reviewed the spreadsheet to determine whether or not patients were adherent with the therapy.

#### 3.6 Data sets and statistical analyses

Completed health surveys were either scanned or manually entered to the patients' database. The only people with access to this database were the members of the research team (i.e. researchers and support personnel). We further maintained the patients' confidentiality by removing the identifying information including names and personal health numbers from the main dataset. Doing so allowed us to identify the patients by a code number and maintain their confidentiality. Patients with an AHI of  $\geq$  5 events/hour were divided into "prescribed CPAP" or "not prescribed CPAP" depending on their extracted subjective and/or objective data. Patients that were

prescribed CPAP were then divided into "adhered to CPAP" or "did not adhere to CPAP" based on the defined definition of CPAP adherence. There is inconsistency in how CPAP adherence is defined, both empirically and clinically. However, the most common definition of CPAP adherence is to use the device for  $\geq 4$  hours for  $\geq 70\%$  of the nights monitored. Early in the 1990s, papers reported CPAP was used for 4.7 hours/night on average. However, the authors did not indicate this number to be an adequate amount of CPAP use, a common assumption then emerged wherein use of CPAP for 4 hours per night on 70% of nights was adequate to reverse the negative effects of OSA and as a result, it was established as a clinical and empiric benchmark of CPAP adherence. Furthermore, our definition of CPAP adherence was in accordance to other studies and was defined as using the device for  $\geq 4$  hours for  $\geq 70\%$  of the nights monitored. CPAP adherence was measured at one point in time (i.e. when the patients went back for a follow-up with their respiratory physician). To begin the analysis, the databases that contained the health survey results and the PSG data were merged. In order to determine the predictors of CPAP adherence in those that were prescribed CPAP, baseline characteristics including age, daytime sleepiness, OSA severity, body mass index (BMI), depression measured by patient health questionnaire (PHQ-9), menopausal status, antidepressant medication, smoking status, restless leg syndrome, marital status, bed partner status, snoring severity, occupational injury, neck circumference, mood disorder, anxiety disorder, periodic limb movement (PLM), hypertension, myocardial infraction, cardiac arrhythmia, stroke, income level, alcohol consumption, education, motor vehicle accident rate, gender, and sleep aid medication were summarized as mean values of proportions as appropriate. For this analysis, education was used as a marker for socioeconomic status because many patients were retired with a low income, but their accumulated asset may have increased their socioeconomic status. Also, higher education is associated with a better health status because those with a higher education tend to understand

the seriousness of OSA and the need to adhere to CPAP better and as a result, adhere more to CPAP compared to those that are not as educated. As a result, using education as a marker of socioeconomic status helped us to determine if people with higher education will be more adherent to CPAP in this study.

We had a relatively large sample size and we wanted to determine which factors, including continuous and categorical variables, would predict CPAP adherence. As a result, a two-sample t-test was performed for continuous variables and  $\chi^2$  test was used for non-continuous or categorical variables between the adherent and the non-adherent groups. Group differences were also tested by using  $\chi^2$  test. Logistic regression was used to model the odds of using CPAP in those that were prescribed CPAP and those that were not prescribed CPAP. A *P* value of < 0.05 was considered to be statistically significant. Lastly, statistical analysis was performed with SAS software, version 9.4 (SAS Institute, Inc., Cary. NC).


# 3.7 Potential predictors of CPAP adherence

Various CPAP adherence predictors and determinants were evaluated within this cohort in this research (Table 1.2).

# **Chapter 4: Results**

# 4.1 The study population

2,281 patients were enrolled in the original study. We were able to identify 1,250 patient charts for analysis. These patients were all recruited at UBC sleep clinic, underwent PSG, filled out study questionnaire and had available clinic follow-up data within the reviewed charts. Of these 1,250 patients, 844 were prescribed CPAP by a sleep physician and 406 were not prescribed CPAP (Figure 4.1). Since we wanted to investigate the predictors and determinants of CPAP adherence, we focused on patients that were prescribed CPAP. Moreover, out of those that were prescribed CPAP, 421 (49.9%) adhered to CPAP and 423 (50.1%) did not adhere to CPAP.



**Figure 4.1.** Cohort sample size. Overall, 1,250 patients were identified, of which 844 patients were prescribed CPAP and 406 were not prescribed CPAP. Out of those that were prescribed CPAP, 421 patients adhered to CPAP and 423 patients did not adhere to CPAP.

# 4.2 The correlation between different predictors of CPAP adherence in those that adhered and did not adhere to CPAP

Multivariate analysis (Table 4.2.1 and Table 4.2.2) indicated that CPAP adherence was only associated with one independent variable, which was the severity of OSA as measured by AHI (p=<0.0001).

Table 4.2.1. The correlation between continuous variables and CPAP adherence in adherent and non-adherent group using t-test.

Variables	t-test P	Adherent patients	Non-adherent patients	Standard deviation (Adherent)	Standard deviation (Non- adherent)
AHI (apneic events/hour)	< 0.0001	34	27	25	21
Age (years)	0.80	51	51	11	11
ESS (points)	0.63	11	11	4.6	4.4
BMI (kg/m <sup>2</sup> )	0.38	33	32	6.8	7.2
Depression-PHQ-9 (points)	0.68	16	16	5.6	5.7
Menopausal status (years)	0.78	3.6	3.4	6.0	4.5

*Definition of abbreviations*: AHI = apnea-hypopnea index; BMI = body mass index; ESS = Epworth Sleepiness Scale; PHQ-9=patient health questionnaire.

<sup>\*</sup>Results presented as mean unless otherwise indicated. Significant level for p-value: <0.05.

Table 4.2.2. Frequency of categorical variables by adherent and non-adherent using Chisquared test.

Variables	Chi-Squared P	Adherent patients (%)	Non-adherent patients (%)
Antidepressant medication (< 1 night/week)	0.57	95	96
Gender (males)	0.97	73	73
Smoking status (current smokers)	0.91	16	16
Restless leg syndrome	0.76	4.2	4.6
Marital status (living with spouse)	0.21	67	71
Bed partner status (living with a partner)	0.20	70	74
Snoring severity (≥ once/week)	0.52	95	94
Occupational injury	0.16	9.5	13
Neck circumference (17"≥)	0.97	74	74
Mood disorder	0.17	22	18
Anxiety disorder	0.84	7.1	7.5
Periodic limb movement	0.44	0.6	1.2
Hypertension	0.05	25	19
Myocardial infraction	0.78	4.2	3.8
Cardiac arrhythmia	0.75	6.6	7.2
Stroke	0.29	1.8	0.87
Income level (>\$40,000)	0.19	70	75
Alcohol consumption (1 drink/day ≥)	0.13	93	90
Education (≥ college degree)	0.15	63	57
Car crashes	0.96	16	16
Sleep aid medication (< 1 night/week)	0.30	88	85

<sup>\*</sup>Results presented as frequency unless otherwise indicated. Significant level for p-value: <0.05.

#### 4.2.1 AHI and adherence


There was a strong association between AHI and CPAP adherence within our cohort. In fact, AHI was the only predictor significantly associated with CPAP adherence. After performing a two-sample t-test, we found the p-value between the two variables to be <0.0001 with average mean in the adherers and non-adherers to be  $34 \pm 25$  apneic events/hour and  $27 \pm 21$  apneic events/hour respectively (Table 4.2.1). We also categorized patients into mild OSA (5-14), moderate OSA (15-29), severe OSA ( $\geq$ 30) (Figure 4.2.1). <sup>104</sup> The correlation was tested once more and a p-value of <0.0001 was observed, which suggests that there is a correlation between OSA severity categories and CPAP adherence (Table 4.2.3). In figure 4.2.1, we can see the overall pattern of adherence increases as the severity of OSA increases. 57% of those with an AHI  $\geq$ 30 events/hour were adherent to CPAP compared to 50% of those with AHI  $\geq$ 15, but <30 and 32% of those with AHI  $\geq$ 5, but <15.

Table 4.2.3. OSA severity categories and CPAP adherence.

AHI (apneic events/hour)	Adherent	Non-adherent	
Mild: ≥5, but <15	16%	34%	
Moderate: $\geq$ 15, but $\leq$ 30	28%	28%	
Total AHI <30	57%	68%	
Severe: ≥30	43%	32%	

*Definition of abbreviations*: AHI = apnea/hypopnea index; CPAP = continuous positive airway pressure; OSA = obstructive sleep apnea.

Adherent patients: continuous positive airway pressure (CPAP) use ≥4hours/night for 70%. Non-adherent patients: CPAP use <4 hours/night.



**Figure 4.2.1. Association between AHI and CPAP adherence.** As shown in the figure, the more severe the OSA, the more patients adhered to CPAP.

# 4.2.2 Age and adherence

The age of the patients was obtained from the patients' medical history charts and their age at the time of their PSG test was recorded. However, similar to many other studies, we found no association between age in those that adhered and did not adhere to CPAP (p=0.60) (Table 4.2.1). The mean age of those who adhered to CPAP was  $51 \pm 11$  years, which was the same as the mean age of those who did not adhere to CPAP ( $51 \pm 11$ ).

#### 4.2.3 ESS and adherence

ESS, which is a validated self-administered questionnaire, was used to measure sleepiness in this population. Mean ESS scores are shown in Table 4.2.1. The mean ESS for subjects that adhered to CPAP was  $11 \pm 4.6$  points, which was similar to those that did not adhere to CPAP  $11 \pm 4.4$  points. Having an ESS of 11 points means that the patients had mild EDS. There was no significant difference in ESS mean among those that adhered and those that did not adhere

(p=0.63). one of the reasons as to why ESS was not one of the predictors of CPAP adherence could be due to the fact that sleepiness was self-reported so the results may not have been accurate whereas, for AHI, we had the actual results from the PSG test, and it was more accurate. The other reason could be because sleepiness may have been caused from other factors other than OSA such as PLM or the patients may have been tired for other reasons. As a result, they were put on CPAP, so their sleepiness improved. One downside is that we do not know how much their sleepiness improved on CPAP.

#### 4.2.4 BMI and adherence

Subjects' BMI was calculated from reviewing their medical charts and obtaining their weight and height. However, after analysing the data, we found no significant association between BMI and CPAP adherence (p=0.38) (Table 4.2.1). Those that adhered to CPAP had a BMI of 33 ± 6.8 kg/m², whereas those that did not adhere had a BMI of 32 ± 7.28 kg/m². Furthermore, similar to previously mentioned variables, the BMI was categorised into normal (18.5-24.9), overweight (25-29.9), obese I (30-34.9), obese II (35-39.9) and obese III (>40) (Table 4.2.4). There was no association between different BMI categories and CPAP adherence (p=0.20). Surprisingly, those who were overweight (BMI=25-29.9) were much less adherent to CPAP compared to the other BMI categories. However, the results were again insignificant. The cause of OSA in those that have normal BMI or are underweight is multifactorial, but it is mainly due to having enlarged large tonsils. Those with a larger tonsil tend to have greater resistance in their upper airway, which leads to its collapse. <sup>105</sup> Also, abnormal anatomic structure of the nose, in particular septal deviation, may cause symptoms of OSA due to the reduction of nasal airflow and elevated nasal airway resistance. <sup>106</sup>

Table 4.2.4. BMI categories and CPAP adherence.

BMI category (kg/m²)	Adherent	Non-adherent	
Normal weight:18.5-24.9	18%	14%	
Over weight: 25.0-29.9	27%	33%	
Obese I: 30.0-34.9	26%	26%	
Obese II: 35-39.9	15%	15%	
Obese III: >40	14%	12%	

*Definition of abbreviations*: BMI = body mass index; CPAP = continuous positive airway pressure.

# 4.2.5 Depression and adherence

PHQ-9 is a validated questionnaire-based depression scale, which is commonly used to assist primary care practitioners to diagnose depression.  $^{107}$  Subjects' PHQ-9 scores were obtained through patients' health questionnaire. In this study, there was no significant difference in PHQ-9 scores between those who adhered and did not adhere to CPAP (p=0.68) (Table 4.2.1). The score of those that adhered to CPAP was  $16 \pm 5.6$  points, which was similar to those that did not adhere to CPAP  $16 \pm 5.7$  points. A PHQ-9 score of 16 indicates moderately severe depression.

# 4.2.6 Menopausal status and adherence

Questions were asked regarding menopausal status in the health questionnaire. Subjects menopausal status were classified as follows: 1) regular menstrual cycles, irregular menstrual cycles—classified as premenopausal, 2) complete absence of periods—post-menopausal, and 3) peri-menopausal. After performing statistical analysis, there was no significant association between menopausal status in women and CPAP adherence (p=0.78) (Table 4.2.1). Women who adhered to CPAP were on average  $3.6 \pm 6.0$  years to their post-menopausal stage, which was very similar to those who did not adhere to CPAP with a mean of  $3.4 \pm 4.5$  years.

# 4.2.7 Antidepressant medication and adherence

Patients were divided into two groups including those who took antidepressant medication <1 night/week and those who took antidepressant medication ≥ 1 night/week. After testing for the amount of antidepressant medication intake in those who adhered and did not adhere to CPAP, we concluded that there was no significant association between antidepressant medication intake and CPAP adherence (p=0.57) (Table 4.2.2). Of those that adhered to CPAP, 95% of them took antidepressant medication < 1 night/week, whereas 96% of those who did not adhere to CPAP took antidepressant medication < 1 night/week.

#### 4.2.8 Gender and adherence

In general, men have less stable sleep architecture, and greater number of sleep-wake transitions than women, which may predispose to respiratory control system instability and eventually apneas. There were 226 females and 618 males in this study. Similar to many other studies, we did not find a significant association between males and females and whether one is more adherent than the other (p=0.97) (Table 4.2.2). We found 73% of the population in the adherent group to be male, which was similar to the non-adherent group.

# 4.2.9 Smoking status and adherence

In general, those who smoke tend to be less adherent to CPAP. Those who smoke are more susceptible to upper airway discomfort and as a result, less likely to take advice from the healthcare providers. <sup>108</sup> This latter theory is also confirmed by McArdle and colleagues, who reported that those who smoked were more likely to refuse CPAP therapy. <sup>22</sup> After performing chi-squared test between smoking status of those who adhered and did not adhere to CPAP, not surprisingly, we found no significant association between the two variables (p=0.91) (Table

4.2.2). We found that 16% of those in the adherent group smoked. Similarly, 16% of those in the non-adherent group smoked.

## 4.2.10 Restless leg syndrome and adherence

After analysing for restless leg syndrome from the health questionnaire, we found that there was no significant association between CPAP adherence and restless leg syndrome (p=0.76) (Table 4.2.2). 4.2% of those that adhered to CPAP had restless leg syndrome. Similarly, 4.6% of those that did not adhere to CPAP had restless leg syndrome.

#### 4.2.11 Marital status and adherence

A patient's perception of the spouse's support increases their adherence to CPAP therapy and as a result, married patients tend to be more adherent to CPAP than those that either live alone or are not married. <sup>36</sup> Consequently, many studies have identified being married as a factor for CPAP adherence. However, in this study, it was not one of the predictors of CPAP adherence (p=0.21) (Table 4.2.2). Patients were divided to the "not married" category if they were: 1) widowed, 2) divorced, 3) separated but not divorced, and 4) never married. In this study, we found that of those who adhered to CPAP, 67% of them were married, whereas, 71% of those in the non-adherent group were married.

#### 4.2.12 Bed partner and adherence

Reports in the literature indicate that those who have a bed partner tend to adhere more to CPAP because they have more social support. <sup>63</sup> Also, sleeping with a bed partner who provides feedback on the symptomatic improvements such as snoring of the patient may increase CPAP adherence. <sup>25</sup> However, in this study, we found no significant association between CPAP adherence and having a bed partner (p=0.20) (Table 4.2.2). In our cohort, 70% of those who

adhered to CPAP had a bed partner, whereas, 74% of those that did not adhered to CPAP had a bed partner.

## 4.2.13 Snoring severity and adherence

Snoring was divided according to whether the patient had been told by others that they snore more than once a week or more. In this study, snoring was not found to be a predictor of CPAP (p=0.52) (Table 4.2.2). 95% of those in the adherent group snored  $\geq$  once/week. Similarly, 94% of those in the non-adherent group snored  $\geq$  once/week.

# 4.2.14 Occupational injury and adherence

We did not find a significant association between CPAP adherence and occupational injury (p=0.16) (Table 4.2.2). Of those in the adherent group, 9.5% had occupational injury whereas, 13% of those in the non-adherent group had occupational injury.

#### 4.2.15 Neck circumference and adherence

From the results of this study, we found no significant association between neck circumference and CPAP adherence (p=0.97) (Table 4.2.2). A neck circumference of 17" was chosen as the cutoff point as the neck circumference of men is normally around 20" and for women around 14". Consequently, there was no difference in CPAP adherence between those who had a neck circumference of less than 17" and those who had a neck circumference of 17" or more.

#### 4.2.16 Mood disorder and adherence

In general, those who enjoy the activities in their life tend to have greater motivation to adhere to CPAP and those with mood disorder have poorer adherence. Similar to many studies that found no association between the two variables, us too did not find a significant association between mood disorder (i.e. depression) and CPAP adherence (p=0.17) (Table 4.2.2). In the adherent

group, 22% had mood disorder whereas, in the non-adherent group, 18% of the subjects had mood disorder.

## 4.2.17 Anxiety disorder and adherence

We found no correlation between CPAP adherence and anxiety disorder (i.e. panic disorder) (p=0.84) (Table 4.2.2). 7.1% of those in the adherent group had anxiety disorder compared to 7.5% in the non-adherent group.

#### 4.2.18 Periodic limb movement and adherence

Periodic limb movement (PLM) is very common in those who have OSA. However, in this study, we did not find an association between CPAP adherence and PLM (p=0.44) (Table 4.2.2). Only 0.6% of those in the adherent group had PLM compared to 1.2% in the non-adherent group.

# 4.2.19 Hypertension and adherence

Many studies have reported a correlation between hypertension and CPAP adherence and consider it as a strong predictor of CPAP adherence. These studies have shown that OSA may contribute to poor control of blood pressure and that 70% of patients with hypertension have OSA. <sup>109</sup>Accordingly, international guidelines have recognized OSA as one of the most common risk factors of hypertension. However, studies are inconsistent with their findings regarding whether or not those with hypertension adhere to CPAP. In our study, the association between adherence and hypertension was not strong enough for us to count hypertension as one of the predictors of CPAP (p=0.05) (Table 4.2.2). This may be due to the fact that hypertension was self-reported and that patients may have misreported whether or not they had hypertension. In addition, 25% of those in the adherent group had hypertension compared to 19% in the non-adherent group. This may be due to the fact that those with higher hypertension might have been

trying to lower their high blood pressure by using CPAP and trying to improve their quality of life in general.

## 4.2.20 Myocardial infraction and adherence

According to the results from this study, myocardial infraction (i.e. heart attack) was not one of the predictors of CPAP adherence (p=0.78) (Table 4.2.2). According to our results, 4.2% of those in the adherent group had heart attack compared to 3.8% in the non-adherent group.

# 4.2.21 Cardiac arrhythmia and adherence

According to the results from this study, no association was found between cardiac arrhythmia (i.e. irregular heartbeat) and CPAP adherence (p=0.75) (Table 4.2.2). 6.6% of those in the adherent group had irregular heart beat compared to 7.2% in the non-adherent group.

#### 4.2.22 Stroke and adherence

OSA has been suggested as a modifiable and independent risk factor for stroke, and some studies have demonstrated that those with stroke and OSA are at increased risk of death or new vascular events. <sup>110</sup> However, after analysing our results, we found no significant association between stroke and CPAP adherence (p=0.29) (Table 4.2.2). However, based on our results, 1.8% of those in the adherent group had stroke compared to only 0.87% in the non-adherent group.

#### 4.2.23 Income level and adherence

Many studies have found income level to be a predictor of CPAP adherence. Low income status is correlated with poorer health literacy, and difficulty accessing healthcare (i.e. sleep clinics and PSG test), but more specifically co-pays, needing childcare and other support, getting time off work and transport to appointments. According to Statistics Canada, the average income in British Columbia in 2015 was around \$50,000 before tax and around \$40,000 after tax. <sup>111</sup>As a result, the income in our study cohort was divided into those who earned \$40,000 or less and

those who earned more than \$40,000. However, after analysing the data, we found no significant association between income and CPAP adherence (p=0.19) (Table 4.2.2). In addition, 70% of those in the adherent group earned >\$40,000 compared to 75% in the non-adherent group.

# 4.2.24 Alcohol consumption and adherence

The role of alcohol as a predictor for CPAP adherence is still controversial as the results from various studies are inconsistent. However, one thing that is certain is that alcohol consumption increases nasal resistance. <sup>112</sup> Depending on the type of alcohol consumed, patients were divided into three groups including: 1) wine, 2) beer and 3) liquor. Alcohol consumption was further divided into those who had one glass or less per day and those who consumed more than one glass per day. After analysing the data, there was no significant association between wine consumption, beer consumption and liquor consumption and CPAP adherence with p-values being 0.46, 0.10 and 0.95 respectively. Moreover, the different alcohol consumptions were then combined to test for the overall alcohol consumption and CPAP adherence. Again, we found no significant association between the two variables (p=0.13) (Table 4.2.2). 93% of those in the adherent group drank one glass or less per day compared to 90% in the non-adherent group.

#### 4.2.25 Education and adherence

Patients were divided into two groups, those who had a college degree or higher and those without a college degree (i.e. high school diploma or lower). In total, 451 patients had a college degree or higher, whereas, 302 patients did not. After analysing the results, there was no correlation between education and CPAP adherence (p=0.14) (Table 4.2.2). Unfortunately, postsecondary education is expensive in Vancouver and many patients may not be able to afford to go to college. As a result, they may not get a high paying job and may not be able to afford

CPAP. Similarly, those who can afford to buy CPAP may not understand the importance of adhering to it and consequently, they may have lower adherence rate.

#### 4.2.26 Car crashes and adherence

Many of the published studies found a correlation between the rate of car crashes and CPAP adherence. This may be due to the fact that those who have OSA do not get sufficient sleep during the night and as a result, they get sleepy while behind the wheel. It is also possible that car crashes result from a failure of reaction, which eventually leads to driving off the road or crashing at higher speeds. <sup>113</sup> However, in our study, no significant association was found between the two variables (p=0.96) (Table 4.2.2). 16% of patients in the adherent group were in a car crash, which is similar to those in the non-adherent group.

# 4.2.27 Sleeping aid medication and adherence

Patients were divided into two groups including: 1) those who took sleeping aid medication < 1 night/week, and 2) those who took sleeping aid medication  $\ge 1$  night/week. We further found no significant association between CPAP adherence and the intake of sleeping aids (p=0.30) (Table 4.2.2).

# **Chapter 5: Discussion**

# 5.1 Summary of the results

The main findings from this single centered retrospective cohort study that included 1,250 patients were that 1) only 49.9% of the patients adhered to CPAP and 2) the only predictor that affects CPAP adherence was the severity of OSA, which was measured by AHI. There was surprisingly no significant association between the other variables including: age, daytime sleepiness (ESS), BMI, PHQ-9, menopausal status, antidepressant medication, smoking status, restless leg syndrome, marital status, bed partner status, snoring severity, occupational injury, neck circumference, mood disorder, anxiety disorder, PLM, hypertension, myocardial infraction, cardiac arrhythmia, stroke, income level, alcohol consumption, education, motor vehicle rate, gender, and sleep aid medication and CPAP adherence. However, some of these variables did trend towards significance and might have been with a larger sample size (though this is speculative).

#### **5.2 CPAP adherence**

CPAP adherence was defined as using the device for ≥ 4 hours per day for 70% of the time being used. Only 421 patients out of 844 adhered to CPAP, which indicates that only 49.9% of the patients in this study adhered to CPAP. Published CPAP adherence rates range from 30-60%, and of those who continue to use the device, 20-30% apply it insufficiently. <sup>10</sup> Moreover, over the past twenty years or so, there has been no significant improvements in CPAP adherence. Even though our result of 49.9% is within the higher range of published studies, it is still low knowing the fact that CPAP is the main treatment for OSA.

# **5.3 OSA severity**

OSA severity, as measured by AHI, is one of the most commonly examined factors in the study of sleep apnea. AHI is commonly defined as the number of apneas and hypopneas per hour while asleep. Although AHI is commonly identified as influential on CPAP adherence, the relationships are relatively inconsistent. Adding to this pool of knowledge, as predicted in our hypothesis, there was a correlation between AHI and CPAP adherence within our cohort (p=<0.0001). In other words, as the patients' OSA was more severe, their adherence to CPAP improved. The mean AHI for those who adhered to CPAP was 34 whereas the mean AHI for those who did not adhere to CPAP was 27. Moreover, 43% of those with AHI ≥30 events/hour were adherent to CPAP compared to 32% in the non-adherent group. On the other hand, 57% of those with AHI <30 events/hour were adherent to CPAP compared to 68% in the non-adherent group. Moreover, our findings support the results of other studies in the literature. <sup>20, 22, 36, 44, 114</sup> According to McArdle and colleagues, AHI was a significant determinant of CPAP adherence (p=0.004) and as the severity of OSA increased in their cohort, the use of CPAP also increased. <sup>22</sup> Similarly, in another study done by Gagnadoux and colleagues, the severity of OSA was a determinant of CPAP adherence increased (p=0.003). <sup>36</sup>

# **5.4 Strengths**

This study has a number of strengths. The most important strength of this study is that it is the first and largest of its kind in Canada. No other study regarding the predictors of CPAP adherence has been done in Canada and within a Canadian context. As a result, this study may help in improving patient care and lower OSA burden in Canada. Similarly, this study was done from 2003 to 2009, which allowed us to look at the patients' medical charts and health surveys comprehensively. By doing so, we were able to accumulate data for a large number of patients and collect as much information as we could from the charts and the health surveys. Moreover,

we uniformly investigated every patient's medical chart who came to UBC sleep clinic for a PSG test. Consequently, we were able to characterise our patients in a consistent manner. In other words, we were able to collect a subset of the "real world" as the patients were not selected for. Lastly, we found a between AHI and CPAP adherence, which is consistent with other published studies.

#### 5.5 Limitations

There are multiple important limitations within this study. The most important limitation of this study is that most of the collected data was of subjective (i.e. the sleep physician's notes from the patient's last visit), we did not have objective data, as this was not downloaded data from the CPAP devices or from PSG tests. This was mainly because during these times, objective adherence from chips in the machines was not commonly available in BC. This could have potentially removed data for patients who may have used the device for ≥ 4 hours when they used CPAP at home, as the doctor may have noted them as being non-adherent. Another limitation of this study was that it was from a single canter, meaning that it was conducted only in one clinic in one geographic area (UBC sleep clinic), which may limit its generalizability to other health regions. Additionally, the majority of the comorbidities were self-reported from the health questionnaire, which could potentially introduce reporting bias in this study. Lastly, based on the literature, CPAP adherence is established fairly early after initiating the therapy and after 2-3 months of using the device, adherence does not decrease. However, in this study, it is unclear whether the patients that adhered to CPAP will continue to use it after a few years of initiating it because we only measured adherence at one point in time (i.e. patients' follow-up appointment).

#### 5.6 Conclusions

OSA is a common respiratory disorder and commonly characterised as repetitive episodes of obstruction of the upper airway while the patient is asleep, which potentially leads to the fragmentation and hypoxemia of sleep. <sup>10</sup> The most common symptoms of OSA include silent pauses in breathing while asleep, loud snoring, excessive daytime sleepiness and frequent nocturnal awakening. <sup>10</sup> OSA affects 3-9% of the general population and is demonstrated to be a risk factor for cardiovascular diseases, increased rate of motor vehicle crashes, low quality of life, daytime sleepiness and impaired neurocognitive performance. <sup>115</sup> Direct expenses of OSA burden in Canada alone in 2008 was around 145 million dollars. <sup>5</sup> Since the pervasive health effects of untreated OSA are well described, CPAP, which improves numerous OSA-specific health outcomes, should be considered as its primary treatment. 116,117 CPAP adherence rates in the literature range from 4 to 6 hours per night. However, most studies have defined CPAP adherence as using the device  $\geq 4$  hours for  $\geq 70\%$  of the time when using the machine. Moreover, a significant proportion of those who have tried CPAP cannot adhere to the device and eventually abandon it and seek for alternate treatments. <sup>115</sup> Despite numerous technological advancements such as quieter pumps, softer and more comfortable masks, and improved portability, CPAP adherence still remains a challenge to OSA patients ranging from 30-60%, and of those who continue to use the device, 20-30% apply it insufficiently. <sup>10</sup> There are many reasons as to why patients may not adhere to CPAP. Some of the most common reasons include comfort, convenience, claustrophobia, socioeconomic status, gender, mask side effects and severity of OSA. Also, many patients remain undiagnosed as the current health care system infrastructure cannot keep up with the increasing demand for diagnostic in-laboratory PSG. <sup>118</sup>As a result, it is critical to study access to sleep clinics along with diagnostic sleep laboratories given then recent trends. 117

None of the studies that investigated the predictors of CPAP adherence were done in Canada. As a result, this study is the first and the largest of its kind that includes Canadian OSA patients. The goal of this retrospective cohort study was to extensively look at various predictors including severity of OSA, daytime sleepiness, age, BMI, PHQ-9, menopausal status, antidepressant medication, smoking status, restless leg syndrome, marital status, bed partner status, snoring severity, occupational injury, neck circumference, mood disorder, anxiety disorder, PLM, hypertension, myocardial infraction, cardiac arrhythmia, stroke, income level, alcohol consumption, education, motor vehicle rate, gender, and sleep aid medication. Out of 844 patients that were prescribed CPAP, only 421 (49.9%) adhered to CPAP. Also, only severity of OSA was correlated with CPAP adherence (p < 0.0001). In other words, as the severity of OSA increased, patients adhered more to CPAP. The findings of this study may result in the improvement of patient care and patients' overall quality of life. More importantly, we can lower OSA burden in Canada.

#### **5.7 Future studies**

As mentioned previously, OSA has many negative health outcomes. Some of the most common ones include increased risk for CV co-morbidity <sup>4</sup>, motor vehicle crashes <sup>119</sup> and hypertension <sup>120</sup>. In the short term, the next steps following this study would be to identify: 1) whether CPAP adherence results in better outcomes (i.e. cardiovascular outcomes, a lower risk of motor vehicle crashes and occupational injuries), and 2) whether other interventions (i.e. telemedicine, education, technological and behavioral) can improve adherence in Canadians. Doing so may enhance patient care and quality of life within the Canadian population and eventually lower OSA burden.

# **Bibliography**

\_\_\_\_

<sup>&</sup>lt;sup>1</sup> Somers, M. L., Peterson, E., Sharma, S., & Yaremchuk, K. (2011). Continuous positive airway pressure adherence for obstructive sleep apnea. *ISRN otolaryngology*, 2011.

<sup>&</sup>lt;sup>2</sup> Hussain, S. F., Irfan, M., Waheed, Z., Alam, N., Mansoor, S., & Islam, M. (2014). Compliance with continuous positive airway pressure (CPAP) therapy for obstructive sleep apnea among privately paying patients-a cross sectional study. *BMC pulmonary medicine*, *14*(1), 188.

<sup>&</sup>lt;sup>3</sup> Peppard, P. E., Young, T., Barnet, J. H., Palta, M., Hagen, E. W., & Hla, K. M. (2013). Increased prevalence of sleep-disordered breathing in adults. *American journal of epidemiology*, *177*(9), 1006-1014.

<sup>&</sup>lt;sup>4</sup> Golbidi, S., Badran, M., Ayas, N., & Laher, I. (2012). Cardiovascular consequences of sleep apnea. *Lung*, *190*(2), 113-132.

<sup>&</sup>lt;sup>5</sup> AlGhanim, N., Comondore, V. R., Fleetham, J., Marra, C. A., & Ayas, N. T. (2008). The economic impact of obstructive sleep apnea. *Lung*, *186*(1), 7-12.

<sup>&</sup>lt;sup>6</sup> Interventions for the treatment of obstructive sleep apnea in adults: a health technology assessment. *CADTH optimal use report*, Vol. **6**, No. 1b. Ottawa: Canadian Agency for Drugs and Technologies in Health (CADTH); 2017. Available: <a href="https://www.cadth.ca/interventions-obstructive-sleep-apnea">https://www.cadth.ca/interventions-obstructive-sleep-apnea</a>(accessed 2018 Oct. 10).

<sup>&</sup>lt;sup>7</sup> McDaid, C., Griffin, S., Weatherly, H., Durée, K., Van der Burgt, M., Van Hout, S., ... & Westwood, M. (2009). Continuous positive airway pressure devices for the treatment of obstructive sleep apnoea–hypopnoea syndrome: a systematic review and economic analysis.

<sup>&</sup>lt;sup>8</sup> Hillman, D., Mitchell, S., Streatfeild, J., Burns, C., Bruck, D., & Pezzullo, L. (2018). The economic cost of inadequate sleep. *Sleep*.

<sup>&</sup>lt;sup>9</sup> Sullivan, C., Berthon-Jones, M., Issa, F., & Eves, L. (1981). Reversal of obstructive sleep apnoea by continuous positive airway pressure applied through the nares. *The Lancet*, *317*(8225), 862-865.

<sup>&</sup>lt;sup>10</sup> Rotenberg, B. W., Murariu, D., & Pang, K. P. (2016). Trends in CPAP adherence over twenty years of data collection: a flattened curve. *Journal of Otolaryngology-Head & Neck Surgery*, *45*(1), 43.

<sup>&</sup>lt;sup>11</sup> Parish, J. M., & Somers, V. K. (2004, August). Obstructive sleep apnea and cardiovascular disease. In *Mayo Clinic Proceedings* (Vol. 79, No. 8, pp. 1036-1046). Elsevier.

<sup>&</sup>lt;sup>12</sup> Kakkar, R. K., & Berry, R. B. (2007). Positive airway pressure treatment for obstructive sleep apnea. *CHEST Journal*, *132*(3), 1057-1072.

<sup>13</sup> Budhiraja, R., Parthasarathy, S., Drake, C. L., Roth, T., Sharief, I., Budhiraja, P., ... & Hudgel, D. W. (2007). Early CPAP use identifies subsequent adherence to CPAP therapy. *Sleep*, *30*(3), 320-324.

- <sup>14</sup> Sin, D. D., Mayers, I., Man, G. C., & Pawluk, L. (2002). Long-term compliance rates to continuous positive airway pressure in obstructive sleep apnea: a population-based study. *CHEST Journal*, *121*(2), 430-435.
- <sup>15</sup> Anttalainen, U., Saaresranta, T., Kalleinen, N., Aittokallio, J., Vahlberg, T., & Polo, O. (2007). CPAP adherence and partial upper airway obstruction during sleep. *Sleep and Breathing*, *11*(3), 171-176.
- <sup>16</sup> Billings, M. E., Auckley, D., Benca, R., Foldvary-Schaefer, N., Iber, C., Redline, S., ... & Kapur, V. K. (2011). Race and residential socioeconomics as predictors of CPAP adherence. *Sleep*, *34*(12), 1653-1658.
- <sup>17</sup> Simon-Tuval, T., Reuveni, H., Greenberg-Dotan, S., Oksenberg, A., Tal, A., & Tarasiuk, A. (2009). Low socioeconomic status is a risk factor for CPAP acceptance among adult OSAS patients requiring treatment. *Sleep*, *32*(4), 545-552.
- <sup>18</sup> Gulati, A., Ali, M., Davies, M., Quinnell, T., & Smith, I. (2017). A prospective observational study to evaluate the effect of social and personality factors on continuous positive airway pressure (CPAP) compliance in obstructive sleep apnoea syndrome. *BMC Pulmonary Medicine*, 17(1), 56.
- <sup>19</sup> Campos-Rodriguez, F., Martinez-Alonso, M., Sanchez-de-la-Torre, M., Barbe, F., & Spanish Sleep Network. (2016). Long-term adherence to continuous positive airway pressure therapy in non-sleepy sleep apnea patients. *Sleep medicine*, *17*, 1-6.
- <sup>20</sup> Kohler, M., Smith, D., Tippett, V., & Stradling, J. R. (2010). Predictors of long-term compliance with continuous positive airway pressure. *Thorax*, 65(9), 829-832.
- <sup>21</sup> Krieger, J., Kurtz, D., Petiau, C., Sforza, E., & Trautmann, D. (1996). Long-term compliance with CPAP therapy in obstructive sleep apnea patients and in snorers. *Sleep*, *19*(suppl\_9), S136-S143.
- <sup>22</sup> McArdle, N., Devereux, G., Heidarnejad, H., Engleman, H. M., Mackay, T. W., & Douglas, N. J. (1999). Long-term use of CPAP therapy for sleep apnea/hypopnea syndrome. *American Journal of Respiratory and Critical Care Medicine*, *159*(4), 1108-1114.
- <sup>23</sup> Rudkowski, J. C., Verschelden, P., & Kimoff, R. J. (2001). Efficacy of daytime continuous positive airway pressure titration in severe obstructive sleep apnoea. *European Respiratory Journal*, *18*(3), 535-541.
- <sup>24</sup> Weaver, T. E., & Grunstein, R. R. (2008). Adherence to continuous positive airway pressure therapy: the challenge to effective treatment. *Proceedings of the American Thoracic Society*, *5*(2), 173-178.

<sup>25</sup> McArdle, N., Kingshott, R., Engleman, H. M., Mackay, T. W., & Douglas, N. J. (2001). Partners of patients with sleep apnoea/hypopnoea syndrome: effect of CPAP treatment on sleep quality and quality of life. *Thorax*, *56*(7), 513-518.

- <sup>26</sup> Pépin, J. L., D., Langevin, B., Robert, D., & Lévy, P. (1995). Side effects of nasal continuous positive airway pressure in sleep apnea syndrome: study of 193 patients in two French sleep centers. *Chest*, 107(2), 375-381.
- <sup>27</sup> Virk, J. S., & Kotecha, B. (2016). When continuous positive airway pressure (CPAP) fails. *Journal of Thoracic Disease*, 8(10), E1112.
- <sup>28</sup> Berthon-Jones, M., Calluaud, M., Lynch, C. E., & Hely, K. F. (1996). *U.S. Patent No.* 5,560,354. Washington, DC: U.S. Patent and Trademark Office.
- <sup>29</sup> Rowland, S., Aiyappan, V., Hennessy, C., Catcheside, P., Chai-Coezter, C. L., McEvoy, R. D., & Antic, N. A. (2018). Comparing the Efficacy, Mask Leak, Patient Adherence, and Patient Preference of Three Different CPAP Interfaces to Treat Moderate-Severe Obstructive Sleep Apnea. *Journal of Clinical Sleep Medicine*, *14*(01), 101-108
- <sup>30</sup> Billings, M. E., Rosen, C. L., Wang, R., Auckley, D., Benca, R., Foldvary-Schaefer, N., ... & Kapur, V. K. (2013). Is the relationship between race and continuous positive airway pressure adherence mediated by sleep duration?. *Sleep*, *36*(2), 221-227.
- <sup>31</sup> Joo, M. J., & Herdegen, J. J. (2007). Sleep apnea in an urban public hospital: assessment of severity and treatment adherence. *Journal of Clinical Sleep Medicine*, *3*(03), 285-288.
- <sup>32</sup> Collen, J., Holley, A., Lettieri, C., Shah, A., & Roop, S. (2010). The impact of split-night versus traditional sleep studies on CPAP compliance. *Sleep and Breathing*, *14*(2), 93-99.
- <sup>33</sup> Palm, A., Midgren, B., Theorell-Haglöw, J., Janson, C., & Lindberg, E. (2017). Factors influencing compliance to continuous positive airway pressure treatment in obstructive sleep apnea and mortality associated with treatment failure. *Sleep Medicine*, 40, e250.
- <sup>34</sup> Woehrle, H., Graml, A., & Weinreich, G. (2011). Age-and gender-dependent adherence with continuous positive airway pressure therapy. *Sleep medicine*, *12*(10), 1034-1036.
- <sup>35</sup> Means, M. K., Ulmer, C. S., & Edinger, J. D. (2010). Ethnic differences in continuous positive airway pressure (CPAP) adherence in veterans with and without psychiatric disorders. *Behavioral sleep medicine*, 8(4), 260-273.
- <sup>36</sup> Gagnadoux, F., Le Vaillant, M., Goupil, F., Pigeanne, T., Chollet, S., Masson, P., ... & Meslier, N. (2011). Influence of marital status and employment status on long-term adherence with continuous positive airway pressure in sleep apnea patients. *PLoS One*, 6(8), e22503.
- <sup>37</sup> Stuck, B. A., Leitzbach, S., & Maurer, J. T. (2012). Effects of continuous positive airway pressure on apnea—hypopnea index in obstructive sleep apnea based on long-term compliance. *Sleep and Breathing*, *16*(2), 467-471.

<sup>38</sup> Pamidi, S., Knutson, K. L., Ghods, F., & Mokhlesi, B. (2012). The impact of sleep consultation prior to a diagnostic polysomnogram on continuous positive airway pressure adherence. *Chest*, *141*(1), 51-57.

- <sup>39</sup> Campbell, A., Neill, A., & Lory, R. (2012). Ethnicity and socioeconomic status predict initial continuous positive airway pressure compliance in New Zealand adults with obstructive sleep apnoea. *Internal medicine journal*, 42(6), e95-e101.
- <sup>40</sup> Platt, A. B., Field, S. H., Asch, D. A., Chen, Z., Patel, N. P., Gupta, R., ... & Kuna, S. T. (2009). Neighborhood of residence is associated with daily adherence to CPAP therapy. *Sleep*, *32*(6), 799-806.
- <sup>41</sup> Wallace, D. M., Vargas, S. S., Schwartz, S. J., Aloia, M. S., & Shafazand, S. (2013). Determinants of continuous positive airway pressure adherence in a sleep clinic cohort of South Florida Hispanic veterans. *Sleep and Breathing*, *17*(1), 351-363.
- <sup>42</sup> Galetke, W., Puzzo, L., Priegnitz, C., Anduleit, N., & Randerath, W. J. (2011). Long-term therapy with continuous positive airway pressure in obstructive sleep apnea: adherence, side effects and predictors of withdrawal–a 'real-life' study. *Respiration*, 82(2), 155-161.
- <sup>43</sup> Waldhorn, R. E., Herrick, T. W., Nguyen, M. C., O'Donnell, A. E., Sodero, J., & Potolicchio, S. J. (1990). Long-term compliance with nasal continuous positive airway pressure therapy of obstructive sleep apnea. *Chest*, *97*(1), 33-38.
- <sup>44</sup> Meurice, J. C., Dore, P., Paquereau, J., Neau, J. P., Ingrand, P., Chavagnat, J. J., & Patte, F. (1994). Predictive factors of long-term compliance with nasal continuous positive airway pressure treatment in sleep apnea syndrome. *Chest*, *105*(2), 429-433.
- <sup>45</sup> Engleman, H. M., Asgari-Jirhandeh, N., McLeod, A. L., Ramsay, C. F., Deary, I. J., & Douglas, N. J. (1996). Self-reported use of CPAP and benefits of CPAP therapy: a patient survey. *Chest*, *109*(6), 1470-1476.
- <sup>46</sup> Krieger, J., Kurtz, D., Petiau, C., Sforza, E., & Trautmann, D. (1996). Long-term compliance with CPAP therapy in obstructive sleep apnea patients and in snorers. *Sleep*, *19*(suppl\_9), S136-S143.
- <sup>47</sup> PÉpin, J. L., Krieger, J., Rodenstein, D., Cornette, A., Sforza, E., Delguste, P., ... & LÉvy, P. (1999). Effective compliance during the first 3 months of continuous positive airway pressure: a European prospective study of 121 patients. *American journal of respiratory and critical care medicine*, *160*(4), 1124-1129.
- <sup>48</sup> Hui, D. S., Choy, D. K., Li, T. S., Ko, F. W., Wong, K. K., Chan, J. K., & Lai, C. K. (2001). Determinants of continuous positive airway pressure compliance in a group of Chinese patients with obstructive sleep apnea. *Chest*, *120*(1), 170-176.
- <sup>49</sup> McARDLE, N. I. G. E. L., & Douglas, N. J. (2001). Effect of continuous positive airway pressure on sleep architecture in the sleep apnea–hypopnea syndrome: a randomized controlled trial. *American journal of respiratory and critical care medicine*, *164*(8), 1459-1463.

<sup>50</sup> Montserrat, J. M., Ferrer, M., Hernandez, L., FarrÉ, R., Vilagut, G., Navajas, D., ... & Puig, M. (2001). Effectiveness of CPAP treatment in daytime function in sleep apnea syndrome: a randomized controlled study with an optimized placebo. *American Journal of Respiratory and Critical Care Medicine*, *164*(4), 608-613.

- <sup>51</sup> Barnes, M., McEvoy, R. D., Banks, S., Tarquinio, N., Murray, C. G., Vowles, N., & Pierce, R. J. (2004). Efficacy of positive airway pressure and oral appliance in mild to moderate obstructive sleep apnea. *American journal of respiratory and critical care medicine*, 170(6), 656-664.
- <sup>52</sup> Rosenthal, L., Gerhardstein, R., Lumley, A., Guido, P., Day, R., Syron, M. L., & Roth, T. (2000). CPAP therapy in patients with mild OSA: implementation and treatment outcome. *Sleep Medicine*, *1*(3), 215-220.
- <sup>53</sup> Budhiraja, R., Kushida, C. A., Nichols, D. A., Walsh, J. K., Simon, R. D., Gottlieb, D. J., & Quan, S. F. (2017). Predictors of sleepiness in obstructive sleep apnoea at baseline and after 6 months of continuous positive airway pressure therapy. *European Respiratory Journal*, 50(5), 1700348.
- <sup>54</sup> Marin, J. M., Agusti, A., Villar, I., Forner, M., Nieto, D., Carrizo, S. J., ... & Jelic, S. (2012). Association between treated and untreated obstructive sleep apnea and risk of hypertension. *Jama*, 307(20), 2169-2176.
- <sup>55</sup> Tarasiuk, A., Reznor, G., Greenberg-Dotan, S., & Reuveni, H. (2012). Financial incentive increases CPAP acceptance in patients from low socioeconomic background. *PLoS One*, 7(3), e33178.
- <sup>56</sup> Wild, M. R., Engleman, H. M., Douglas, N. J., & Espie, C. A. (2004). Can psychological factors help us to determine adherence to CPAP? A prospective study. *European Respiratory Journal*, *24*(3), 461-465.
- <sup>57</sup> Hayes MJ, McGregor FB, Roberts DN, Schroter RC, Pride NB. Contin- uous nasal positive airway pressure with a mouth leak: effect on nasal mucosal blood flux and nasal geometry. Thorax 1995;50:1179–1182.
- <sup>58</sup> Demolles, D. A., Sparrow, D., Gottlieb, D. J., & Friedman, R. (2004). A pilot trial of a telecommunications system in sleep apnea management. *Medical care*, 42(8), 764-769.
- <sup>59</sup> Smith, C. E., Dauz, E. R., Clements, F., Puno, F. N., Cook, D., Doolittle, G., & Leeds, W. (2006). Telehealth services to improve nonadherence: a placebo-controlled study. *Telemedicine Journal & e-Health*, *12*(3), 289-296.
- <sup>60</sup> Stepnowsky, C. J., Marler, M. R., & Ancoli-Israel, S. (2002). Determinants of nasal CPAP compliance. *Sleep medicine*, *3*(3), 239-247.
- <sup>61</sup> Kjelsberg, F. N., Ruud, E. A., & Stavem, K. (2005). Predictors of symptoms of anxiety and depression in obstructive sleep apnea. *Sleep Medicine*, *6*(4), 341-346.

<sup>62</sup> Ayalon, L., Ancoli-Israel, S., Stepnowsky, C., Marler, M., Palmer, B. W., Liu, L., ... & Cooke, J. (2006). Adherence to continuous positive airway pressure treatment in patients with Alzheimer disease and obstructive sleep apnea. *The American journal of geriatric psychiatry*, *14*(2), 176-180.

- <sup>63</sup> Lewis, K. E., Seale, L., Bartle, I. E., Watkins, A. J., & Ebden, P. (2004). Early predictors of CPAP use for the treatment of obstructive sleep apnea. *Sleep*, *27*(1), 134-138.
- <sup>64</sup> Stepnowsky Jr, C. J., Bardwell, W. A., Moore, P. J., Ancoli-Israel, S., & Dimsdale, J. E. (2002). Psychologic correlates of compliance with continuous positive airway pressure. *Sleep*, *25*(7), 758-762.
- <sup>65</sup> Pieh, C., Bach, M., Popp, R., Jara, C., Crönlein, T., Hajak, G., & Geisler, P. (2013). Insomnia symptoms influence CPAP compliance. *Sleep and Breathing*, 17(1), 99-104.
- <sup>66</sup> Philip, P., Bioulac, S., Altena, E., Morin, C. M., Ghorayeb, I., Coste, O., ... & Micoulaud-Franchi, J. A. (2018). Specific insomnia symptoms and self-efficacy explain CPAP compliance in a sample of OSAS patients. *PloS one*, *13*(4), e0195343.
- <sup>67</sup> Micoulaud-Franchi, J. A., Coste, O., Bioulac, S., Guichard, K., Monteyrol, P. J., Ghorayeb, I., ... & Philip, P. (2018). A French update on the Self-Efficacy Measure for Sleep Apnea (SEMSA) to assess continuous positive airway pressure (CPAP) use. *Sleep and Breathing*, 1-10.
- <sup>68</sup> Bakker, J. P., O'Keeffe, K. M., Neill, A. M., & Campbell, A. J. (2011). Ethnic disparities in CPAP adherence in New Zealand: effects of socioeconomic status, health literacy and self-efficacy. *Sleep*, *34*(11), 1595-1603.
- <sup>69</sup> Law, M., Naughton, M., Ho, S., Roebuck, T., & Dabscheck, E. (2014). Depression may reduce adherence during CPAP titration trial. *Journal of Clinical Sleep Medicine*, 10(02), 163-169.
- <sup>70</sup> Kaplan, J. L., Chung, S. A., Fargher, T., & Shapiro, C. M. (2007). The effect of one versus two nights of in-laboratory continuous positive airway pressure titration on continuous positive airway pressure compliance. *Behavioral sleep medicine*, *5*(2), 117-129.
- <sup>71</sup> Poulet, C., Veale, D., Arnol, N., Levy, P., Pepin, J. L., & Tyrrell, J. (2009). Psychological variables as predictors of adherence to treatment by continuous positive airway pressure. *Sleep medicine*, *10*(9), 993-999.
- <sup>72</sup> Campos-Rodriguez, F., Martinez-Garcia, M. A., Reyes-Nuñez, N., Caballero-Martinez, I., Catalan-Serra, P., & Almeida-Gonzalez, C. V. (2014). Role of sleep apnea and continuous positive airway pressure therapy in the incidence of stroke or coronary heart disease in women. *American journal of respiratory and critical care medicine*, *189*(12), 1544-1550.
- <sup>73</sup> McEvoy, R. D., Antic, N. A., Heeley, E., Luo, Y., Ou, Q., Zhang, X., ... & Chen, G. (2016). CPAP for prevention of cardiovascular events in obstructive sleep apnea. *New England Journal of Medicine*, *375*(10), 919-931.

<sup>74</sup> Nguyên, X. L., Chaskalovic, J., Rakotonanahary, D., & Fleury, B. (2010). Insomnia symptoms and CPAP compliance in OSAS patients: A descriptive study using Data Mining methods. *Sleep medicine*, *11*(8), 777-784.

- <sup>75</sup> Ritterband, L. M., Thorndike, F. P., Gonder-Frederick, L. A., Magee, J. C., Bailey, E. T., Saylor, D. K., & Morin, C. M. (2009). Efficacy of an Internet-based behavioral intervention for adults with insomnia. *Archives of general psychiatry*, 66(7), 692-698.
- <sup>76</sup> Gooneratne, N. S., Gehrman, P., Gurubhagavatula, I., Al-Shehabi, E., Marie, E., & Schwab, R. (2010). Effectiveness of ramelteon for insomnia symptoms in older adults with obstructive sleep apnea: a randomized placebo-controlled pilot study. *Journal of clinical sleep medicine: JCSM: official publication of the American Academy of Sleep Medicine*, 6(6), 572.
- <sup>77</sup> Scharf, S. M., Seiden, L., DeMore, J., & Carter-Pokras, O. (2004). Racial differences in clinical presentation of patients with sleep-disordered breathing. *Sleep and Breathing*,  $\delta$ (04), 173-183.
- <sup>78</sup> Edinger, J. D., & Means, M. K. (2005). Cognitive—behavioral therapy for primary insomnia. *Clinical psychology review*, 25(5), 539-558.
- <sup>79</sup> Schwab, R. J., Badr, S. M., Epstein, L. J., Gay, P. C., Gozal, D., Kohler, M., ... & Strohl, K. P. (2013). An official American Thoracic Society statement: continuous positive airway pressure adherence tracking systems. The optimal monitoring strategies and outcome measures in adults. *American journal of respiratory and critical care medicine*, *188*(5), 613-620.
- <sup>80</sup> Bachour, A., & Maasilta, P. (2004). Mouth breathing compromises adherence to nasal continuous positive airway pressure therapy. *Chest*, *126*(4), 1248-1254.
- <sup>81</sup> Mortimore, I. L., Bradley, P. A., Murray, J. A., & Douglas, N. J. (1996). Uvulopalatopharyngoplasty may compromise nasal CPAP therapy in sleep apnea syndrome. *American journal of respiratory and critical care medicine*, *154*(6), 1759-1762.
- <sup>82</sup> Weaver, T. E., & Sawyer, A. M. (2010). Adherence to continuous positive airway pressure treatment for obstructive sleep apnea: implications for future interventions. *The Indian journal of medical research*, 131, 245.
- <sup>83</sup> Lai, A. Y., Fong, D. Y., Lam, J. C., Weaver, T. E., & Ip, M. S. (2014). The efficacy of a brief motivational enhancement education program on CPAP adherence in OSA: a randomized controlled trial. *Chest*, *146*(3), 600-610.
- <sup>84</sup> Neill, A. M., Wai, H. S., Bannan, S. P. T., Beasley, C. R., Weatherall, M., & Campbell, A. J. (2003). Humidified nasal continuous positive airway pressure in obstructive sleep apnoea. *European Respiratory Journal*, *22*(2), 258-262.
- <sup>85</sup> Borel, J. C., Tamisier, R., Dias-Domingos, S., Sapene, M., Martin, F., Stach, B., ... & Pepin, J. L. (2013). Type of mask may impact on continuous positive airway pressure adherence in apneic patients. *PloS one*, 8(5), e64382.

<sup>86</sup> Kreivi, H. R., Virkkula, P., Lehto, J., & Brander, P. (2010). Frequency of upper airway symptoms before and during continuous positive airway pressure treatment in patients with obstructive sleep apnea syndrome. *Respiration*, 80(6), 488-494.

- <sup>87</sup> Kalan, A., Kenyon, G. S., Seemungal, T. A. R., & Wedzicha, J. A. (1999). Adverse effects of nasal continuous positive airway pressure therapy in sleep apnoea syndrome. *The Journal of Laryngology & Otology*, *113*(10), 888-892.
- <sup>88</sup> Pepin, J. L., Leger, P., Veale, D., Langevin, B., Robert, D., & Lévy, P. (1995). Side effects of nasal continuous positive airway pressure in sleep apnea syndrome: study of 193 patients in two French sleep centers. *Chest*, 107(2), 375-381.
- <sup>89</sup> Reid, J. (1996). The History of Telemedicine (in brief): a telemedicine primer. *Billings: Innovative Medical Communications*, 5-27.
- <sup>90</sup> Kwiatkowska, M., & Ayas, N. (2010). Can telemedicine improve CPAP adherence? *Thorax*, 65(12), 1035-1036. doi:10.1136/thx.2010.140897
- <sup>91</sup> Stepnowsky Jr, C. J., Palau, J. J., Marler, M. R., & Gifford, A. L. (2007). Pilot randomized trial of the effect of wireless telemonitoring on compliance and treatment efficacy in obstructive sleep apnea. *Journal of medical Internet research*, 9(2).
- <sup>92</sup> Wiese, H. J., Boethel, C., Phillips, B., Wilson, J. F., Peters, J., & Viggiano, T. (2005). CPAP compliance: video education may help!. *Sleep medicine*, *6*(2), 171-174.
- <sup>93</sup> Golay, A., Girard, A., Grandin, S., Métrailler, J. C., Victorion, M., Lebas, P., ... & Rochat, T. (2006). A new educational program for patients suffering from sleep apnea syndrome. *Patient education and counseling*, 60(2), 220-227.
- <sup>94</sup> Sparrow, D., Aloia, M., DeMolles, D. A., & Gottlieb, D. J. (2010). A telemedicine intervention to improve adherence to continuous positive airway pressure: a randomised controlled trial. *Thorax*, *65*(12), 1061-1066.
- <sup>95</sup> Hwang, D., Chang, J. W., Benjafield, A. V., Crocker, M. E., Kelly, C., Becker, K. A., ... & Derose, S. F. (2018). Effect of telemedicine education and telemonitoring on continuous positive airway pressure adherence. The tele-OSA randomized trial. *American journal of respiratory and critical care medicine*, 197(1), 117-126.
- <sup>96</sup> Richards, D., Bartlett, D. J., Wong, K., Malouff, J., & Grunstein, R. R. (2007). Increased adherence to CPAP with a group cognitive behavioral treatment intervention: a randomized trial. *Sleep*, *30*(5), 635-640.
- <sup>97</sup> Aloia, M. S., Smith, K., Arnedt, J. T., Millman, R. P., Stanchina, M., Carlisle, C., ... & Borrelli, B. (2007). Brief behavioral therapies reduce early positive airway pressure discontinuation rates in sleep apnea syndrome: preliminary findings. *Behavioral sleep medicine*, *5*(2), 89-104.

<sup>98</sup> Malhotra, A., Crocker, M. E., Willes, L., Kelly, C., Lynch, S., & Benjafield, A. V. (2018). Patient engagement using new technology to improve adherence to positive airway pressure therapy: a retrospective analysis. *Chest*, *153*(4), 843-850.

- <sup>99</sup> Mador, M. J., Krauza, M., Pervez, A., Pierce, D., & Braun, M. (2005). Effect of heated humidification on compliance and quality of life in patients with sleep apnea using nasal continuous positive airway pressure. *Chest*, 128(4), 2151-2158.
- <sup>100</sup> Hussain, S. F., Irfan, M., Waheed, Z., Alam, N., Mansoor, S., & Islam, M. (2014). Compliance with continuous positive airway pressure (CPAP) therapy for obstructive sleep apnea among privately paying patients-a cross sectional study. *BMC pulmonary medicine*, *14*(1), 188.
- <sup>101</sup> Peppard, P. E., Young, T., Barnet, J. H., Palta, M., Hagen, E. W., & Hla, K. M. (2013). Increased prevalence of sleep-disordered breathing in adults. *American journal of epidemiology*, *177*(9), 1006-1014.
- <sup>102</sup> Berry, R. B., Budhiraja, R., Gottlieb, D. J., Gozal, D., Iber, C., Kapur, V. K., ... & Redline, S. (2012). Rules for scoring respiratory events in sleep: update of the 2007 AASM manual for the scoring of sleep and associated events. *Journal of clinical sleep medicine*, 8(05), 597-619)
- <sup>103</sup> Mbata, G., & Chukwuka, J. (2012). Obstructive Sleep Apnea Hypopnea Syndrome. *Annals of Medical and Health Sciences Research*, 2(1), 74–77. http://doi.org/10.4103/2141-9248.96943
- <sup>104</sup> (Sleep-related breathing disorders in adults: recommendations for syndrome definition and measurement techniques in clinical research. (1999). The report of an American Academy of Sleep Medicine task force. *Sleep*, 22(5), 667-689)
- <sup>105</sup> Park, P., Kim, J., Song, Y. J., Lim, J. H., Cho, S. W., Won, T. B., ... & Kim, H. J. (2017). Influencing factors on CPAP adherence and anatomic characteristics of upper airway in OSA subjects. *Medicine*, *96*(51).
- <sup>106</sup> Kohler, M., Bloch, K. E., & Stradling, J. R. (2007). The role of the nose in the pathogenesis of obstructive sleep apnoea and snoring. *European Respiratory Journal*, *30*(6), 1208-1215.
- <sup>107</sup> Edwards, C., Mukherjee, S., Simpson, L., Palmer, L. J., Almeida, O. P., & Hillman, D. R. (2015). Depressive symptoms before and after treatment of obstructive sleep apnea in men and women. *Journal of clinical sleep medicine*, *11*(09), 1029-1038.
- <sup>108</sup> Jacobsen, A. R., Eriksen, F., Hansen, R. W., Erlandsen, M., Thorup, L., Damgård, M. B., . . . Hansen, K. W. (2017). Determinants for adherence to continuous positive airway pressure therapy in obstructive sleep apnea. *PloS One*, *12*(12), e0189614. doi:10.1371/journal.pone.0189614
- <sup>109</sup> Martínez-García, M. A., Capote, F., Campos-Rodríguez, F., Lloberes, P., de Atauri, M. J. D., Somoza, M., ... & Durán-Cantolla, J. (2013). Effect of CPAP on blood pressure in patients with obstructive sleep apnea and resistant hypertension: the HIPARCO randomized clinical trial. *Jama*, *310*(22), 2407-2415.

- <sup>110</sup> Martínez-García, M. Á., Soler-Cataluña, J. J., Ejarque-Martínez, L., Soriano, Y., Román-Sánchez, P., Illa, F. B., ... & Durán-Cantolla, J. (2009). Continuous positive airway pressure treatment reduces mortality in patients with ischemic stroke and obstructive sleep apnea: a 5-year follow-up study. *American journal of respiratory and critical care medicine*, *180*(1), 36-41.
- <sup>111</sup> Statistics Canada. (2018). *Table 11-10-0008-01* Tax filers and dependants with income by total income, sex and age. Retrieved from: https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=1110000801
- <sup>112</sup> Jeong, J. I., Kim, H. Y., Hong, S. D., Ryu, G., Kim, S. J., Lee, K. E., ... & Chung, S. K. (2016). Upper Airway Variation and Frequent Alcohol Consumption Can Affect Compliance With Continuous Positive Airway Pressure. *Clinical and experimental otorhinolaryngology*, *9*(4), 346.
- <sup>113</sup> Tregear, S., Reston, J., Schoelles, K., & Phillips, B. (2010). Continuous positive airway pressure reduces risk of motor vehicle crash among drivers with obstructive sleep apnea: systematic review and meta-analysis. *Sleep*, *33*(10), 1373-1380.
- <sup>114</sup> Hui, D. S. C., Choy, D. K. L., Li, T. S. T., Ko, F. W. S., Wong, K. K., Chan, J. K. W., & Lai, C. K. W. (2001). Determinants of continuous positive airway pressure compliance in a group of chinese patients with obstructive sleep apnea. *Chest*, *120*(1), 170-176. doi:10.1378/chest.120.1.170)
- <sup>115</sup> Young T, Peppard PE, Gottlieb DJ. Epidemiology of obstructive sleep apnea: a population health perspective. Am J Respir Crit Care Med. 2002;165(9):1217–39.
- <sup>116</sup> Aurora, R. N., Casey, K. R., Kristo, D., Auerbach, S., Bista, S. R., Chowdhuri, S., ... & Morgenthaler, T. I. (2010). Practice parameters for the surgical modifications of the upper airway for obstructive sleep apnea in adults. *Sleep*, *33*(10), 1408-1413.
- <sup>117</sup> Yaremchuk, K., Tacia, B., Peterson, E., & Roth, T. (2011). Change in Epworth Sleepiness Scale after surgical treatment of obstructive sleep apnea. *The Laryngoscope*, *121*(7), 1590-1593.
- <sup>118</sup> Evans, J., Skomro, R., Driver, H., Graham, B., Mayers, I., McRae, L., ... & Fleetham, J. (2014). Sleep laboratory test referrals in Canada: sleep apnea rapid response survey. *Canadian respiratory journal*, *21*(1), e4-e10.
- <sup>119</sup> Tregear, S., Reston, J., Schoelles, K., & Phillips, B. (2009). Obstructive sleep apnea and risk of motor vehicle crash: systematic review and meta-analysis. *Journal of clinical sleep medicine*, *5*(06), 573-581.
- <sup>120</sup> Torres, G., Sánchez-de-la-Torre, M., & Barbé, F. (2015). Relationship between OSA and hypertension. *Chest*, *148*(3), 824-832.

## Appendices

pp	en	dix A: Health quest	ionnaire					- 10g
		ID Number	То	day's Date /	$\Box'\Box$	MM/DD/	w [	<b>82</b>
		Vanco	uver Co	oastal				
			F	ealth	Authori	ty		
				aribaldi, Vancouv				
M	OGF	RAPHICS:						
	1.	What is your date of birth:	/		MM/DD/YYY	<b>Y</b>		
	2.	What is your gender:						
		☐ Male ☐ Fe	male					
	3.	What is your current Marital	status:					
		☐ never married ☐ married ☐ separated but no ☐ divorced ☐ widowed/widowe	t divorced					
	4.	Does another person regula	rly share your be	ed/sleep with you?:				
		□ Yes □ N	o					
	5.	What is the highest level of	education you ha	ave completed?				
		☐ Less than high s	chool diploma					
		☐ High school diple						
		☐ College or Unive						
		☐ Masters or abov	е					
<u>.E</u> l	EP I	HABITS:						
	6.	After you have turned out the	ne lights to fall as	sleep, how long does it <u>usu</u>	ally take you to fa	all asleep?		
		☐ 0-15 minutes						
		☐ 16-30 minutes						
		31-60 minutes						
		☐ 61-120 minutes						
		more than 2 hou						
	7.	On average, how many hou	irs of sleep do yo	ou get per night on weekda	ays?			
		□≤5 □6	<b>0</b> 7	□8	□9	□ 10+	hours	
		On weekends?						
		□≤5 □6	<b>0</b> 7	□8	□9	□ 10+	hours	
	8	On average, how many hou	urs do you spend	I napping during the daytin	ne on weekdays?			
		☐ None or nearly none	□ 1-2	□ 3-4	□ 5-6	□7+	hours	
		On weekends?						
		☐ None or nearly none	□ 1-2	□ 3-4	□ 5-6	<b>17</b> +	hours	
								Q#

	ver the last 24 months, which of the following, if any, did you have to do to keep from falling asleep at wor
9. C	hoose all that apply.
	☐ I don't have any problems staying awake at work
S	☐ chew gum or candy
	□ eat
	□ smoke
	☐ chew tobacco
	☐ drink coffee
	☐ drink other caffeinated beverages
	☐ listen to radio/music
	☐ moves around/keep physically active
	other (please describe)
	Other (please describe)
SLEEP-REL	ATED SYMPTOMS:
10. Do	you have difficulty falling or staying asleep?
	never
	☐ rarely (less than one night per week)
	sometimes (1-2 nights/week)
	☐ frequently (3-4 nights/week)
	☐ almost always (5-7 nights/week)
11. Or	average, how many days/nights during the last month have you snored or been told you snored?
	□ never
	☐ rarely (less than once per week)
	sometimes (1-2/week)
	☐ frequently (3-4/week)
	☐ almost always (5-7/week)
	☐ not sure
12. Wh	en falling asleep, how often do you have "restless legs" (a feeling of crawling, aching, or inability to p legs still)?
	never
	rarely (less than one night per week)
	☐ sometimes (1-2 nights/week)
	☐ frequently (3-4 nights/week)
	☐ almost always (5-7 nights/week)
13. In ti	e last 24 months, have you found that you awaken and briefly cannot move your entire body (i.e. felt layzed as opposed to feeling very tired)?
	□ Yes
	□ No
	U NO
14. In t	ne last 24 months, have you seen or heard things that don't exist upon waking up or falling asleep?
	□No
	in last 24 marks.
15 In t	
15. In t	e last 24 months, have you had sudden muscular weakness (e.g. buckling of your knees) associated
15. In to 6705	emotion (e.g. while laughing or after hearing a joke)?

ф

DAYTIME A	CTIVITY
-----------	---------

j	in the last 30 days asleep in the follo ust tired)? This re times. <u>Even if you</u> recently try to wor	wing situat efers to you have not d	ions (in c ir usual v done som	contrast vay of lift as of the	to feeling fe in rece	) int	HIGH CHAN		ODERATI CHANCE		GHT ANCE	NO CHANCE
a. \$	Sitting and readin	g					0		۵		0	
b. \	Watching TV							1	0		o -	0
	Sitting inactive in					)		-	0		0	
	As a passenger i				a break	60 S.5 P.T.	Contract Con		0	n dispersion.	0	0
e.	Lying down to res circumstances p	it in the afte ermit.	ernoon w	hen					0		0	0
	Sitting and talking							)	0		0	0
g. :	Sitting quietly after	er lunch wit	thout alco	ohol.					0		0	0
h.	In a car while sto	pped for a	few mini	ites in t	raffic.				0	The state of	0	
a. Cola or carbonated with caffein Pepsi, Mou	that per other beverage se, e.g. Coke,		How man How man use of the 1-3 per month	ny years	s?		iring the	e past ye 2-3 per day		6-10   per day	11-20 per day	more than 20 day
b. Tea or	iced tea with cup or glass), teas	0	0	a	0	0	0	۵	۵	0	0	0
c. Coffee (1 cup)	with caffeine	۵	0	0	0		0		п	۵	0	0
d. Beer (1 or can)	glass, bottle,	0	۵		0	۵	0	۵	0	۵	0	0
e. Wine (4	4 oz glass)	0		0	۵	۵	0	۵	0	٥		а
	e.g. vodka, drink or shot)	0	0	0	0	0	0	0	0	a	0	0

Ô.



n	^	CI	IP	ΔT	'n	NA	L:

	y months did you work?
eks that	
	you worked, approximately how many hours did you work
e last 24	months (describe)?
0 1001 2	menure (decarbe).
characte	erized your occupation over the last 24 months:
<u>_</u>	Forestry
	Oil and Gas or Mineral Resources
	Petroleum, Coal, Rubber, Plastic, and Chemical Production
ses)	Road Construction or Maintenance
	Transportation and Related Services
	Wholesale
g for the	federal government government)
	lover(milent)
es 🗆	Healthcare and Social Assistance Professional, Scientific, and Technical Services
	Utilities
	characte

s a percentage, how many of each of the fol otal should equal 100%)	llowing si	hifts did you	work over th	e last 24 mo	enths	
Day shift (e.g. 8am -5 pm)	<u></u> %					
Evening shift (e.g. 3 pm-11pm)	<u></u> %					
Nightshift (e.g. 11 pm to 7 am)	%					
	Y Too					
iven the hours you work, how much on average a year?	rage wou	ıld someone	employed in	a field like	ours eam	
□ \$0-20 000						
□ 21,000 -40,000						
□ 41,000-60,000						
□ 61,000-80,000						
□ 81,000-100,000						
□ >100,000						
over the last 24 months, did you suffer an oc	ccupation	al injury?				
over the last 24 months, did you suffer an od Yes No  If YES, please answer the following clust If NO, please skip to question 27.						
If YES, please answer the following clust		estions.	2	3	44	
☐ Yes ☐ No  If YES, please answer the following clust	ter of que	estions.	2	3	4+	
☐ Yes ☐ No  If YES, please answer the following clust If NO, please skip to question 27.	ter of que	estions.	2	3	4+ □	
☐ Yes ☐ No  If YES, please answer the following clust If NO, please skip to question 27.  a. If so, how many injuries did you have?	ter of que	estions.				
☐ Yes ☐ No  If YES, please answer the following clust If NO, please skip to question 27.  a. If so, how many injuries did you have?  b. How many of these, if any, were reported to the Workers Compensation Board?	ter of que	estions.	0	0	0	
<ul> <li>Yes □ No</li> <li>If YES, please answer the following clust If NO, please skip to question 27.</li> <li>a. If so, how many injuries did you have?</li> <li>b. How many of these, if any, were reported to the Workers Compensation Board?</li> <li>c. In how many of these, if any, do you think sleepiness or fatigue in you</li> </ul>	0	1	0	0	0	

٧	_	
1		

	☐ Yes ☐ No						
	If YES, please answer the following of If NO, please skip to question 28.	luster of q	uestions.				
		0	1	2	3	4+	
a.	How many injuries did you cause?	0	0	0	0		
b.	How many of these, if any, were reported to the Workers Compensation Board?	0	0	0	0	0	
C.	In how many of these, if any, do you think sleepiness or fatigue in you contributed?	0	0		0	0	
d.	In how many of these, if any, was an emergency room visit required?		0	0		0	
e.	Please take a moment and tell us a few words about each incident						
	lew words about each incident						
CAL I	HISTORY:						
CAL I	HISTORY:						
	HISTORY:	Inches	s OR	c	m		
28. W					m s OR		kg
28. W 29. W	hat is your height?	st pound)?					kg
28. W 29. W 30. W	hat is your height? Feet hat is your current weight (to the neares hen was your weight last measured? W	it pound)?	st	Pound			kg
28. W 29. W 30. W	hat is your height? Feet feet hat is your current weight (to the neares	st pound)? ithin the la	stthe nearest	Pound	s OR	121 22	kg □ 22+
28. W 29. W 80. W	That is your height? Feet hat is your current weight (to the neares hen was your weight last measured? What is your neck circumference (e.g. shi	st pound)? ithin the la rt size) to t	the nearest	Pound Months inch?	s OR	121 🗆 22	
28. W 29. W 30. W 31. W	That is your height? Feet hat is your current weight (to the nearest hen was your weight last measured? What is your neck circumference (e.g. shill ≤11	st pound)? ithin the la rt size) to t  16	the nearest	Pound Months inch? 18 □ 19	s OR □ 20 □		
28. W 29. W 30. W 31. W	That is your height? Feet hat is your current weight (to the nearest hen was your weight last measured? What is your neck circumference (e.g. shith of the same weight last describes your neck the sentence that best describes your last last last last last last last last	st pound)?  ithin the la  rt size) to t    16  our situatio	the nearest	Pound Months inch? 18 □ 19	s OR □ 20 □		
28. W 29. W 30. W 31. W	That is your height? Feet hat is your current weight (to the nearest hen was your weight last measured? What is your neck circumference (e.g. shill ≤11	ithin the la rt size) to t    16	the nearest 17	Pound Months inch? 18 □ 19	s OR □ 20 □		
28. W 29. W 30. W 31. W	that is your height? Feet that is your current weight (to the nearesthen was your weight last measured? What is your neck circumference (e.g. shings of the sentence that best describes your neck the sentence that best describes you have remained about the same weight last measured? I have gained 5 or more pounds in the Approximately how many pounds of 5-10 11-15	st pound)?  ithin the la  rt size) to t  i	the nearest 17	Pound Months inch? 18 19 ds) over the	s OR □ 20 □		
28. W 29. W 30. W 31. W	that is your height? Feet hat is your current weight (to the nearest hen was your weight last measured? What is your neck circumference (e.g. shill should be should b	st pound)?  ithin the la  rt size) to t  i	the nearest 17	Pound Months inch? 18 19 ds) over the	s OR		
28. W 29. W 30. W 31. W	that is your height? Feet that is your current weight (to the nearesthen was your weight last measured? What is your neck circumference (e.g. shings of the sentence that best describes your neck the sentence that best describes you have remained about the same weight last measured? I have gained 5 or more pounds in the Approximately how many pounds of 5-10 11-15	st pound)?  ithin the la  rt size) to t  i	the nearest 17	Pound Months inch? 18 19 ds) over the	s OR		

Has a physician diagnosed you with any of the following disorders (checles psychiatric:	
A major mood disorder (e.g. depression)	
A major anxiety disorder (e.g. panic disorder)	
Sleep Disorders:	
□ Narcolepsy	
☐ Restless legs syndrome	
Periodic Limb Movements	
Respiratory:	
☐ Asthma	
☐ Emphysema	
☐ Chronic Bronchitis	
Cardiovascular:	
☐ Hypertension	
☐ Myocardial infarction (heart attack)	
☐ Cardiac arrythmias (irregular heartbeat)	
☐ Angina	
☐ Congestive heart failure	
Endocrine:	
□ Diabetes	
☐ Hypothyroidism (low thyroid level)	
☐ Polycystic ovarian disease	
Neurology:	
☐ Stroke	
□ TIA	
☐ Seizure	
Have you been previously diagnosed with obstructive sleep apnea by a ph	nysician?
☐ Yes ☐ No	
If YES, please answer the following questions.  If NO, please advance to question 35.	
a. Was this diagnosis based on a full sleep study in which you slepi	t in the sleep laboratory overnight?
□ Yes □ No	
b. Was this diagnosis based on a home oximetry study (i.e. where	you wore a probe on your finger overnight)
☐ Yes ☐ No	
	- have a she diagnosis
c. If you answered NO to the above two questions, please describe	e now your physician made the diagnosis

																					. 1
☐ Positiv	ve airway pres	ssure (e	e.g. C	PAP,	BIPA	AP, sr	nart (	CPA	P)												
	appliance																				
	(please descr	ibe)																			
П	TIT	TI	T																		
35. Have you had	d surgery spe	cifically	to tre	eat yo	ur sle	ep a	pnea	or s	no	ring?											
ΠVaa	<u> </u>							19 P. S													
☐ Yes	If YES	, please hat kind	e ans	wer ti	ne fol	lowing	g que	stio	n: Ma	rk al	l that	t ar	nolv)								
□No				July	C. y C	ia you	a nav	٠. ر				٠ -	ر ر. ۳								
	sinu	s surge	ry																		
	☐ lase	r surge	ry of t	the pa	late/	throat															
	□ som	noplast	ty (rac	diofre	quen	cy abl	lation	) of t	the	tong	jue										
	☐ form	nal uvule	opala	topha	ryng	oplast	y (UF	PP	) ur	nder	gene	eral	ane	sthe	sia	in	the	op	erat	ina	FOO
		er. Pleas				$\Box$	ĹΤ	Ť	_			_	Т	Т	Т	7	_	Ė	1	T	
													1			-			1	1	
Over the last 24 Any type of the									g a	iđs)											
Any type of the 36. Prescription n	following to								g a	iđs)											
Any type of the 36. Prescription n	following to nedication:	get to	or st	ay as					g a	ids)											
Any type of the 36. Prescription no never very rare	following to nedication: ely (less than	get to	or st	ay as	leep	? (i.e.	. slee	pinę													
Any type of the 36. Prescription none rever very rare	following to nedication: ely (less than ess than one	get to once ponight pe	or st	ay as onth) ek bu	leep	? (i.e.	. slee	pinę			)										
Any type of the 36. Prescription non never very rare rarely (le	following to nedication: ely (less than ess than one les (1-2 night	once ponight pe	or st er mo er wee	enth) ek bu	leep	? (i.e.	. slee	pinę			)										
Any type of the 36. Prescription non-confidence of the 36. Prescription non-confidence of the 36. Prescription of the 36. Pres	following to nedication: ely (less than ess than one	once ponight person or days	er moer wee	enth) ek bur ek) k)	leep	? (i.e.	. slee	pinę			)										
Any type of the 36. Prescription non-compared very rare rarely (let sometime frequenting almost a	following to nedication: ely (less than ess than one les (1-2 nights ly (3-4 nights lways (5-7 nig	once prinight person days or days ghts or d	er moer wee	enth) ek bur ek) k)	leep	? (i.e.	. slee	pinę			)							7			
Any type of the 36. Prescription non-compared very rare rarely (let sometime frequenting almost a	following to nedication: ely (less than ess than one les (1-2 nights	once prinight person days or days ghts or d	er moer wee	enth) ek bur ek) k)	leep	? (i.e.	. slee	pinę			)	I	I	I			I	]			
Any type of the 36. Prescription non-compared very rare rarely (let sometime frequenting almost a	following to nedication: ely (less than ess than one les (1-2 nights ly (3-4 nights lways (5-7 nig	once prinight person days or days ghts or d	er moer wee	enth) ek bur ek) k)	leep	? (i.e.	. slee	pinę			)	I	I	<u> </u>	<u> </u>		Γ	]			
Any type of the 36. Prescription in never very rare rarely (le sometim frequent almost a	following to nedication: ely (less than ess than one les (1-2 nights ly (3-4 nights lways (5-7 nights	once ponight person days	er moer wee	enth) ek bur ek) k)	leep	? (i.e.	. slee	pinę			)	I	<u> </u>	1	1		Ι	]			
Any type of the 36. Prescription in never very rare rarely (le sometim frequent almost a	following to nedication: ely (less than ess than one les (1-2 nights ly (3-4 nights lways (5-7 nights	once ponight person days	er moer wee	enth) ek bur ek) k)	leep	? (i.e.	. slee	pinę			)	<u> </u>	I	1	1		Ι	]			
Any type of the 36. Prescription in never rarely (le sometime frequent almost a Name of street rarely (le le sometime frequent almost a never very rare	following to nedication: ely (less than ess than one less (1-2 nights ly (3-4 nights llways (5-7 nights f medication:	once ponight person days ghts or days	er mo	ay as onth) ek bur ek) k) week	t more	? (i.e.	once	ping	er m	oonth	Ι	I	I	I	1		Γ	]			
Any type of the 36. Prescription in never rarely (le sometime lamost a Name of 37. Non-prescriptic rarely (les rar	following to nedication: ely (less than ess than one i les (1-2 nights dy (3-4 nights dways (5-7 nights) f medication: on medication	once penight person or days ghts or days	er mo	onth) ek bur ek) k) week	t more	? (i.e.	once	ping	er m	oonth	Ι	I	<u> </u>	I	1		Ι	]			
Any type of the 36. Prescription in ever every rare frequent almost a Name of 37. Non-prescription in ever every rare rarely (less sometimes) sometimes	following to nedication: ely (less than ess than one les (1-2 nights ly (3-4 nights lways (5-7 nights f medication: on medication ely (less than one ss than one nies (1-2 nights	once prinight personned prince personned perso	er mo	onth) ek bur ek) k) week  muth) kbut	t more	? (i.e.	once	ping	er m	oonth	Ι	I	1	1	I		Γ	]			
Any type of the 36. Prescription in ever every rare frequent almost a Name of 37. Non-prescription ever every rare rarely (le sometime frequent) frequent frequent frequent frequent frequent frequent every frequent frequent frequent frequent every frequent	following to nedication: ely (less than ess than one les (1-2 nights ly (3-4 nights lways (5-7 nights f medication: on medication ly (less than one ss than one nies (1-2 nights y (3-4 nights of	once prinight personned pe	er mo	onth) ek bur ek) k) week inth) kbut kk)	t more	? (i.e.	once	ping	er m	oonth	Ι	I	I	I			Γ	]			
Any type of the 36. Prescription in ever every rare frequent almost a Name of 37. Non-prescription ever every rare rarely (le sometime frequent) frequent frequent frequent frequent frequent frequent every frequent frequent frequent frequent every frequent	following to nedication: ely (less than ess than one les (1-2 nights ly (3-4 nights lways (5-7 nights f medication: on medication ely (less than one ss than one nies (1-2 nights	once prinight personned pe	er mo	onth) ek bur ek) k) week inth) kbut kk)	t more	? (i.e.	once	ping	er m	oonth	Ι	I	I	1			Γ	]			

						D		·DA	D)												
☐ Positive a		sure (e.	g. CPA	P, B	IPA	P, sn	nan C	PA	(۲												
☐ dental ap																					
other (ple	ase descril	oe)																			
						882	105														
		1-1						لـــ													
35. Have you had su	irgery spec	ifically t	to treat	your	sle	ep ap	onea	or s	noi	ing?	,										
□Yes ⊏	) IF VEC	please	000140	r tha	foll	ouin		ctio	٠.												
	Wh	at kind(	answei s) of su	ırger	y di	d you	ı have	suo ∋? (I	n. Ma	rk al	l tha	t a	pply	)							
L NO		- Louis de la company																			
	sinus																				
	☐ laser	surgery	y of the	pala	te/ti	hroat															
	☐ somr	oplasty	(radiof	requ	enc	y abl	ation	of t	he	tong	gue										
	☐ forma	al uvulo	palatop	hary	ngo	plast	y (UP	PP)	ur	der	gene	era	l an	esth	esia	in	the	ор	erat	ing	FOC
	☐ other	. Please	e descri	be:				T				Г	T	T	Т				T	1	
												ŀ									
Over the last 24 mo	onths, on								1 2	ide)	L		1		701			L	1		
Any type of the following 36. Prescription med	onths, on								g a	ids)	<b>L</b>	-	•	1				I_	I		
Any type of the foll 36. Prescription med	onths, on a lowing to lication:	get to c	or stay	asle					ga	ids)								I_		ı	
Any type of the foll 36. Prescription med never very rarely	onths, on lowing to lication:	get to d	or stay	asle	ep?	? (i.e.	slee	ping					1					I_			
Any type of the foll 36. Prescription med never very rarely rarely (less	onths, on lowing to lication:  (less than one means that the means the means the means that the means that the means the	get to conce pe	or stay	asle  i)  out m	ep?	? (i.e.	slee	ping					1					L	<u>I</u>		
Any type of the folion 36. Prescription median never very rarely rarely (less sometimes	onths, on a lowing to lication:  (less than one in (1-2 nights)	get to conce peright period	or stay or month r week t s/week)	asle  i)  out m	ep?	? (i.e.	slee	ping						<b>.</b>				I		1	
Any type of the folion 36. Prescription median never very rarely rarely (less sometimes frequently (see the folion).	onths, on a lowing to lication: (less than or than one no (1-2 nights of 3-4 nights of 5-4 nights of	get to conce per ight per or days	or stay or month r week to s/week)	asle  i)  out m	ep?	? (i.e.	slee	ping					•					I	1	I	
Any type of the folional and the foliona	onths, on lowing to lication: (less than of than one modern incomplete incomp	get to conce per ight per or days	or stay or month r week to s/week)	asle  i)  out m	ep?	? (i.e.	slee	ping				_					_	I	1	I	
Any type of the foll 36. Prescription med never very rarely rarely (less sometimes frequently (:	onths, on lowing to lication: (less than of than one modern incomplete incomp	get to conce per ight per or days	or stay or month r week to s/week)	asle  i)  out m	ep?	? (i.e.	slee	ping				I					I	]	1	L	
Any type of the foli 36. Prescription med never very rarely rarely (less sometimes frequently (including the content of the co	onths, on lowing to lication: (less than of than one modern incomplete incomp	get to conce per ight per or days	or stay or month r week to s/week)	asle  i)  out m	ep?	? (i.e.	slee	ping				1					I	]	1		
Any type of the foll 36. Prescription med never very rarely rarely (less sometimes frequently (including almost always) Name of med	lowing to lication: (less than or than one no (1-2 nights 3-4 nights 6 bys (5-7 nights dedication:	get to conce pe ight per or days/ hts or d	or stay or month r week to s/week)	asle  i)  out m	ep?	? (i.e.	slee	ping				I					I	]	1	1	
Any type of the foll 36. Prescription med never very rarely rarely (less sometimes frequently (i) almost alway Name of med 37. Non-prescription never	lowing to dication: (less than or than one in (1-2 nights or less (5-7 nigredication:	ponce pe ight per or days or days/ hts or d	or stay	asle  i)  out m	ep?	? (i.e.	slee	ping				<u> </u>					I	]		1	
Any type of the folion 36. Prescription medians are left in the folion and in the fo	lowing to lication: (less than or than one in (1-2 nights and a lication: (5-7 nights displayed (5-7 nights)) (bedication:	pnce peright per or days/hts or d	or stay	asle  i)  ek)	ep?	? (i.e.	slee	ping	r m	onth	)	<u> </u>					I	]		1	
Any type of the folion 36. Prescription medians are left in the folion and in the fo	lowing to lication: (less than of than one in less than of the ledication:  medication:	pnce peright per or days/hts or d	or stay  or month r week t s/week) days/wee month week b	asle  i)  ek)	ep?	? (i.e.	slee	ping	r m	onth	)	<u> </u>					Ι	]			
Any type of the folion 36. Prescription median rever very rarely (less sometimes frequently (including almost always).  37. Non-prescription never very rarely (less sometimes	lowing to lication: (less than or than one in less than or han one ni 1-2 nights or han one ni 1-2 nights	pnce peright per or days/hts or days/nce per ght per or days/	or stay  or month r week t s/week) ays/wee month week b	asle  i)  ek)	ep?	? (i.e.	slee	ping	r m	onth	)	<u> </u>					Ι	]			
Any type of the folion 36. Prescription median in ever rarely (less sometimes frequently (in almost always) almost always.  37. Non-prescription rever rarely (less sometimes (in proper sometimes (in prequently (in prepared (in prepa	lowing to lication: (less than of than one in the less than of the less than of than one in the less than of the less than	pnce per days/hts or days/r days/r days/r days/r days/r days/r days/v	r month week) week) week) month week b week)	ek)	ep?	? (i.e.	slee	ping	r m	onth	)	1					Ι	]			
Any type of the folion 36. Prescription median rever very rarely (less sometimes frequently (including almost always).  37. Non-prescription never very rarely (less sometimes	lowing to lication: (less than of than one in that of the than one in than one in that of the than one in the than	pnce per days/hts or days/r days/r days/r days/r days/r days/r days/v	r month week) week) week) month week b week)	ek)	ep?	? (i.e.	slee	ping	r m	onth	)	I					I	]			



	u. dication
8. Herbal or health food	medication
never	s than once per month)
- July (less than	n one night per week but more than once per month)
times (1-2	nights of days/week)
c contly (3-4 f	nights or days/week)
I frequency (	(5-7 nights or days/week)
Name of medic	;ation:
grange de la companya	ns, on average how often did you take:
Over the last 24 month	ing to stay awake? (i.e. etimulants)
Any type of the followi	ing to stay awake? (i.e. stimulants)
	tion: (eg. Modafinil, Ritalin)
never	ss than once per month)
very rarely (less	an one night per week but more than once per month)
rarely (less that	In one hight per week but more than once per means,
sometimes (1-2	2 nights or days/week)
frequently (3-4	nights or days/week)
almost always	(5-7 nights or days/week)
Name of medic	cation:
40 Non-prescription me	edication (eg. caffeine pills)
□ never	
	ss than once per month)
□ rarely (less tha	an one night per week but more than once per month)
☐ sometimes (1-	-2 nights or days/week)
☐ frequently (3-4	4 nights or days/week)
□ almost always	s (5-7 nights or days/week)
Name of medi	
Over the last 24 month	ths, on average how often did you take:
	medications (for any reason)
41. Anti-histamines	
never	
	ess than once per month)
D very rarely (le	than once per morning
very rarely (less that	an one night per week but more train over
rarely (less that	ess than once per month) nan one night per week but more than once per month) 1-2 nights or days/week)
☐ rarely (less that ☐ sometimes (1-	nan one night per week but more trial of the last of t

4	n . 4 . 4 . 4 . 4 . 4 . 4 . 4 . 4 . 4 .						
	2. Antidepressant						
	never never						
	very rarely (less than once per month) rarely (less than one night per week but more to	han ond	ce per mor	ith)			
	rarely (less than one night per week but more to	nun on					
	☐ sometimes (1-2 nights or days/week)						
	☐ frequently (3-4 nights or days/week) ☐ almost always (5-7 nights or days/week)						
4	3. Other medications: (please list)						
			3 - 9		A Paragonia	_	
						_	
	Questions:						
4	<ol> <li>On average over the last 24 months, how many kilom</li> </ol>	eters d	id you drive	e per week	?	kilo	meter
45	5. Over the last 24 months, did <u>vou</u> have any motor vehi	cle acc	idents or c	rashes in v	vhich you	were drivin	g?
	☐ Yes ☐ No						
	If YES, please answer the following cluster of que: If NO, please skip to question 47.	stions.					
	How many crashes did you have over the last 24	0	1	2	3	4	5
•	now many crashes did you have over the last 24			_		_	
a.	months?				·		
				0	0		
b.	months?						E
b.	months?  In how many of these, if any, was a police report filed?  In how many of these, if any, do you think sleepiness or fatigue in you contributed?  In how many of these, if any, was total property	_ 	0	0	0	0	
b. c. d.	months? In how many of these, if any, was a police report filed? In how many of these, if any, do you think sleepiness or fatigue in you contributed? In how many of these, if any, was total property damage greater than \$1000?					0	<u> </u>
b. c. d.	months?  In how many of these, if any, was a police report filed?  In how many of these, if any, do you think sleepiness or fatigue in you contributed?  In how many of these, if any, was total property	_ 	0	0	0	0	0
b. c. d. e.	months?  In how many of these, if any, was a police report filed?  In how many of these, if any, do you think sleepiness or fatigue in you contributed?  In how many of these, if any, was total property damage greater than \$1000?  In how many of these, if any, were any vehicles involved towed away?  In how many of these, if any, were emergency room		0	0	0	0	E
b. c. d. e.	months?  In how many of these, if any, was a police report filed?  In how many of these, if any, do you think sleepiness or fatigue in you contributed?  In how many of these, if any, was total property damage greater than \$1000?  In how many of these, if any, were any vehicles involved towed away?		0	0	0	0	0

over the last two weeks, how often have you been bothered b	Not at	onowing pro	blems?	
	all	several days	more than half the days	nearly every
47. Little interest or pleasure in doing things?	0	0	0	day
s seeling down, depressed, or hopeless?		0		
Trouble falling/staying asleep, sleeping too much?		0		
50. Feeling tired or having little energy?	0	0		
51. Poor appetite or overeating?	0		0	0
51. Poor appeare or over the type are a failure		0	0	0
52. Feeling bad about yourself, or that you are a failure, or have let yourself or your family down?	0	מ	0	0
53. Trouble concentrating on things, such as reading the newspaper or watching TV	0	۵	۵	0
54. Moving or speaking so slowly that other people could have noticed. Or the opposite; being so fidgety or restless that you have been moving around more than usual.	0	0	0	0
55. Thoughts that you would be better off dead or of hurting yourself in some way.			0	0
55. Thoughts that you would be better off dead or of hurting yourself in some way.  56. If you have checked off any problem on these questions (for you to do your work, take care of things at home, or get all \( \subseteq \text{Not difficult} at all \( \subseteq \text{very difficult} \) very difficult \( \subseteq \text{ extremely difficult} \)	47-55), how	difficult have		
yourself in some way.  56. If you have checked off any problem on these questions (for you to do your work, take care of things at home, or get all  \text{\tex{\tex	47-55), how	difficult have		
yourself in some way.  56. If you have checked off any problem on these questions (for you to do your work, take care of things at home, or get all Not difficult at all somewhat difficult very difficult extremely difficult	47-55), how	difficult have		
yourself in some way.  56. If you have checked off any problem on these questions (for you to do your work, take care of things at home, or get all Not difficult at all somewhat difficult very difficult extremely difficult extremely difficult  57. Women Only: What is your menopausal status:  1 Premenopausal  1 If premenopausal: Are your periods  1 regular	47-55), how ong with oth	difficult have er people?		s made it



Thank you very much for completing the survey. Your information is very important to us and we really appreciate your time and consideration.



