

**Table 2: Proportion of SLE pregnancies exposed to medications during specific windows before, during, and after pregnancy**

Medication	Before pregnancy			During pregnancy			After pregnancy	
	BC1	BC2	Combined	T1	T2	T3	Combined	PP
<b>Antimalarials</b>	114 (30.3%)	114 (30.3%)	136 (36.2%)	72 (19.1%)	63 (16.7%)	64 (17.0%)	91 (24.3)	117 (31.1%)
<b>Immunosuppressants</b>								
Azathioprine	26 (6.9%)	41 (10.9%)	44 (11.7%)	25 (6.6%)	24 (6.4%)	26 (6.9%)	31 (8.2%)	39 (10.4%)
Methotrexate	10 (2.7%)	7 (1.9%)	13 (3.5%)	*	0	0	*	10 (2.7%)
Mycophenolate	11 (2.9%)	8 (2.1%)	12 (3.2%)	0	0	0	0	*
Minocycline	7 (1.9%)	*	11 (2.9%)	0	0	0	0	*
Cyclosporine	*	*	*	*	*	*	*	*
Cyclophosphamide	*	0	*	0	0	0	0	0
Leflunomide	*	0	*	0	0	0	0	0
Sulfasalazine	*	*	*	*	*	*	*	*
<b>Other medications</b>								
Glucocorticosteroids	101 (26.9%)	89 (23.7%)	125 (33.2%)	50 (13.3%)	56 (14.9%)	50 (13.3%)	74 (19.7%)	95 (25.3%)
NSAIDs	91 (24.2%)	84 (22.3)	131 (34.8%)	16 (4.3%)	*	*	18 (4.8%)	72 (19.1%)
Low dose aspirin	*	*	*	*	*	*	*	*

\* Cell counts  $\leq 6$ , suppressed as per data use agreement with Popdata BC

Abbreviations: BC1 – 24-12 months before conception

BC2 – 12 months before conception

T1 – first pregnancy trimester

T2- second pregnancy trimester

T3- third pregnancy trimester

PP- first year post partum

NSAID- Nonsteroidal anti-inflammatory drug