The University of British Columbia

FACULTY OF GRADUATE STUDIES

PROGRAMME OF THE

FINAL ORAL EXAMINATION.

FOR THE DEGREE OF

DOCTOR OF PHILOSOPHY

of

EWART WILLIAM BLACKMORE

B.Sc.(Eng.), Queen's University, 1963

M.Sc., University of British Columbia, 1965

THURSDAY, AUGUST 17, 1967 AT 3:30 P.M.

IN ROOM 301, HENNINGS BUILDING

COMMITTEE IN CHARGE

Chairman: I. McT. Cowan

J.B. Warren

G.M. Griffiths

B.L. White

E. Vogt

M. McMillan

F.K. Bowers

Research Supervisor: J.B. Warren

External Examiner: T.A. Tombrello

Department of Physics

California Institute of Technology

Pasadena

California

U.S.A.

FINAL STATE INTERACTIONS IN THE REACTIONS

$\text{He}^3(\text{He}^3, 2\text{p})\text{He}^4$ and $\text{T}(\text{He}^3, \text{np})\text{He}^4$.

ABSTRACT

Final state interactions in the p-p and p-n systems corresponding to the formation of the singlet states of the diproton and the deuteron have been observed in the reactions $\text{He}^3(\text{He}^3, 2\text{p})\text{He}^4$ and $\text{T}(\text{He}^3, \text{np})\text{He}^4$, by measuring the energy spectra of two of the final state particles detected in coincidence. These interactions appear as regions of high density of events along the kinematic contours in two dimensional energy space. Both p-p and p- He^4 coincidences from the reaction $\text{He}^3(\text{He}^3, 2\text{p})\text{He}^4$ and p- He^4 coincidences from the reaction $\text{T}(\text{He}^3, \text{np})\text{He}^4$ were studied. The dominant mechanism producing the three particle final states was found to be the sequential breakup through the states of Li^5 and He^5 , in agreement with the results of previous experiments. The interaction between the two nucleons affects the distribution of coincidence events only in the region of two dimensional energy space corresponding to a breakup in which the two nucleons emerge with low relative energy. The relative contributions of the different processes in the reaction $\text{He}^3(\text{He}^3, 2\text{p})\text{He}^4$ was determined for incident He^3 energies from 1.15 MeV to 5.0 MeV. It was found that the nucleon-nucleon interaction becomes less important as the incident energy increases. The validity

of two approximate theories of final state interactions, the Watson theory and the Phillips, Griffy and Biedenharn density of states formalism, was checked by comparing their predictions with the observed energy spectra.

GRADUATE STUDIES

Field of Study: Experimental Nuclear Physics

Nuclear Physics	E. Vogt, J.B. Warren
Theoretical Nuclear Physics	M. McMillan
Cosmic Rays and High Energy Physics	J.B. Warren
Elementary Quantum Mechanics	F.A. Kaempffer
Electronic Instrumentation	F.K. Bowers
Electromagnetic Theory	G. Volkoff
Digital Computer Programming	H. Dempster
Low Temperature Physics	J.B. Brown
Group Theory Methods in Quantum Mechanics	W. Opechowski

PUBLICATIONS AND PAPERS

E.W. Blackmore and J.B. Warren. "Final State Interactions in the Reaction $\text{He}^3(\text{He}^3, 2p)\text{He}^4$ ". Phys. Rev. Letters 16, 520 (1966).

R.J.A. Levesque, R.W. Ollerhead, E.W. Blackmore and J.A. Kuehner. "Properties of Levels at 6.69 and 6.88-6.89 MeV in Si^{28} ". Can. J. Phys. 44, 1087 (1966).

E.W. Blackmore and J.B. Warren. "Final State Interactions in the Reactions $\text{He}^3(\text{He}^3, 2p)\text{He}^4$ and $\text{T}(\text{He}^3, np)\text{He}^4$ ". Can. J. Phys. (submitted).

FINAL STATE INTERACTIONS IN THE REACTIONS

$\text{He}^3(\text{He}^3, 2p)\text{He}^4$ AND $\text{T}(\text{He}^3, np)\text{He}^4$

by

EWART WILLIAM BLACKMORE

B.Sc.(Eng) Queen's University, Kingston, 1963

M.Sc., University of British Columbia, 1965

A THESIS SUBMITTED IN PARTIAL FULFILMENT
OF THE REQUIREMENTS FOR THE DEGREE OF

DOCTOR OF PHILOSOPHY

in the department of

PHYSICS

We accept this thesis as conforming to the
required standard

THE UNIVERSITY OF BRITISH COLUMBIA

July 1967

In presenting this thesis in partial fulfilment of the requirements for an advanced degree at the University of British Columbia, I agree that the Library shall make it freely available for reference and Study. I further agree that permission for extensive copying of this thesis for scholarly purposes may be granted by the Head of my Department or by his representatives. It is understood that copying or publication of this thesis for financial gain shall not be allowed without my written permission.

Department of PHYSICS

The University of British Columbia
Vancouver 8, Canada

Date Aug 17, 1967

ABSTRACT

Final state interactions in the p-p and p-n systems corresponding to the formation of the singlet states of the di-proton and the deuteron have been observed in the reactions $\text{He}^3(\text{He}^3, 2p)\text{He}^4$ and $\text{T}(\text{He}^3, np)\text{He}^4$, by measuring the energy spectra of two of the final state particles detected in coincidence. These interactions appear as regions of high density of events along the kinematic contours in two dimensional energy space. Both p-p and p- He^4 coincidences from the reaction $\text{He}^3(\text{He}^3, 2p)\text{He}^4$ and p- He^4 coincidences from the reaction $\text{T}(\text{He}^3, np)\text{He}^4$ were studied. The dominant mechanism producing the three particle final states was found to be the sequential breakup through the states of Li^5 and He^5 , in agreement with the results of previous experiments. The interaction between the two nucleons affects the distribution of coincidence events only in the region of two dimensional energy space corresponding to a breakup in which the two nucleons emerge with low relative energy. The relative contributions of the different processes in the reaction $\text{He}^3(\text{He}^3, 2p)\text{He}^4$ was determined for incident He^3 energies from 1.15 MeV to 5.0 MeV. It was found that the nucleon-nucleon interaction becomes less important as the incident energy increases. The validity of two approximate theories of final state interactions, the Watson theory and the Phillips, Griffy and Biedenharn density of states formalism, was checked by comparing their predictions with the observed energy spectra.

TABLE OF CONTENTS

		<u>Page</u>
CHAPTER I	INTRODUCTION	1
	A. General Introduction	1
	B. Review of Previous Work	8
	C. Present Work	13
CHAPTER II	KINEMATICS AND EXPERIMENTAL TECHNIQUE	16
	A. Kinematics of three particle breakup reactions	16
	B. Experimental Technique	20
CHAPTER III	EXPERIMENTAL ARRANGEMENT	24
	A. Introduction	24
	B. Geometry #1 Target Chamber	27
	C. Geometry #2 Target Chamber	29
	D. Tritium Handling System	31
	E. Charged Particle Detectors	33
	F. Electronics	34
	G. On-line Computer Techniques	36
CHAPTER IV	EXPERIMENTAL PROCEDURE AND RESULTS	40
	A. Geometry #1 Measurements	40
	B. Geometry #2 Measurements	43
	C. The $\text{He}^3(\text{He}^3, 2p)\text{He}^4$ Reaction	44
	D. The $\text{T}(\text{He}^3, np)\text{He}^4$ Reaction	47
CHAPTER V	THEORY OF FINAL STATE INTERACTIONS	51
	A. Introduction	51
	B. Analysis of Coincidence Proton Spectra	57
	C. Geometry #1 Analysis of Results	63
	D. Geometry #2 Analysis of Results	69

<u>Contents continued</u>		<u>Page</u>
CHAPTER VI	DISCUSSION	73
	A. Present Results	73
	B. Future Work	75
APPENDIX A	REACTION KINEMATICS FOR THREE FINAL PARTICLES	78
APPENDIX B	COMPUTER PROGRAM LISTINGS	87
APPENDIX C	SEMI-EMPIRICAL EXPRESSION FOR ENERGY LOSS IN THIN FOILS	89
REFERENCES		92

LIST OF FIGURES

To follow
page

Figure 1.	Relationship between velocities in lab. and scm. coordinate systems.	17
2.	Relationship between velocities in scm. and rcm(1) coordinate systems.	17
3.	Kinematic contours for p-p and p-He ⁴ coincidences from the reaction He ³ (He ³ ,2p)He ⁴ .	20
4.	Kinematic contours for p-He ⁴ coincidences from the reaction He ³ (He ³ ,2p)He ⁴ .	23
5.	Schematic diagram of Geometry #1 and Geometry #2.	26
6.	Kinematic contours for p-p, p-He ⁴ and He ⁴ -p coincidences from the reaction He ³ (He ³ ,2p)He ⁴ .	26
7.	Kinematic contours for p-p coincidences from the reaction He ³ (He ³ ,2p)He ⁴ .	26
8.	Geometry #1 target-detector arrangement.	27
9.	Geometry #2 target-detector arrangement (side-view).	29
10.	Geometry #2 target-detector arrangement (rear-view).	29
11.	Diagram of pumping and gas handling system.	31
12.	Block diagram of electronics used in on-line data acquisition at Chalk River.	35
13.	Block diagram of electronics used in data acquisition at U.B.C.	35
14.	Time distribution spectrum of coincidences from the reaction He ³ (He ³ ,2p)He ⁴ .	35
15.	Two dimensional energy spectrum taken with Geometry #1 and $\Delta_{12} = 6^\circ$.	43

Figures continued

to follow
page

Figure 16.	Coincidence proton spectra obtained by summing events along appropriate contour.	43
17.	Coincidence proton spectra obtained by summing events along appropriate contour.	43
18.	Coincidence proton spectra obtained by summing events along appropriate contour.	43
19.	Coincidence proton spectra obtained by summing events along appropriate contour.	43
20.	Coincidence proton spectra obtained by summing events along appropriate contour.	43
21.	Coincidence proton spectra obtained by summing events along appropriate contour.	43
22.	Single particle spectra in both detectors from the reaction $\text{He}(\text{He}, 2\text{p})\text{He}$ at 90° to the incident beam direction.	44
23.	Two dimensional contour plot of p-p and p- He^4 coincidence events from the reaction $\text{He}^3(\text{He}^3, 2\text{p})\text{He}^4$ at $\Delta_{12} = 152^\circ$.	45
24.	Coincidence proton spectrum from the reaction $\text{He}^3(\text{He}^3, 2\text{p})\text{He}^4$ obtained by summing over p- He^4 contour for $\Delta_{12} = 152^\circ$.	46
25.	Coincidence proton spectrum from the reaction $\text{He}^3(\text{He}^3, 2\text{p})\text{He}^4$ obtained by summing over p- He^4 contour for $\Delta_{12} = 165^\circ$.	46
26.	Single particle spectra in both detectors from the reaction $\text{T}(\text{He}^3, \text{np})\text{He}^4$ at 90° to the incident beam direction.	47

Figures continued

to follow
page

- Figure 27. Two dimensional contour plot of p-He⁴ and d-He⁴ coincidence events from the reaction T(He³,np)He⁴ at $\Delta_{12} = 151.4^\circ$. 48
28. Coincidence proton spectrum from the reaction T(He³,np)He⁴ obtained by summing over p-He⁴ contour for $\Delta_{12} = 151.4^\circ$. 49
29. Coincidence proton spectrum from the reaction T(He³,np)He⁴ obtained by summing over p-He⁴ contour for $\Delta_{12} = 165^\circ$. 49
30. Excitation functions for states of Li⁵ predicted by Watson and PBG theories. 62
31. Excitation functions for states of He⁵ predicted by Watson and PBG theories. 62
32. Excitation functions for singlet state of p-p system predicted by Watson and PBG theories. 62
33. Excitation functions for singlet state of n-p system predicted by Watson and PBG theories. 62
34. Comparison of experimental coincidence proton spectra with Watson and PBG theories for $T_0 = 1.15$ MeV. 64
35. Comparison of experimental coincidence proton spectrum with Watson and PBG theories for $T_0 = 5.00$ MeV. 64
36. Comparison of predictions of statistical breakup with Watson and PBG theories for Li⁵ excited state. 66
37. Comparison of experimental coincidence proton spectra at small angles with Watson and PBG theories. 67

Figures continued

to follow
page

Figure 38.	Coincidence proton spectrum from the reaction $\text{He}^3(\text{He}^3, 2\text{p})\text{He}^4$ for $\Delta_{12} = 152^\circ$ divided by the available phase-space.	70
39.	Comparison of experimental coincidence proton spectra with Watson theory for the p-p final state interaction.	71
40.	Coincidence proton spectrum from the reaction $\text{T}(\text{He}, 2\text{p})\text{He}$ for $\Delta_{12} = 151.4^\circ$ divided by the available phase-space.	71
41.	Comparison of experimental coincidence proton spectra with Watson theory for the n-p final state interaction.	71
A1.	Relationship between velocities in lab. and scm. coordinate systems.	79
A2.	Relationship between velocities in scm. and rcm(i) coordinate systems.	79
C1.	Energy loss curves for protons and deuterons in nickel foils.	91
C2.	Energy loss curves for He^3 and He^4 in nickel foils.	91

LIST OF TABLES

	<u>To follow page</u>
Table 1. Symbols used in describing a three particle breakup.	16
2. Commercial electronic units used in the experimental arrangement.	35
3. Detector configurations used with Geometry #1 measurements.	41
4. Parameters obtained for states of Li^5 and He^5 from single level dispersion theory.	59
5. Effective range theory parameters for nucleon-nucleon scattering phase shifts.	59

ACKNOWLEDGEMENTS

I am deeply grateful to my supervisor, Dr. J. B. Warren, for his interest and encouragement throughout the course of this work. The assistance of Drs. B. L. White and G.M. Griffiths who acted as interim supervisors in Dr. Warren's absence is also greatly appreciated.

The many discussions with Dr. Erich Vogt on some of the theoretical problems have been a tremendous help. The willing assistance of the members of Van de Graaff group, both faculty, students and technicians is acknowledged with thanks.

I wish to thank the tandem group at the Chalk River Nuclear Laboratories for making it possible to carry out some of the experimental measurements at higher energies.

Finally, I am indebted to the National Research Council for a Bursary and three Studentships received during my graduate work and for their continued financial support in postdoctoral studies.

CHAPTER I
INTRODUCTION

A. General Introduction

Nuclear reactions in which three or more particles are produced in the final state have recently become the subject of a great deal of interest both experimentally and theoretically. This has come about as a result of the development of multi-dimensional pulse height analysers and on-line computers which has made possible useful experimental studies of these reactions. Although these reactions are very complicated in comparison with the usual nuclear reaction with two particles in both the initial and final states, and hence the need for more sophisticated analysing equipment, there are a number of advantages to be gained in their investigation. Before enumerating these advantages it is worthwhile to discuss some of the present aims of nuclear research at low energies.

The fundamental problem of nuclear physics is to understand the forces acting between the nuclear particles. In the investigation of this interaction early researchers followed the obvious course and studied intensively the simplest nuclear systems, those in which only two nucleons are interacting. They reasoned that a complete knowledge of the two nucleon systems would allow one to compute all nuclear properties, i.e. that the forces between nucleons are simply additive. Present evidence indicates that this may not be completely true and that the nuclear force may involve terms which arise only when

a third or fourth particle is present. These "many-body" forces have not been studied in great detail as yet due to their mathematical complexity and to the difficulty in designing experiments to determine their properties unambiguously. Since many features of nuclear behaviour are conveniently explained using two body forces it is expected that the study of this simple interaction will continue to be profitable for some time. In fact it will probably be necessary to obtain a better understanding of the two body interaction before any quantitative conclusions can be drawn about the role of many body forces in determining nuclear properties.

There are three nuclear systems consisting of two nucleons, the di-proton, the di-neutron, and the deuteron. The possible states of these nucleon-nucleon systems may be classified using the concepts of ordinary spin and isospin. The neutron-proton system in the S state can have its ordinary spins either parallel corresponding to a triplet spin state, ^3S , or antiparallel corresponding to a singlet spin state, ^1S . The two protons and two neutrons on the other hand can exist only with opposite spins as a result of the exclusion principle for identical particles and therefore must be in a singlet state ^1S . The singlet states of the three systems belong to the same isospin triplet $T=1$, while the triplet state of the deuteron is in the isospin singlet state $T=0$.

In general, simple scattering experiments have been used to uncover most of the properties of these systems. However, only the proton-proton and proton-neutron systems are

accessible with these scattering experiments due to the problem of obtaining a target of free neutrons. In these experiments a target of essentially free protons, such as hydrogen gas, is bombarded with a beam of protons from an accelerator, or neutrons from a reactor or neutron generator, and the scattering of the incoming particles is monitored. The quantities which are measured are the total cross section for scattering as a function of the relative nucleon-nucleon energy or the excitation function, the differential cross section or the probability that the incoming nucleon is scattered through a given angle, and in more recent experiments the polarization or preferred spin direction of the scattered nucleon. Information about the neutron-neutron system has been obtained indirectly from the scattering of neutrons on deuterons and from the reaction of negative pions on deuterons.

Of the possible two nucleon systems only the neutron-proton system is found to exist in a stable bound state, the ground state of the deuteron. Measurements of the magnetic moment of the deuteron indicate that it is predominantly in the triplet ^3S state. The singlet states belonging to the isospin triplet are not bound pointing out one property of the nuclear force, spin dependence.

If the nuclear interaction is described in terms of a simple potential it is found that the low energy scattering experiments cannot determine the shape of this potential but merely two quantities which describe its gross features, the scattering length 'a' and the effective range 'r'. The values of these parameters for the singlet states are presently reported

as follows:

n-p	p-p
$a_s = -23.71 \pm .07 \text{ fm.}$	$a_p = -7.72 \pm .04 \text{ fm.}$
$r_{os} = 2.4 \pm .3 \text{ fm.}$	$r_{op} = 2.72 \pm .10 \text{ fm.}$

The similarity of the two effective ranges suggests that the nuclear forces are charge independent. Further evidence for this is the fact that the concept of isospin, in which the neutron and proton are regarded as two almost degenerate states of the same particle, the nucleon, is extremely successful in predicting properties of mirror nuclei. However the question remains, how good is the approximation of charge independence. The scattering lengths for the two systems are quite different but most of this difference can be attributed to the Coulomb force in the case of the p-p system. Approximate calculations of the scattering length for two protons in the absence of the Coulomb force yields a value of -17 fm. The best estimates of the scattering length for the n-n system range from -17 fm. to -24 fm.

One property of the nuclear force which is very much in evidence in the vast amount of experimental information available is its extremely complicated nature. To explain one of the simplest properties of a nuclear system, the quadrupole moment of the deuteron, it is necessary to assume the presence of a non-central or tensor component in the nuclear force. Even more complications such as repulsive cores and velocity dependent interactions are used to explain the results of scattering

experiments carried out at higher energies.

It is not expected that the study of multiparticle breakup reactions will immediately solve this complicated situation. However this approach should allow a more accurate determination of the neutron-neutron scattering length and it offers the remote possibility of obtaining evidence for or against the presence of "many-body" forces. In addition to obtaining information about the nucleon-nucleon interaction, multiparticle breakup reactions may be used to probe the clustered nature of nuclei containing many nucleons. In high energy physics the analysis of reactions of this type has led to the discovery of many of the short-lived strange particle interactions, such as the pion resonances. As a result of this success, high energy physics has led the way in determining the complicated kinematic and phase space calculations applicable to multiparticle breakup reactions.

In low energy nuclear physics, three particle breakup reactions may be used to give similar information about the nucleon-nucleon interaction or any other particle-particle interaction as scattering experiments. Here the interaction appears as a "final state interaction" influencing the energy and angular distributions of the breakup particles. If the final state interaction is of sufficiently long duration then the breakup may also be considered as a series of two body decays. In this sequential breakup the product of the first decay is a virtual particle which subsequently decays into two stable particles.

The types of measurements used in the analysis of these reactions fall into two groups. The first group consists of

those measurements which are made by monitoring the reaction with only one detector. These include determinations of the total yield as a function of the incident energy, and the various energy distributions of the reaction products. These measurements tend to be ambiguous unless some sort of particle identification system is used. Reasons for this ambiguity are given in the discussion of the reaction kinematics in Chapter II.

The second class of measurements requires the use of two detectors to monitor the energies and directions of two of the final state particles detected in coincidence and hence from the same breakup. A detailed description of this technique requires a discussion of the reaction kinematics. In the next chapter it will be shown that by choosing appropriate polar and azimuthal angles of the two detectors it is possible to observe a particular interaction as a function of the relative energy of the corresponding two particles. Therefore with a fixed beam energy it is possible to determine the same sort of excitation function for two particles as one would obtain by studying the scattering between the same particles at various incident energies.

One advantage of the three particle breakup reaction is that it is possible to observe several final state interactions in a single experiment and therefore it is a simple matter to determine the relative intensities of the interactions. This, of course, would not be possible if the three particles in the final state were identical. In some cases this advantage might be a disadvantage if the contributions from the various interactions cannot be separated. However in most reactions it is possible to

choose detector configurations in which the interaction of interest dominates. Another property of these reactions is that the centre of mass motion of the intermediate virtual system can be used as a kind of "energy amplifier". This means that a range of relative or internal energies in the two particle system of a few hundred keV might correspond to a range of a few MeV in the laboratory energy of one of the breakup particles. This feature allows a more accurate determination of the excitation function especially at low excitation energies. From an experimental point of view, the fact that a coincidence experiment is being performed is useful in reducing the background from contaminant reactions.

At the present time the main problem faced in obtaining useful information from the experimental results is the mathematical complexity of the three body problem. Although there is considerable effort being made towards finding techniques to include the three body effects exactly, and some of these methods appear promising, the state of the existing theories is still not sufficiently advanced to extract nuclear parameters from the experimental results with much confidence. Several formalisms for final state interactions have been developed, each method being applicable only to a certain type of problem. A general discussion of this work is given in Chapter V. These formalisms have been partly successful in explaining certain features of the experimental results. However the present situation is to use the experimental results to check on the existing theories. Perhaps the most important result of studying the nucleon-nucleon interaction in this way is that it serves as one of the most stringent tests on

the theories.

With the development of a precise theory, otherwise inaccessible parameters such as the neutron-neutron singlet scattering length may be determined accurately. The theory should also be quite useful in high energy physics where the multiparticle breakup reaction is quite common and the interactions between the particles are in the process of being investigated.

B. Review of Previous Work

A general conclusion which can be drawn from the results of the experimental work to date is that final state interactions or their kinematic equivalent, sequential decays through intermediate states, play a dominant role in multiparticle breakups. This is indeed fortunate as a no nuclear information could be obtained otherwise. The other possible reaction mode is the direct or instantaneous breakup in which the energy and angular distributions of the reaction products depend only on the available phase-space. This type of breakup has a coincidence yield distribution characterized by a slow variation with particle energy and the presence of peaks at the maximum particle energies. The effect of final state interactions is to modify this phase-space distribution and by performing a coincidence experiment it is usually possible to distinguish between the reaction mechanisms.

The proceedings of the Gatlinburg Conference on Correlations of Particles Emitted in Nuclear Reactions (1) contain the best summary at the present time of the work being undertaken on these reactions. In addition to several review articles on

the experimental and theoretical techniques, many interesting experimental results have been reported. It is beyond the scope of this review to give a complete up to date summary of the experimental work on all these reactions. The reactions which will be mentioned are those which have been used to gain information about the nucleon-nucleon interactions, in particular the reactions of interest $\text{He}^3(\text{He}^3, 2p)\text{He}^4$ and $\text{T}(\text{He}^3, np)\text{He}^4$.

One interesting result of the second type of measurement discussed in the general introduction is the discovery of the levels in He^4 at 19.94 and 21.24 MeV by Parker et al at Brookhaven (2). A 31.8 MeV He^3 beam was used to bombard a deuterated polyethylene target and two three particle states were produced, one consisting of $\text{T}+p+p$ and the other He^3+p+n . The final state interactions were observed using two dimensional analysis of the energies of two coincident particle from the reaction, in this case both He^3-p and $\text{T}-p$ coincidences were studied. The energy spectrum of the protons in coincidence with a triton contained two peaks corresponding to the excited states in He^4 . A broad peak was also observed in the $p-\text{He}^3$ coincidence spectra due to the singlet state of the deuteron. No similar peak was found in the $p-\text{T}$ coincidence spectra indicating that at this bombarding energy the $p-p$ singlet state interaction is relatively much weaker.

The nucleon-nucleon interactions have also been observed in several three body reactions (3-6) by measuring the energy spectrum of the third particle, the other two particles being the two nucleons. Tombrello and Bacher (5) have made a systematic survey of several of these reactions. The proton-proton

interaction was studied in the reactions $\text{He}^3(\text{d},\text{t})2\text{p}$, $\text{He}^3(\text{He}^3,\text{He}^4)2\text{p}$ and $\text{He}^3(\text{p},\text{d})2\text{p}$ by observing at forward angles the energy spectra of the triton, alpha particle and deuteron respectively. The proton-neutron interaction was studied in a similar manner in the reaction $\text{He}^3(\text{d},\text{He}^3)\text{pn}$. The incident energies used in these measurements were typically 10-12 MeV. The two nucleon interactions appeared as peaks at the high energy end of the particle spectra and only at very forward angles. The general conclusion drawn from these studies was that although the interactions do play a part in the breakup, the present sequential decay theories could only give qualitative agreement with the observed spectra and no nuclear parameters could be extracted. The most interesting result of the investigation was the absence of any structure in the reaction $\text{He}^4(\text{d},\text{He}^4)\text{pn}$. In this reaction the singlet $T=1$, p-n interaction is inhibited by isospin conservation as the total isospin of the initial state is zero. This is convincing evidence that the interactions being observed are in fact due to the singlet states.

Using a similar experiment, Jarmie and Allen (6) observed the di-neutron interaction in the alpha particle energy spectrum from the reaction $\text{T}(\text{T},\text{He}^4)2\text{n}$. No evidence for a bound di-neutron state was found. Again no quantitative information could be obtained as the peak due to the n-n interaction was superimposed on a continuum from the states of He^5 . These measurements, although interesting since they indicate that the nucleon-nucleon interaction can be conveniently observed as a final state interaction, also point out the need for a more

sophisticated experimental approach.

Coincidence experiments using two dimensional energy analysis have been used by several groups in studying the reaction $\text{He}^3(\text{He}^3, 2\text{p})\text{He}^4$. The possible reaction mechanisms are the following:

Aldridge et al (7) at Rice University have measured the energy spectra of both p-p and p- He^4 coincidences from this reaction at a bombarding energy of 4.9 MeV. The theoretical predictions for these energy spectra were calculated using the generalized density of states formalism of Phillips, Griffy and Biedenharn (8). This formalism will be discussed in some detail in Chapter V. It treats the three particle breakup as a series of two body decays through intermediate states with the energy dependence of the interaction determined from the corresponding elastic scattering phase shifts. The intermediate systems considered in this analysis were the ground state and first excited state of Li^5 and the singlet two proton state. The dominant process in the reaction was found to be the sequential decay through the ground state of Li^5 . No good evidence for the p-p interaction was seen in the spectra.

Bacher and Tombrello (9) have also observed the p-p and p- He^4 coincidences from this reaction at an incident He^3 energy of 9.91 MeV. They found that at this energy the ground state transition was again the dominant process. Proton-alpha coincidences were analysed with a geometry favourable to observing

the p-p interaction at low relative energies. Only a small number of counts were found in this region of the kinematic contour indicating the lack of importance of this final state interaction. The results of Zurmuhle (10) at an incident energy of 15 MeV are in agreement with the two previously mentioned experiments. In this experiment p-p coincidences were observed with a small angular separation of the two detectors but poor statistics prevented any conclusions from being drawn regarding the p-p interaction.

No similar experiments using coincidence techniques have been published for the reaction $T(\text{He}^3, np)\text{He}^4$. The energy spectra of the charged particles from this reaction have been observed by Moak (11) and Kuhn et al (12) for incident energies below 1 MeV. The neutron spectrum have been observed by Barry et al (13) at an incident energy of 3.23 MeV. The possible reaction mechanisms are the following:

		$Q(\text{MeV})$
(1)	$\text{He}^3 + T \rightarrow \text{He}^4 + d$	<u>14.320</u>
(2)	$\rightarrow \text{Li}^5(\text{g.s.}) + n \rightarrow \text{He}^4 + p + n$	10.128
(2a)	$\rightarrow \text{Li}^{5*} + n \rightarrow \text{He}^4 + p + n$	$E_x = 5-10 \text{ MeV}$
(3)	$\rightarrow \text{He}^5(\text{g.s.}) + p \rightarrow \text{He}^4 + n + p$	11.138
(3a)	$\rightarrow \text{He}^{5*} + p \rightarrow \text{He}^4 + n + p$	$E_x = 2-4 \text{ MeV}$
(4)	$\rightarrow \text{He}^4 + p + n$	12.095

The results of Moak and Kuhn indicate that for incident energies less than 1 MeV the reaction channels 1 and 4 are of approximately the same intensity, with channel 3 a factor of 6-8 lower. Barry found that reaction channel 2 has about the same intensity as the sum of 3 and 4 for the incident energy 3.23 MeV. The separation

of the relative contributions is made difficult by the fact that mechanism 4 produces similar single particle spectra as the breakup through the first excited states of Li^5 and He^5 . In all the observed energy spectra there was no evidence for the formation of the singlet state of the deuteron.

C. Present Work

The work which will be described in this thesis was originally undertaken to determine the importance of the p-p interaction in the reaction $\text{He}^3(\text{He}^3, 2\text{p})\text{He}^4$ at bombarding energies around 1 MeV. At the time the work was begun no coincidence measurements on this reaction had been reported. The first series of measurements, described in the author's Masters thesis (14) consisted of observing the angular distribution of coincidence events between the two protons as a function of the angle between the protons. Both detectors were kept in a plane perpendicular to the incident beam direction. The reaction mechanism was determined by comparing the experimental angular distribution with those calculated for the various possible processes obtained from kinematic and phase-space arguments. These calculations predicted what is intuitively obvious, that the distribution of p-p coincidences from the breakup through the states of Li^5 is peaked at an angle of 180° between the protons. Also the only process which produces an increase in the number of coincidence events with decreasing angular separation of the protons is the final state p-p interaction.

The experimental distribution of coincidence events showed that at this incident energy the dominant process is again

the sequential decay through the ground state of Li^5 . A more interesting result was that there was a definite increase in the distribution at small angles between the detectors which could only be attributed to the p-p final state interaction. It was decided to continue the investigation of this reaction using a recently acquired two dimensional kicksorter to analyse the proton-proton events. These experiments were carried out first using the U.B.C. Van de Graaff generator and the results indicated that it would be useful to study the effect of increasing the incident He^3 energy. The measurements at higher energies were made during the summer of 1965 using the Chalk River tandem accelerator. The results of these measurements have been reported (15) and will be discussed more fully in this thesis. It was found that the singlet state p-p interaction was very evident in the coincidence spectra taken at an angular separation of the proton detectors of 6° . The contribution of this interaction relative to the contribution from the ground state of Li^5 decreased as the bombarding energy increased.

The next step in the investigation was to study p- He^4 coincidences from the same reaction with a target-detector geometry arranged to observe the p-p system at low relative energies of the two protons. The same experiment was then carried out using a tritium target in place of the He^3 . The reactions $\text{He}^3(\text{He}^3, 2p)\text{He}^4$ and $\text{T}(\text{He}^3, np)\text{He}^4$ have similar Q-values of 12.859 and 12.095 respectively and therefore the same experimental arrangement allows the singlet di-proton state and the singlet deuteron state to be observed and compared.

This thesis presents the description of the experimental arrangement and the technique used in performing the measurements along with a discussion of the results. Two approximate theories of final state interactions, the Watson theory (16) and the Phillips, Griffy and Biedenharn generalized density of states formalism (8) are used to fit the observed spectra. Statements are made about the validity of these two formalisms and the suitability of this experimental technique in determining nuclear parameters.

CHAPTER II

KINEMATICS AND EXPERIMENTAL TECHNIQUE

A. Kinematics of Three Particle Breakup Reactions

Before a systematic investigation of the final state interactions involved in a three particle breakup reaction can be carried out, it is necessary to have a complete understanding of the reaction kinematics. The reason for choosing a coincidence type of measurement rather than observing the single particle energy spectra becomes evident once the kinematics of the breakup are understood. In addition a number of kinematic calculations are necessary before one can choose the appropriate target-detector geometry to scan the desired range of internal or excitation energies of the possible two particle systems.

Several coordinate systems may be used to describe a three body decay. These include the laboratory system (lab) in which the target nucleus is at rest, the system centre of mass (scm) and the various recoil centres of mass (rcm(i)). The latter are useful in the theoretical analysis of the breakup when the sequential two body decay mode is important. The general non-relativistic kinematic formula for a three particle breakup in these coordinate systems have been derived in several recent publications (17,18). The important expressions are given for reference in Appendix A.

Some of the expressions useful in the design of the experiment will be derived in this chapter. The symbols for the various breakup parameters which will be used are those given by Bronson (17) and are summarized in Table 1. The relationships

TABLE 1

Symbols used in describing a three particle breakup in the three relevant coordinate systems. (reference Figure 1 and 2).

Coordinate system		Definition
lab.	scm.	
T_i	t_i	kinetic energy of particle i.
V_i	v_i	velocity of particle i.
P_i	p_i	momentum of particle i.
Θ_i	θ_i	polar angle of velocity of particle i with respect to the beam direction.
Φ_i	φ_i	azimuthal angle of velocity of particle i with respect to the beam direction.
Δ_{ij}	δ_{ij}	angle between the two velocities i and j.
rcm. system for particle i emitted first		
$p_j^{(i)}$		momentum of particle j in system (i).
$\Theta_j^{(i)}$		polar angle of particle j with respect to the beam direction.
$\varphi_j^{(i)}$		azimuthal angle of particle j with respect to the beam direction.
$\eta_j^{(i)}$		polar angle of particle j with respect to the recoil axis.
V_{scm}		velocity of scm. system in lab. system.
$v_{rcm}^{(i)}$		velocity of rcm(i) system in scm. system.
Q		Q-value for overall reaction.
E_{jk}		internal or excitation energy of cluster (j+k).
m_i		mass of particle i.
M		total mass of system

Useful relations:

$$\cos \Delta_{ij} = \cos \Theta_i \cos \Theta_j + \sin \Theta_i \sin \Theta_j \cos (\Phi_i - \Phi_j)$$

$$\cos \delta_{ij} = \cos \theta_i \cos \theta_j + \sin \theta_i \sin \theta_j \cos (\varphi_i - \varphi_j)$$

between the parameters in the three coordinate systems are shown in Figures 1 and 2. The subscripts 0, 1, 2, 3 are used to designate the incident particle, the two detected particles and the undetected particle respectively. Since it is possible that different pairs of particles may be detected with the two detectors, the subscripts 1, 2 will refer specifically to the detector positions while letter subscripts b, c, d will be used to differentiate between the particles.

A three particle final state is completely described by nine quantities, for instance the nine components of the three particle momenta or the energy and the two polar coordinate angles of each of the three particles. These are reduced to five independent quantities by imposing energy and momentum conservation.

$$\vec{P}_0 = \vec{P}_1 + \vec{P}_2 + \vec{P}_3 \quad 2-1$$

$$T_0 + Q = T_1 + T_2 + T_3 \quad 2-2$$

with

$$P_i^2 = 2m_i T_i$$

Solving equations 2-1 and 2-2 for P_3 and regrouping the terms gives the quadratic equation

$$AP_2^2 - 2BP_2 + C = 0 \quad 2-3$$

where

$$A = (m_2 + m_3) / m_2$$

$$B = P_0 \cos \theta_2 - P_1 \cos \Delta_{12}$$

$$C = (1 + m_3/m_1) P_1^2 + (1 - m_3/m_0) P_0^2 - 2P_0 P_1 \cos \theta_1 - 2m_3 Q$$

Figure 1. Relationship between velocities in lab. and scm. coordinate systems.

Figure 2. Relationship between velocities in scm. and rcm(i) coordinate systems.

The solutions of the quadratic equation are

$$P_2^{\pm} = (B \pm \sqrt{B^2 - AC}) / A \quad 2-4$$

with the range of values of P_1 determined by the requirement that

$$B^2 - AC \geq 0$$

Coincidence events associated with a particular breakup (for instance particle b $\rightarrow$ detector 1, particle c $\rightarrow$ detector 2) must be on a curved locus described by equation 2-4 in the P_2 vs. P_1 plane. A similar locus can be calculated for the related T_2 vs. T_1 plane. The kinetic energy T_3 of the third particle and the laboratory polar angles Θ_3, Φ_3 may be calculated from the following expressions.

$$T_3 = T_0 + Q - T_1 - T_2 \quad 2-5$$

$$\cos \Theta_3 = (P_0 - P_1 \cos \Theta_1 - P_2 \cos \Theta_2) / P_3 \quad 2-6$$

$$\cos \Phi_3 = -(P_1 \sin \Theta_1 \cos \Phi_1 + P_2 \sin \Theta_2 \cos \Phi_2) / P_3 \sin \Theta_3 \quad 2-7$$

Since observing only one particle in the breakup determines three of the five quantities necessary to describe the breakup, the energy spectrum of one particle will be an average over the two unobserved quantities. An experiment of this type is meaningful only if the breakup occurs sequentially through a reasonably well defined intermediate state and if the spectrum of the first emitted particle is observed. This reduces the breakup to the usual two body decay. If no particle identification is used, then the averaging process tends to cover up much of the

structure which might be expected to appear in the spectrum if an intermediate state or strong final state interaction is present. Several improper conclusions have been made in the past of the dominant decay mode in this type of reaction due to a failure to observe the above requirements. This indeterminacy can be removed by using two detectors to observe two of the final state particles in coincidence.

Since the angular coordinates of the two detectors determine the polar and azimuthal angles $\Theta_1, \Phi_1, \Theta_2, \Phi_2$, it is necessary to determine only one more quantity to specify the breakup completely. This quantity is usually the energy of one of the detected particles. However experimentally it is desirable to determine both T_1 and T_2 for the following reasons:

- a) The background from accidental coincidences and contaminant two body reactions is greatly reduced.
- b) From equation 2-4 it is evident that for certain detector angles P_2 , and therefore T_2 , is a double valued function of P_1 . These two solutions can be separated by observing the energies of both particles.
- c) If b,c,d are distinct particles of different mass then there exist six distinct loci T_2 vs. T_1 corresponding to the six permutations of three particles with two detectors. The problem of associating each event with the appropriate locus is somewhat simplified by using two dimensional energy analysis. If the loci overlap considerably then some form of particle identification is necessary as well. Another solution is to perform a triple coincidence

measurement with three detectors, together with two dimensional energy analysis as has been done by Reimann in the reaction $\text{Li}^6(\text{He}^3, p)2\text{He}^4$ (19).

The experimental work described in this thesis uses two detectors to perform the two dimensional energy analysis. The parameters which are determined are $\Theta_1, \Phi_1, \Theta_2, \Phi_2, T_1$, and T_2 .

B. Experimental Technique.

Typical loci of coincidence events or "kinematic contours" in the T_2 vs. T_1 plane are shown in Figure 3 for p-p and p- He^4 coincidences from the reaction $\text{He}^3(\text{He}^3, 2p)\text{He}^4$. These contours have been calculated for an incident beam energy $T_0=1.50$ MeV and for the case with the two detectors coplanar with the beam. To simplify the situation detector 1 is assumed to observe only protons while detector 2 can observe both protons and alpha particles. Detector 2 is fixed at $\Theta_2 = 90^\circ$ and detector 1 can be rotated in the plane of the beam with the angular separation of the detectors $\Delta_{12} = \Theta_1 + \Theta_2$. The graph shown has been computer generated using the U.B.C. IBM 7040 computer and a Calcomp plotter. A general program for calculating kinematic contours is listed in Appendix B. The number adjacent to each contour refers to the value of Δ_{12} .

The distribution of coincidence events along a kinematic contour is determined by the interactions which take place between the particles during breakup. The usual method of comparing the experimental coincidence yield with theoretical predictions is to use the triple correlation distributions (20)

Figure 3. Kinematic contour for p-p and p-He⁴ coincidences from the reaction $\text{He}^3(\text{He}^3, 2p)\text{He}^4$. The number adjacent to each curve is the corresponding value of Δ_{12} .

given by

$$\sigma(T_1) = \frac{d^3\sigma}{dT_1 d\Omega_1 d\Omega_2} \quad 2-8$$

or

$$\sigma(T_2) = \frac{d^3\sigma}{dT_2 d\Omega_1 d\Omega_2} \quad 2-9$$

These distributions are obtained by summing the coincidence events along a particular contour for constant T_1 in the case of $\sigma(T_1)$ and T_2 in the case of $\sigma(T_2)$. These sums are usually displayed in the form of a histogram. A typical example of this is shown later in Figure 15.

As mentioned in the introduction, two reaction mechanisms are possible for a three body decay: direct breakup into three particles and sequential breakup through an intermediate state. In the first case, if it is assumed that the breakup is purely statistical, that is the decay probability depends only on the available phase-space, then the triple correlation has the form (see Appendix A):

$$\frac{d^3\sigma}{dT_1 d\Omega_1 d\Omega_2} = \frac{K P_1 P_2^2}{AP_2 + P_1 \cos \Delta_{12} - P_0 \cos \Theta_2} \quad 2-10$$

where K is a constant depending on the particle masses. This yields a broad structureless distribution which is characteristic of statistical breakups.

If the sequential decay process takes place, then there will be an increase in the coincidence yield at points in the T_2 vs. T_1 plot where the internal or excitation energy of one of the two particle clusters, either E_{12} , E_{13} , or E_{23} , corresponds

to a state of that particular system. The width of this state will be reflected in the width of the peak in the coincidence distribution. A well defined state will appear as a point in the T_2 vs. T_1 plot while a broad state will produce coincidence events over a segment of the contour. Since the excitation energies which are accessible along a given contour depend on the detector angles, it is worthwhile working out the relations between the excitation energy and the other breakup parameters before designing the target chamber for any particular reaction.

Two cases must be considered in this calculation:

- 1) the first emitted particle in the breakup is detected along with one of the particles from the breakup of the intermediate system.
  - 2) both detected particles come from the intermediate system.
- In the first case ordinary two body kinematic relations can be used. If we assume the intermediate system is (2,3) then the momentum and energy relations are given by

$$\vec{P}_0 = \vec{P}_1 + \vec{P}_{23} \quad 2-11$$

$$T_0 + Q - E_{23} = T_1 + T_{23} \quad 2-12$$

where $\vec{P}_{23}$ and T_{23} are the momentum and energy of the virtual system (2,3). Solving both equations 2-11 and 2-12 for P_{23} and rearranging terms gives

$$E_{23}(T_1) = Q + \frac{1}{m_2+m_3} \left[2\sqrt{m_0 m_1 T_0 T_1} \cos \Theta_1 - T_0(m_0 - m_2 - m_3) - T_1 M \right] \quad 2-13$$

A similar result may be obtained for $E_{13}(T_2)$ by interchanging the

subscripts 1 and 2.

In the second case the intermediate system is (1,2). The momentum and energy relations may be written

$$\vec{P}_{12} = \vec{P}_1 + \vec{P}_2 \quad 2-14$$

$$T_{12} + E_{12} = T_1 + T_2 \quad 2-15$$

which yields on eliminating $\vec{P}_{12}$

$$E_{12} = \frac{1}{m_1 + m_2} \left[m_2 T_1 + m_1 T_2 - 2 \sqrt{m_1 m_2 T_1 T_2} \cos \Delta_{12} \right] \quad 2-16$$

Curves of fixed excitation energy in one of the two particle systems may be drawn on the T_2 vs. T_1 plot. If it is desired to study a particular system at a given excitation energy then it is necessary to choose a detector arrangement which yields a contour intersecting the excitation curve. The contour should be chosen so that no other intersection points of the excitation curves from the other possible systems are near to the one of interest. Figure 4 shows the p-He⁴ contours from Figure 3 along with the loci of constant excitation in the three intermediate systems. The curves marked $E_{23} = 1.97$ MeV and $E_{12} = 1.97$ MeV correspond to the ground state of Li⁵ and a breakup proceeding through this state would produce events along these curves. The cross-hatched area represents the region where events from the breakup of a two proton system with relative energy less than 500 keV would be expected to appear.

Figure 4. Kinematic contours for p- He^4 coincidences from the reaction $\text{He}^3(\text{He}^3, 2p)\text{He}^4$. Also shown are some of the corresponding excitation energies in the two particle systems.

CHAPTER III

EXPERIMENTAL ARRANGEMENT

A. Introduction

As helium-3 and tritium are both gases at room temperature, it is an obvious choice to use a gas target in this particular experiment. Gas targets have several advantages in angular distribution measurements or measurements requiring more than one detector. Target uniformity is inherent except for small beam heating effects and there are no problems with target backings. Contaminant reactions can be held to a minimum by using collimators to determine the active volume of the target gas. Gas targets may be divided into two types: those in which the beam enters from the high vacuum region through a thin window and those in which the beam enters through a differentially pumped capillary. The first type was used in this experiment as it was desirable to use as little target gas as possible, particularly in the case of tritium. Nickel foil was chosen as the material for the entrance window because of its high mechanical strength and the commercial availability of high quality vacuum-tight foils of the required thicknesses.

The charged particles from the reaction are monitored with solid state detectors. The requirements on these detectors are linear response for protons over a range of energies from 500 keV to 14 MeV and for alpha particles from 500 keV to 5 MeV along with an intrinsic resolution of less than 35 keV. Surface barrier detectors using high resistivity (12,000 ohm-cm) n-type silicon were found to meet these requirements. Two methods of

detecting the reaction products are possible when gas targets are employed. The detectors may be immersed in the target gas using a series of collimators and defining slits to determine the active target volume or they may be placed in the vacuum outside the gas cell. In the latter case the gas cell must have exit windows of very thin foils through which the charged particles pass. The second method is preferable if large target gas pressures are used as the energy loss in traversing the target gas may then be of more concern than the energy loss in the exit foil. In addition very small quantities of target gas may be used with this arrangement. Both arrangements were used in course of these experimental measurements.

The prime consideration in the chamber design other than the requirement that gas targets and solid state detectors be used is the angular variation of the detectors. The polar and/or azimuthal angles of the two detectors are required to be variable in such a way that the desired range of excitation energies of the intermediate systems can be obtained. In general there is more than one detector configuration which can be used to yield the same information about the intermediate states present, and there is no *a priori* reason for choosing one configuration over another. The decision can therefore be made on the basis of simplicity in chamber construction.

Two reaction chambers were employed in the experiment. The first target-detector configuration called Geometry #1 was used in determining the relative contributions of the states of Li^5 and the two proton system in the reaction $\text{He}^3(\text{He}^3, 2p)\text{He}^4$ by

studying proton-proton coincidences. In this geometry, shown schematically in Figure 5a, the polar angles of both detectors are kept at 90° with the angular separation of the detectors variable in the azimuthal direction from 180° to 6° . Figure 6 shows the kinematic contours for p-p, p-He⁴ and He⁴-p coincidences from this reaction at an incident energy of 1.15 MeV and for angular separations $\Delta_{12} = 6^\circ$ and $\Delta_{12} = 180^\circ$. The p-He⁴ and He⁴-p coincidences may be removed by placing a foil in front of each detector to degrade the energy of the alpha particles. This results in only a slight change to the proton-proton contours. The regions of interest in the T_2 vs. T_1 plot of p-p coincidences are shown in Figure 7. The crosshatched area indicates the region which will be populated by p-p coincidence events corresponding to a relative energy in the two proton system of less than 500 keV. Coincidence events lying along the two lines marked E_{13} and E_{23} represent a relative energy in the p-He⁴ system of 1.97 MeV corresponding to the ground state of Li⁵. It is evident that this geometry gives good separation between the contributions from both intermediate systems.

The second target-detector configuration called Geometry #2 was used in the investigation of the p-p and p-n final state interactions in the reactions $\text{He}^3(\text{He}^3, 2p)\text{He}^4$ and $\text{T}(\text{He}^3, np)\text{He}^4$. As it was necessary to use p-He⁴ coincidences to compare the interactions, Geometry #1 did not allow the desired range of excitation energies to be studied. In Geometry #2 as shown in Figure 5b both detectors are kept in the plane of the beam with one detector fixed at an angle of 90° with respect to the beam

(a) GEOMETRY #1

(b) GEOMETRY #2

Figure 5. Schematic diagram of Geometry #1 and Geometry #2.

Figure 6. Kinematic contours for $p-p$, $p-\text{He}^4$ and He^4-p coincidences from the reaction $\text{He}^3(\text{He}^3, 2\text{p})\text{He}^4$. The number adjacent to each curve is the corresponding value of Δ_{12} .

Figure 7. Kinematic contours for p-p coincidences from the reaction $\text{He}^3(\text{He}^3, 2\text{p})\text{He}^4$. Also shown are some of the corresponding excitation energies in the two particle systems.

direction. The other detector is variable in the polar direction for angular separations of the detectors from 180° to 135° . The kinematic contours and excitation curves for this geometry have been described previously in Figures 3 and 4. In studying both reactions it is necessary to discriminate against He^4 -p coincidences in which the alpha particle enters detector 1. Again the discrimination is made with the use of a thin foil.

The details of the two target chambers and a description of the tritium handling system is given in the first part of this chapter. The rest of the chapter contains a discussion of the electronics used in the two dimensional energy analysis and the use of an on-line computer to simplify the data handling and analysis.

B. Geometry #1 Target Chamber

The target chamber is shown in Figure 8. The external beam, defined by a series of collimators (A), enters the chamber through a thin nickel foil mounted on a removable brass tube (B). The detector holders (C) and (D) are mounted on radial arms, one of which is fixed and the other may be rotated in the plane perpendicular to the beam direction. The angular separation of the detectors is read on the scale (E) on the outside of the chamber. The detector holders may be positioned at any radial distance from the beam axis, within the confines of the chamber, and they may also be rotated about an axis parallel to the beam direction. This rotation serves two purposes: to increase the effective depletion thickness of the detectors and to allow the detectors to be positioned closer together for measurements

Figure 3. Geometry #1 target-detector arrangement (side-view).

Scale $\frac{1}{2} = 1$.

at small angular separations. The measurements at an angle of 6° were made using two rectangular detectors mounted together in a lucite block. The detectors were usually placed at a distance of 5.0 to 5.5 cm. from the beam axis. The active volume of the gas target is determined by a cylindrical slit (F) of width 5 mm. at a distance of 12.7 mm. from the beam axis and by adjustable slits in front of the detectors. The alignment of the entrance apertures, the defining slits and the angular scale was done optically with the angular readings accurate to better than $\pm 1^\circ$.

The incident beam is stopped in a brass cup (G) insulated from the rest of the chamber to allow the beam current to be monitored. This measurement was used only as a means of optimizing the beam focus as the target gas surrounding the beam stop prevented any accurate beam current determination. The nickel foils used in the entrance window are typically 0.025 to 0.050 mils thick. These foils are Grade C (wholly light-tight) $\frac{1}{2}$ " x $\frac{1}{2}$ " square, obtainable from Chromium Corporation of America, Waterbury, Connecticut. Originally the foils were soft-soldered to their brass holders using a high tenacity flux but this technique was found to cause considerable stress to the foils due to the heating. A simpler method of attaching the foils, using an epoxy resin HYSOL type 4307 with type 3404 hardener (obtainable from HYSOL Corporation, Olean, New York) gave considerably better success. The loss of a good heat contact between the foil and its holder did not seem to limit the maximum beam current the foil could withstand indicating that

a good portion of the foil heating is removed by conduction through the target gas.

The target gas pressure is continuously monitored with a Wallace and Tiernan Absolute Pressure Indicator Type FA-160 (0-410 mm. Hg). A target pressure of 100-150 mm. Hg was generally used. The He^3 target gas was obtained from Monsanto Research Corporation and had a specified purity of 99.8% He^3 with less than $3.6 \times 10^{-11}\%$ tritium.

C. Geometry #2 Target Chamber

Several safety features were incorporated in the design of this chamber to allow it to be used with tritium gas. The volume of the gas cell and the filling manifold was kept low so that as little target gas as possible would be used. The gas cell was designed so that it could be easily installed, as it was intended to discard the contaminated gas cell in the event of breakage of the entrance or exit foils. Two views of the target assembly are shown in Figures 9 and 10. The gas cell (A) is mounted on one of the side flanges using Teflon washers (H) and a Neoprene O-ring seal to keep it insulated from the rest of the chamber. The target gas enters the cell through a 1/16" diameter hole drilled lengthwise through the stem of the gas cell and aligned with a similar hole in the mounting flange. A spring clip attached to a Kovar seal on the rear flange (C) is attached to the stem of the cell after installation to allow the beam current to be monitored.

Foils are attached to the gas cell on three sides.

Figure 9. Geometry #2 target-detector arrangement (side-view).

Scale $\frac{1}{2} = 1$.

Figure 10. Geometry #2 target-detector arrangement (rear-view).
Full scale.

The foils for the entrance window and alpha particle exit window are typically 0.025 mils thick and the foil covering the proton exit window is 0.200 mils thick. The proton window has a rectangular cross section so that the protons can be observed at polar angles from 90° to 45° with respect to the beam direction. These foils were attached using the epoxy resin previously mentioned and the gas cell was vacuum tested before being installed using a differential pressure of 200 mm. Hg of hydrogen. A success ratio of approximately 50% was obtained in mounting the foils.

The proton detector (B) is mounted on a brass arm (I) which also contains a pair of collimators to define the solid angle. This arm can be rotated in the plane of the beam by means of a steel rod (F) vacuum coupled through the remaining side flange. A pointer (G), attached to the end of this rod rotates over an angular scale (not shown) mounted on the outside of the chamber to allow the angular separation of the detectors to be determined. The alpha particle detector (D) is fixed at an angle of 90° with respect to the beam direction. The positioning of the two detectors is accurate to $\pm \frac{1}{2}^\circ$.

The active volume of the gas target is determined by a 4.75 mm. diameter aperture (J) at a distance of 16 mm. from the axis of the target and by a rectangular slit (K) in front of each detector. An antiscattering collimator (L) is also provided to prevent the high energy protons from entering the detectors after scattering from the chamber walls. The incident beam enters the chamber through a series of stainless steel collimators (E).

The entire assembly was aligned optically using a light beam from a helium-neon laser.

D. Tritium Handling System

Figure 11 is a diagram of the pumping and gas handling system used in conjunction with the tritium and helium gas targets. The system can be isolated from the vacuum system of the beam handling equipment by using a C.V.C. pneumatically operated gate valve. This valve can be operated manually or actuated by a pressure rise in the target chamber. The valve has its own air supply and is operated in the normally close mode so that an electrical failure will leave the valve closed. The signal to automatically actuate the valve is obtained from an ionization gauge with type UBC-NP-12 control unit situated in front of the target collimators. Normally the pressure in the vacuum system is 3×10^{-6} mm. Hg. The adjustable trip-out of the gauge is set to operate when the pressure rises to 10^{-5} mm. Hg. In order to prevent any gas from the target cell from entering the beam line before the valve is completely closed, a series of baffles is placed in the line between the ionization gauge and the gate valve.

The vacuum system consists of an Edwards High Vacuum 2M4A mercury diffusion pump which pumps into a reservoir of one litre capacity. This booster pump was chosen because it can operate into a backing pressure of 30-35 mm. Hg. The pumping reservoir may be evacuated using a Welsh type 1405 Duo-seal mechanical pump when non-contaminating gases are being used. If

Figure 11. Diagram of pumping and gas handling system.

tritium is present the reservoir is opened to a charcoal trap charged with 225 gms of activated charcoal and immersed in a liquid nitrogen bath. The adsorption relation for hydrogen in charcoal at liquid nitrogen temperatures is given by (21)

$$V_o = 1.6 \times 10^{-2} P_u$$

where V_o is the volume of hydrogen in cc. at S.T.P. adsorbed per gram of charcoal and P_u is the equilibrium pressure in microns.

The gas handling manifold consists of four HOKE diaphragm seal valves of all metal construction silver soldered to a brass block. The tritium container ORNL type HoH-50-3 containing 10cc. of tritium is connected to the manifold through valve A. Valves B and C allow both sides of the gas cell to be evacuated simultaneously to maintain a minimum pressure differential across the foils. Valve D is used to attach the He^3 container, as well as for leak testing using a helium leak detector. The gas pressure in the target cell is continuously monitored with a Bourdon type pressure gauge (0-30" Hg vacuum). The volume of the target system and pressure gauge is 20 cc. with the gauge accounting for most of the volume. A pressure of 50 mm. Hg, equivalent to about 4 curies of tritium, was used during the experimental measurements.

The entire system is enclosed in an air-tight box covered with 1/8" thick lucite and exhausted to the outside open air using a small aerofoil fan of approximately 200 c.f.m. capacity. Ports are cut in the lucite to allow the necessary valves to be manipulated and the gas cell to be replaced with a minimum danger of personal contamination. Tests were carried out on the tritium

handling system using ordinary hydrogen before the tritium container was connected. The tritium level inside the box is continuously measured with a tritium-in-air monitor capable of detecting levels down to 5 micro-curies per cubic meter.

No problems were encountered with the tritium system during the experimental runs. Only one gas cell was required with the tritium as it remained intact for the entire duration of the measurements.

E. Charged Particle Detectors

With the exception of the pair of rectangular detectors employed in the small angle measurements at Chalk River the solid state detectors are type PH-8-35-10 obtained from Nuclear Diodes Inc. These detectors are the surface barrier type consisting of a p-n junction formed at the surface of a slice of high resistivity (10,000 - 12,000 ohm-cm) n- type silicon. Electrical contact is made to the p - type layer through a thin gold film evaporated on the front surface, and to the n - type wafer through a non-rectifying contact on the back surface. A reverse bias of 350-400 volts gives the required depletion thickness of 1000 microns. Larger effective depletion thicknesses are obtained by tilting the detectors at an angle to the direction of the incoming particles. The active area of these detectors is 10 mm. diameter and the intrinsic resolution for 5 MeV alpha particles is about 35 keV. The pair of rectangular detectors were fabricated by the Chalk River Counter Development Group to be used in polarization measurements. These detectors have an active area of 14 mm. by 14 mm.

but this was reduced to 6 mm. by 4 mm. using an aluminum slit arrangement for the measurements reported here. These detectors were mounted together with a separation of 1 mm. between the active areas to allow the measurements to be made at a small angular separation.

F. Electronics

The use of a charge sensitive preamplifier in conjunction with a solid state detector results in a voltage pulse from the amplifier which is linearly proportional to the particle energy, assuming the depletion depth of the detector exceeds the maximum range of the charged particles being detected. The electronic arrangement used in the two dimensional energy analysis of the pulses from two such detector-preamplifier systems is described in this section. Briefly, the function of this arrangement is to amplify the pulses from both detectors, corresponding to particle energies T_1 in detector 1 and T_2 in detector 2, and to store the event in a T_1 vs. T_2 array if certain requirements are met. These requirements for the measurements reported here are that the two pulses be coincident in time within 50 nsec. and that the corresponding energies T_1 and T_2 be within a given range of values. The coincidence requirement ensures that the two particles being analysed come from the same breakup and the energy requirement eliminates random coincidences due to the scattered incident beam.

Two coincidence systems were employed in the course of the experiment as the measurements were carried out using both the University of British Columbia 3 MeV Van de Graaff accelerator

and the Chalk River 12 MeV tandem accelerator. A block diagram of the fast-slow coincidence system used at Chalk River is shown in Figure 12. This arrangement has been described at length by Tomlinson and Brown (22). It incorporates a 100 channel kicksorter to provide a time display of the coincidence events. The two dimensional data acquisition is carried out using a PDP-1 computer. A description of this on-line system is given in the following section.

The coincidence system used at U.B.C. is shown in Figure 13. This arrangement is made up of standard commercial units (see Table 2) with the exception of the charge sensitive preamplifiers. The preamplifiers, described by Whalen (23), produce a linear output pulse with a fast risetime of 10-15 nsec. and a slow decay time of 100 usec. suitable for use in delay line clipped main amplifiers. The outputs of the double delay line clipped amplifiers are fed into single channel analysers which produce an output pulse at the zero-crossover point only if the input pulse falls within the required window. These outputs are used as the start and stop pulses for a time to pulse height converter. The alpha particle pulse is delayed by 50 nsec. to centre the time spectrum in the display. This display is obtained using a Nuclear Data ND 120 kicksorter operating in the 128 channel mode. A further single channel analyser is used to set a window on the time spectrum to minimize the random coincidence events and to remove those events with no stop pulse. A typical time spectrum is shown in Figure 14 for p-p and p-He⁴ coincidences from the reaction $\text{He}^3(\text{He}^3, 2\text{p})\text{He}^4$. The window

TABLE 2

Commercial Electronic Units used in the Experimental Arrangement (see Figure 13).

- 1) Cosmic Model 901 A Linear Amplifier
(Cosmic Radiation Labs. Inc., Bellport, N.Y.).
- 2) Cosmic Model 901 SCA Single Channel Analyser.
- 3) LRS Model 108H Analog Storage Time to Height Converter
(Lecroy Research Systems Corp., Irvington, N.Y.).
- 4) CI Timing Single Channel Analyser
(Canberra Industries Inc., Middletown, Conn.).
- 5) ORTEC Model 430 Scalers
(Oak Ridge Technical Enterprises, Oak Ridge, Tenn.).
- 6) Nuclear Data ND 160 Dual Parameter Analyser
(Nuclear Data Inc., Palatine, Ill.).
- 7) Nuclear Data ND 120 512 Channel Analyser
- 8) Datapulse 106A Pulse Generator
(Datapulse Inc., Inglewood, Calif.).

Figure 13. Block diagram of electronics used in data acquisition at U.B.C.

Figure 14. Time distribution spectrum of coincidences from the reaction $\text{He}^3(\text{He}^3, 2p)\text{He}^4$.

setting is also shown. The two peaks arise from the different response of the solid state detectors to protons and alpha particles. The resulting pulse shapes produce the 10 nsec. time difference between the start and stop pulses.

The two linear pulses from the main amplifiers are fed into the F and M sides of a Nuclear Data ND 160 Dual Parameter Analyser operating in the 64x64 mode and gated with the output of the timing single channel analyser. A coincidence resolving time of 50 nsec. was used as this resulted in a negligible random rate. The memory of the ND 160 kicksorter is transferred into a PDP-8 computer after each run for preliminary analysis.

The timing of the coincidence system and the linearity of the amplifiers were tested by feeding pulses from two Datapulse generators into the test inputs of the preamplifiers. By varying the delay between the pulses the coincidence resolving time could be determined and the timing single channel analyser set to give the desired window. The outputs of the single channel analyser from the alpha particle detector (detector 2) and the timing single channel analyser are connected to scalers to monitor the singles and coincidence rates.

G. On-line Computer Techniques

The use and usefulness of an on-line computer and "data-simulation" techniques in the analysis of three particle breakup reactions has been described by Donovan (24). A similar analysis system was set up using the on-line computer facility

at the Chalk River tandem accelerator. This facility is based on a PDP-1 (Digital Equipment Corporation) computer with four memory modules of 4096, 18 bit words. Along with the standard input-output devices such as tape readers, typewriter and card punch, there is a 16" Cathode ray tube and light pen for visual display. Connected to the computer via a sequence break system are eight 400 channel A.D.C. units for the purpose of pulse height analysis. The sequence break, usually triggered by a coincidence pulse indicating a desired event has taken place, interrupts the computer processor from the operation it is performing at that particular time. In data analysis the program that is interrupted is usually a general purpose spectrum handling routine where it might be displaying some spectrum of interest. The interrupt causes a transfer of control to a data storage routine which deals with the event by reading the appropriate buffers of the A.D.C. units and incrementing the proper memory locations.

During the measurements on the reaction $\text{He}^3(\text{He}^3, 2p)\text{He}^4$ the computer was programmed to store the events from p-p coincidences in a two dimensional 60x60 channel array and to display these events as they accumulated either on a contour T_2 vs. T_1 plot or on an isometric display. A general program for generating the kinematic contours was also written and the appropriate contour was displayed simultaneously with the experimental events. As mentioned previously in Chapter II, the final state interactions appear as regions of high density of events along the kinematic contour. To give an immediate idea of the interactions present, a routine was written to sum the coincidence events along either

energy axis and within a specified number of channels on either side of the predicted kinematic contour. The number of channels summed over could be varied to check the effect of including the few counts lying well off the contour. The two summed spectra and the spectrum obtained by adding these two spectra together were displayed on the 16" CRT as well. The spectra may be added as they are identical due to the symmetry of the experiment with Geometry #1. It was found that the number of random and contaminant events was negligible so that no provision was made for subtracting a background. This facility eliminated the necessity of performing time consuming off-line analysis and removed the guesswork of determining when sufficient statistics had been accumulated.

The pulse height analysis system at the University of British Columbia consists of several conventional kicksorters including a ND 160 dual parameter analyser and a PDP-8 computer of 4096 word memory with a magnetic tape system for data and program storage. The coincidence events, both p-p and p-He⁴ in the case of the reaction $\text{He}^3(\text{He}^3, 2\text{p})\text{He}^4$ and Geometry #2, were accumulated with the ND 160 kicksorter operating in the 64x64 mode. At the time the experiment was performed the PDP-8 computer was not directly interfaced to any analog to digital converters for the purpose of pulse height analysis. Provision was made, however, for the rapid transfer of information from the memory of the ND 160 to the computer memory. A short program was written to transfer a 64x60 channel array from the kicksorter to the computer. The remaining 256 locations in the computer were

used to store the transfer program and a program to type out the information in memory either as three 64×20 arrays of two digit numbers or six 64×10 arrays of four digit numbers. Zero suppression was used in the typeout so that the contours could be easily seen in the arrays of numbers. This form of presentation of the data proved to be quite useful in obtaining a preliminary check on the experiment.

CHAPTER IV

EXPERIMENTAL PROCEDURE AND RESULTS

In the previous chapters the two reaction chambers and data accumulation systems have been described along with the kinematic calculations for both the Geometry #1 and Geometry #2 target-detector configurations. In this chapter an outline of the experimental procedure with both arrangements and the results of the measurements are given. The measurements which were made using the Geometry #1 arrangement are discussed first.

A. Geometry #1 Measurements.

These measurements were carried out at three incident energies using a well collimated He^{3+} beam from the Chalk River tandem accelerator. The beam energies which were chosen for studying the reaction $\text{He}^3(\text{He}^3, 2p)\text{He}^4$ were 1.90 MeV, 3.00 MeV and 5.40 MeV. These energies correspond to incident energies inside the gas target, after correcting for the energy loss in the entrance foil, of 1.15 MeV, 2.40 MeV and 5.0 MeV respectively. The nickel foil used in the entrance window was nominally 0.050 mils thick. The thicknesses of all foils used in the course of the experiment were determined from the energy loss of the 5.477 MeV alpha particles from an Am^{241} source. A computer program described in Appendix C is used to calculate the energy loss of the different particles in the foils. Several curves are also included in this Appendix showing the energy loss as a function of particle energy for p, d, He^3 and He^4 .

Figure 6 shows that the proton-alpha and proton-proton kinematic contours for Geometry #1 do not overlap for a

particular angular separation of the two detectors, although for angles Δ_{12} around 180° the contours lie sufficiently close together that some alpha particle discrimination is required. This was obtained by placing a 3.0 mg/cm^2 aluminium foil in front of each detector. From Figure 7 it is evident that, to observe the p-p interaction at low relative energies with this geometry, it is necessary to use as small an angular separation of the two detectors as possible. The minimum angular separation, $\Delta_{12} = 6^\circ$, which was used in the measurements, was determined by the detectors available and the coincidence counting rate. To scan the entire range of angles from $\Delta_{12} = 180^\circ$ to $\Delta_{12} = 6^\circ$ three different detector configurations were employed. These are listed in Table 3 along with some of the relevant information about the solid angles involved. The solid angles subtended by both detectors were approximately equal.

The single particle yield in each detector is proportional to the geometrical factor $G = L\Omega$ where L is the target thickness and Ω is the effective solid angle subtended by the detector. For a gas target the G factor is given approximately by

$$G = sA/Rh \quad 4-1$$

where s is the width of the first defining slit,

A is the area of the second defining slit,

R is the distance of the second defining slit from the beam axis,

h is the distance between the two slits.

The ratio of the coincidence rate to the singles rate therefore

TABLE 3

Detector configurations used with Geometry #1 measurements.

Detector conf.	Angular range covered	Slit shape in front of detectors	Angular width of detectors		$\frac{G_1 G_2}{G_1 + G_2}$ cm. $\times 10^3$	Foil in front of detectors
			$\Delta\Theta$	$\Delta\Phi$		
#1	180°-22°	Circular	11°	11°	9.50	3.0 mg/cm ² aluminum
#2	22°-13°	Elliptical	11°	6°	5.01	none
#3	6°	Rectangular	7°	4°	3.60	none

depends on the factor $G_1 G_2 / (G_1 + G_2)$ where G_1 and G_2 are the geometrical factors for the two detectors. This factor was determined for detector configuration #1 by using the expression given above and the appropriate dimensions. The values for configurations #2 and #3 were determined relative to configuration #1 by placing an alpha source (ThC') at the centre of the reaction chamber to simulate the active volume of the target gas, and comparing the counting rates for the different configurations. These results are listed in the second last column of Table 3. the same quantities evaluated for detector configurations #2 and #3 using expression 4-1 and the slit dimensions agree within 10%. The normalization between runs at different angular separations and with different detector configurations is obtained by monitoring the singles rate in both detectors and using the corresponding solid angle ratio. The correction to the singles rate due to the insertion of the aluminum foil was determined by making runs at $\Delta_{12} = 22^\circ$ both with and without the foil.

An energy calibration of 250 keV/channel was used in the two dimensional pulse height analysis. This was obtained by adjusting the amplifier gains until the proton peak corresponding to the transition through the ground state of Li^5 was centered about channel 38 for an incident energy of 1.15 MeV. For this energy and angle, $\Theta = 90^\circ$, this peak should appear at 9.46 MeV. The lower level discriminator of the S.C.A.'s was set to reject pulses corresponding to less than 1.25 MeV deposited in the detectors. This was necessary to remove the scattered He^3 events when no foil was placed in front of the detectors. The window

setting of the S.C.A.'s was opened to accept pulses up to 15 MeV.

The counting rates with detector configuration #3, an incident energy of 1.15 MeV, a beam current of 0.5 uamps and a target pressure of 100 mm. Hg were typically 100-150 counts/sec. for the singles rate and 5-10 counts/min. for the coincidence rate. The runs lasted until at least 1000 coincidence events along the desired kinematic contour were accumulated. A typical two dimensional spectrum is shown in Figure 15. The solid curve represents the kinematic contour for p-p coincidences for the angle and energy specified. The events below channel 10, lying well off the p-p contour are due to p-He⁴ and He⁴-p coincidences. The events lying well above the p-p contour are random coincidence events. With a coincidence resolving time of 50 nsec. the expected random rate is approximately 8 events/hour in agreement with the observed rate. Also shown in Figure 15 are the two energy histograms or coincidence proton spectra obtained by summing the p-p events lying within a nine channel interval along the kinematic contour. Because of the symmetry of the experiment these two coincidence spectra may be added to improve the statistics. Similar histograms, normalized to a total singles count in both detectors of 10⁶ events for detector configuration #1, are shown in Figures 16 - 21. In all cases the statistical error in the number of counts per channel is equal to or less than the square root of the number of counts in that particular channel.

B. Geometry #2 Measurements

These measurements were carried out using a 1.88 MeV He³⁺ beam from the University of British Columbia Van de Graaff

Figure 15. Two dimensional energy spectrum taken with Geometry #1 and $\Delta_{12} = 6^\circ$. Also shown are the energy histograms obtained by summing events along contour.

Figure 16. Coincidence proton spectra obtained by summing events along appropriate contour.

Figure 17. Coincidence proton spectra obtained by summing events along appropriate contour.

Figure 18. Coincidence proton spectra obtained by summing events along appropriate contour.

Figure 19. Coincidence proton spectra obtained by summing events along appropriate contour.

$\text{He}^3(\text{He}^3, 2\text{p})\text{He}^4$

$T_0 = 5.00 \text{ MeV}$

$\Delta_{12} = 180^\circ$

Figure 20. Coincidence proton spectra obtained by summing events along appropriate contour.

Figure 21. Coincidence proton spectra obtained by summing events along appropriate contour.

accelerator. The incident energy inside the gas target after correcting for the energy loss in the 0.025 mil nickel window was 1.50 MeV. From the kinematic calculations the detector angles which are required to observe the singlet two nucleon systems at the lowest possible relative energies for this geometry are $\Theta_1 = 62.0^\circ$ and $\Theta_2 = 90.0^\circ$ for the reaction $\text{He}^3(\text{He}^3, 2p)\text{He}^4$ and $\Theta_1 = 61.4^\circ$ and $\Theta_2 = 90.0^\circ$ for the reaction $\text{T}(\text{He}^3, np)\text{He}^4$. The difference is due to the slight difference in the reaction Q-values, 12.859 MeV for the reaction $\text{He}^3(\text{He}^3, 2p)\text{He}^4$ and 12.095 MeV for the reaction $\text{T}(\text{He}^3, np)\text{He}^4$.

Coincidence events in which the alpha particle enters detector 1 and the proton enters detector 2 are removed from the region of interest by the 0.200 mil nickel foil over the proton exit window and a 0.100 mil nickel foil in front of detector 1. The alpha particle exit window is 0.025 mils thick. The dimensions of the slit in front of both detectors for this series of measurements is 1/8" by 1/4" corresponding to an angular width in the polar direction $\Delta\Theta = 4^\circ$ and in the azimuthal direction $\Delta\Phi = 8^\circ$. These solid angles allowed sufficient statistics to be obtained for measurements at a particular pair of angles in three to six hours of running time. This time depended on the angles used, the beam current, target pressure and the target gas.

C. The $\text{He}^3(\text{He}^3, 2p)\text{He}^4$ Reaction.

The single particle spectra in both detectors from this reaction are shown in Figure 22. The peak at the high energy end of both spectra is due to the first emitted proton from the breakup through the ground state of Li^5 . This peak was used

$$T_0 = 1.50 \text{ MeV}$$

DETECTOR 1 SPECTRUM

$$\theta_1 = 90^\circ$$

DETECTOR 2 SPECTRUM

$$\theta_2 = 90^\circ$$

Figure 22. Single particle spectra in both detectors from the reaction $\text{He}^3(\text{He}^3, 2\text{p})\text{He}^4$ at 90° to the beam direction.

to set the gains of both main amplifiers so that the energy calibration was 200 keV/channel. The low energy peak in the spectrum from detector 2 is due to alpha particles and protons from the breakup of the Li^5 ground state. This peak does not appear in the detector 1 spectrum due to the nickel foils in front of this detector. The broad continuum in both spectra is made up of contributions from the sequential decay through the first excited state of Li^5 and the singlet di-proton state and from the direct three particle breakup. The small peak about channel 32 in the detector 2 spectrum is due to protons from the contaminant reaction $\text{He}^3(\text{d},\text{p})\text{He}^4$. Although these protons have an energy of 15.0 MeV at 90° only about 6 MeV is deposited in the alpha detector because of its depletion thickness. This peak is not observed in the same place in the detector 1 spectrum as this detector has a large enough depletion thickness to stop a 14 MeV proton and therefore the peak appears above channel 64. The windows of the single channel analysers were both set to accept particles with energies from 1 MeV to 12.8 MeV.

A two dimensional contour plot of the coincidence events observed for the case of $\Theta_1 = 62^\circ$ and $\Theta_2 = 90^\circ$ is shown in Figure 23. The events along contour A are the desired p- He^4 coincidences. The events along contour B are due to p-p coincidences and in the region marked C due to He^4 -p coincidences in which the alpha particle energy has been degraded by the foils in front of detector 1. The few events in the region marked D are from p- He^4 coincidences from the contaminant reaction $\text{He}^3(\text{d},\text{p})\text{He}^4$.

Figure 23. Two dimensional contour plot of p-p and p- He^4 coincidence events from the reaction $\text{He}^3(\text{He}^3, 2p)\text{He}^4$ at $\Delta_{12} = 152^\circ$.

In this case the deuterium is present as a HD^+ component of the He^3 beam and because of the high cross section for this reaction a few nanoamps of HD^+ produces a significant coincidence rate. Although this reaction leads to a two particle final state and therefore only a particular configuration of the two detectors will observe both particles simultaneously, it is found that this configuration is almost exactly the same configuration required to observe the p-p interaction. Fortunately the separation can be made by using the proton energy as the events from the contaminant reaction have energies $T_p = 16.0$ MeV and $T_\alpha = 3.3$ MeV for this geometry and therefore lie well off the desired contour. However some of the protons lose part of their energy by slit scattering and these produce the coincident events in the region D. Measurements were made with a pure deuterium beam to simulate the HD^+ component in the He^3 beam and these showed that no coincidence events from this reaction produce events in the portion of contour A where the p-p interaction is expected to appear.

The spectrum of protons from detector 1 in coincidence with alpha particles in detector 2 is obtained by summing the coincidence events lying within an eight channel interval of the calculated p- He^4 contour. This spectrum was corrected for the contribution from the reaction $\text{He}^3(d,p)\text{He}^4$ by using the results of the run with the pure deuterium beam to obtain the shape of the contribution. This spectrum is shown in Figure 24. A similar run was made for detector angles $\theta_1 = 75^\circ$ and $\theta_2 = 90^\circ$ and the coincidence proton spectrum obtained by summing over the p- He^4 contour is shown in Figure 25. Both these spectra are normalized

$\text{He}^3(\text{He}^3, 2\text{p})\text{He}^4$

$T_0 = 1.50 \text{ MeV}$

$\theta_1 = 62^\circ \quad \theta_2 = 90^\circ$

Figure 24. Coincidence proton spectrum from the reaction $\text{He}^3(\text{He}^3, 2\text{p})\text{He}^4$, obtained by summing over p-He⁴ contour for $\Delta_{12} = 152^\circ$.

Figure 25. Coincidence proton spectrum from the reaction
 $\text{He}^3(\text{He}^3, 2\text{p})\text{He}^4$ obtained by summing over p-He⁴ contour
 for $\Delta_{12} = 165^\circ$.

to the same number of single counts (2.5×10^6) in the fixed detector. For the angles $\Theta_1 = 75^\circ$ and $\Theta_2 = 90^\circ$ the coincidence events from the contaminant reaction are negligible as this geometry does not permit the alpha particle and proton to be observed simultaneously in both detectors unless one or both of the particles have been scattered.

D. The $T(\text{He}^3, np)\text{He}^4$ Reaction

The single particle spectra from this reaction are shown in Figure 26 for both detectors at 90° to the incident beam direction. The same energy calibration of 200 keV/channel is used for these measurements. The possible processes contributing to these spectra are the following:

	<u>Q (MeV)</u>
(1) $\text{He}^3 + T \rightarrow \text{He}^4 + d$	14.320
(2) $\rightarrow \text{Li}^5 + n \rightarrow \text{He}^4 + p + n$	10.128
(3) $\rightarrow \text{He}^5 + p \rightarrow \text{He}^4 + n + p$	11.138
(4) $\rightarrow \text{He}^4 + n + p$	12.095
(5) $\text{He}^3 + D \rightarrow \text{He}^4 + p$	18.352

The positions of the peaks expected from the various particles and possible final states are shown above each spectrum. The number refers to the process and the subscript to the particle detected. The particle energies have been corrected for the energy loss in the nickel foils. The 0.300 mil thickness of nickel in front of detector 1 prevents the low energy structure from being observed. The continuum in both spectra consists mainly of contributions from the sequential decay through the first excited state of Li^5 and He^5 .

Figure 26. Single particle spectra in both detectors from the reaction $T(\text{He}^3, np)\text{He}^4$ at 90° to the beam direction.

A two dimensional contour plot of the coincidence events for $\Theta_1 = 61.4^\circ$ and $\Theta_2 = 90^\circ$ is shown in Figure 27. As before, the events lying along contour A are the desired p-He⁴ coincidences. The events in region B are from He⁴-p coincidences in which the alpha particle energy has been degraded by the foils in front of detector 1 and the peak in this region is due to He⁴-d coincidences from the two particle final state T(He³,d)He⁴. The events in region C and along the line of constant energy T₂ are d-He⁴ coincidences from the same reaction. Again as this is a two particle final state both particles will only be observed for a particular detector configuration. This is approximately the same configuration as that required to observe the n-p interaction. The events along the line of constant T₂ are due to coincidences in which the deuteron has lost some of its energy by slit scattering. This effect produces an appreciable contribution to the coincidence events as the two particle final state consisting of the deuteron and alpha particle is the dominant decay mode of the reaction. This may be seen in the single particle spectra. The peak channel from this process is some three orders of magnitude above the average contribution per channel from the three particle final states. Considerable effort was made to reduce the effect of slit scattering by using accurately machined slits but it was found impossible to remove it completely. Particle identification techniques may be used to remove this contribution to the two dimensional coincidence spectra. These improvements are discussed in the final chapter. For the purpose of the experiment considered here, that is the observation of the two nucleon interactions, this contri-

Figure 27. Two dimensional contour plot of p- He^4 and d- He^4 coincidence events from the reaction $T(\text{He}^3, np)\text{He}^4$ at

$$\Delta_{12} = 151.4^\circ$$

bution does not affect the results.

The contaminant reaction $D(\text{He}^3, p)\text{He}^4$ also presents a problem in these measurements. This time the deuterium is present in the tritium gas target. There is not sufficient He^3 in the tritium to produce any significant p-p coincidences from the $\text{He}^3(\text{He}^3, 2p)\text{He}^4$ reaction so it is unlikely that it is the HD^+ component in the beam producing the contaminant reaction. Although the p- He^4 coincidence events from this reaction do not appear in the region where the n-p interaction is observed they do present a problem in determining the contribution from the states in He^5 . The technique used to remove the contribution from this reaction was to make the same measurements at this pair of angles for two settings of the bias level of detector 1. The peak from the high energy protons from the contaminant reaction thus appeared at different places in the spectrum and could be subtracted out.

The coincidence proton spectrum obtained by summing over an eight channel interval along the calculated p- He^4 contour and subtracting the contribution from the $D(\text{He}^3, p)\text{He}^4$ reaction is shown in Figure 28. The similar spectrum obtained for the detector positions $\Theta_1 = 75^\circ$ and $\Theta_2 = 90^\circ$ is shown in Figure 29. In this case the contaminant reaction does not present a problem. Again both these spectra are normalized to 2.5×10^6 single counts in the fixed detector.

The theoretical interpretation of these spectra and the spectra from the reaction $\text{He}^3(\text{He}^3, 2p)\text{He}^4$ is discussed in the

Figure 28. Coincidence proton spectrum from the reaction $T(\text{He}^3, np)\text{He}^4$ obtained by summing over p-He⁴ contour for $\Delta_{12} = 151.4^\circ$

Figure 29. Coincidence proton spectrum from the reaction $T(\text{He}^3, np)\text{He}^4$ obtained by summing over p-He⁴ contour for $\Delta_{12} = 165^\circ$

next chapter. In Chapter VI, improvements to the experimental technique are considered which will remove some of the ambiguity resulting from the two particle final states, both from the contaminant reaction and the decay mode $T(\text{He}^3, d)\text{He}^4$.

CHAPTER V

THEORY OF FINAL STATE INTERACTIONS

A. Introduction

The test of a good nuclear theory is its usefulness first of all in extracting nuclear parameters from experimental information and then in predicting what to expect from further experiments on the basis of these results. At the present time no really good theory exists for the three body problem although much of the mathematical groundwork has been laid for the development of such a theory. The immediate applications in low energy nuclear physics and particle physics of a precise three body theory are sufficient to warrant considerable time and effort being spent on this problem. As this thesis is intended to be mainly experimental in content, no complete history of the mathematical techniques used in tackling the three body problem will be given. However two of the approximate methods developed to explain certain features of three particle breakup reactions will be discussed in this chapter along with some of the recent developments of the theory. The accuracy of the two theories will be checked by comparing their predictions with the experimental spectra measured in the reactions $\text{He}^3(\text{He}^3, 2p)\text{He}^4$ and $\text{T}(\text{He}^3, np)\text{He}^4$.

Watson (16) in 1952 introduced the term "final state interaction" and showed that these interactions can greatly enhance transitions where two nucleons emerge in a singlet state of low relative energy. His approach was to separate the

reaction process into a "primary mechanism" in which the final state interaction plays no important part and a "final state interaction" which distorts the energy and angular distributions of the outgoing particles before they can get outside the range of their mutual forces. In the case of two nucleons in the final state, i.e. a reaction of the type

the interaction operator in the Schrodinger equation is separated into two terms, $v + V$, where v is the two nucleon interaction which leads to the correct scattering cross section for the particular pair of particles. The transition amplitude may be written

$$T_{fi} = \langle \phi_f^{(-)} | V | \psi_i^{(+)} \rangle \quad 5-1$$

where $\phi_f^{(-)}$ is the final state wave function under the interaction of v alone. Watson's main assumption is that for k , the relative nucleon momentum, small, the transition amplitude has the same k -dependence as $\phi_f^{(-)*}$.

R.J.N. Phillips (25) uses the exact outside wave function for the two nucleon case and shows that the spectrum of particle X in the breakup may be approximated by

$$\frac{d^2\sigma}{d\Omega_x dE_x} = \frac{W_0}{k} \left[F(kb) \cos \delta + G(kb) \sin \delta \right]^2 \quad 5-2$$

where F and G are the regular and irregular solutions of the radial wave equation evaluated at a radius ' b ', W_0 contains the factors which are not momentum dependent and δ is the S-wave nucleon-nucleon scattering phase shift.

A somewhat different approach is used by G.C. Phillips, Griffy and Biedenharn (8). They express the cross section for the process

where B is formed in a state unstable to particle emission, i.e.

$B \rightarrow c + C$, by

$$\sigma_l(E_B) = \frac{\mu_a \mu_b k_b}{4\pi^2 \hbar^4 k_a} \left| \langle B+b, E_b | H' | A+a, E_a \rangle \right|^2 \rho_l(E_B) \quad 5-3$$

where H' is the interaction Hamiltonian, $|A+a, E_a\rangle$ represents the initial state vector and $|B+b, E_b\rangle$ the final state vector. The initial and final wave numbers are k_a and k_b , and $\rho_l(E_B)$ is called the generalized density of states function for B. For the case of B having only a set of sharp states at energies E_n this function may be expressed as

$$\rho(E_B) = \sum_n \delta(E - E_n) \quad 5-4$$

The density of states function for a system B which is unstable to decay into two particles $c+C$ is obtained by evaluating the number of discrete states per unit energy interval of B lying within two spheres, one with the interaction radius 'b' and the other with an arbitrarily large radius 'R'. This leads to the result

$$\rho_l(E_B) = \frac{1}{\pi} \frac{d}{dE_B} \left(\delta_l(E_B) + \phi_l(E_B, b) \right) \quad 5-5$$

where δ_l is the phase shift for the corresponding elastic scattering of $c+C$ and ϕ_l is the hard sphere phase shift eval-

uated at the radius 'b'.

It is easy to show that the results of Watson and Phillips, Griffy and Biedenharn (hereafter referred to as the PBG theory) lead to a similar expression for the cross section for the case where the scattering process has a resonance with phase shifts which may be described by single level dispersion theory. Then the scattering phase shift may be written as

$$\delta_l = \beta_l - \phi_l$$

where
$$\beta_l = \tan^{-1} \left[\frac{\frac{1}{2} \Gamma_l}{(E_l + \Delta_l - E)} \right] \quad 5-6$$

with
$$\frac{1}{2} \Gamma_l = e \gamma_l^2 / (F_l^2 + G_l^2) = e \gamma_l^2 / A_l^2$$

and
$$\Delta_l = -[\gamma_l^2 / e] \left[(e / A_l) (dA_l / d\rho) + l \right]$$

with $e = kb$. γ_l^2 and E are respectively the reduced width and the characteristic energy used to describe the particular resonance.

Watson's theory as expressed in equation 5-2 may also be written as

$$\sigma_l(E)_{\text{WATSON}} = \frac{\text{const.}}{P_l} \sin^2(\delta_l + \phi_l) \quad 5-7$$

This relationship is true for any form of the phase shifts with P_l and ϕ_l defined as follows:

$$P_l = \frac{e}{F_l^2 + G_l^2} \quad 5-8$$

and
$$\phi_l = \tan^{-1} \frac{F_l}{G_l} \quad 5-9$$

If the phase shifts are described by the single level approximation then equation 5-7 becomes

$$\sigma_{\ell}(E)_{\text{WATSON}} = \frac{\text{const. } \frac{1}{2} \Gamma_{\ell}}{(E_{\ell} + \Delta_{\ell} - E)^2 + (\frac{1}{2} \Gamma_{\ell})^2} \quad 5-10$$

Similarly on differentiating the resonant phase shift 5-6 the cross section obtained from the density of states formalism is given by

$$\sigma_{\ell}(E)_{\text{FBG}} = \frac{\text{const. } \frac{1}{2} \Gamma_{\ell}}{(E_{\ell} + \Delta_{\ell} - E)^2 + (\frac{1}{2} \Gamma_{\ell})^2} \left[1 + \frac{d\Delta_{\ell}}{dE} + \frac{E_{\ell} + \Delta_{\ell} - E}{\Gamma_{\ell}} \frac{d\Gamma_{\ell}}{dE} \right] \quad 5-11$$

The two results are equivalent if Δ_{ℓ} is a linear function of energy and Γ_{ℓ} is independent of energy. Barker and Treacy (26) have shown that this is approximately true for energies well above threshold but for energies near threshold or for phase shifts not exhibiting a resonance the two formalisms produce different results. This will be shown in the next section where both formalisms are used to fit the experimental results.

Another approximation method has been used by Duck (27) explicitly for the three alpha and three nucleon systems and for a coincidence measurement of two of the three final state particles. In the case of the three alpha decay of C^{12} the triple correlation cross section has the form

$$\frac{d^3\sigma}{d\Omega_1 d\Omega_2} = \left| \frac{P_1 P_2^2}{A P_2 + P_1 \cos \Delta_{12} - P_0 \cos \Theta_2} \right| \overline{M}_{JJ'}^2 \quad 5-12$$

where the first term is the phase-space factor described in Appendix A and $\overline{M}_{JJ'}^2$ is the square of the matrix element for C^{12} (spin J) to breakup through Be^8 (spin J') and contains all the

angular dependence. This matrix element was evaluated using the Breit-Wigner resonance theory, the Watson theory of final state interactions and the PBG density of states formalism. All theories predict the distinctive features of the experimental spectra although peak widths and positions are not predicted with any accuracy. A similar expression has also been obtained for the p-D inelastic triple correlation by using a modified impulse approximation. The form of the interactions included in the approximation is that obtained from the Watson theory.

A more "exact" treatment of the three nucleon system has been made by Amado and his co-workers (28,29) based on the fundamental theory of Fadeev and Lovelace. The three body effects are taken completely into account including inelastic processes and the couplings between channels. In order to include these effects and still keep the problem to manageable proportions it is necessary to use a simplified form for the nucleon-nucleon interaction.

The force between the nucleons is taken to arise from a sum of separable interactions, in this case two S-wave interactions are used, one in the spin zero, isospin one state and the other in the spin one, isospin zero state. Even with the simplified form for the interactions the exact treatment of the three body effects requires the solution of a set of coupled integral equations which can only be obtained with a high speed computer. The differential proton spectrum for the inelastic scattering

has been calculated and the general features of the experimental data are reproduced by the theory. The magnitudes are not so well

reproduced but the agreement does indicate considerable promise in this type of treatment of the three body problem. Amado and Aaron (29) also compare their theory with the Watson theory for final state interactions in the region of low relative neutron-neutron momentum. They show that although the assumption used by Watson, that in this region only the interaction between the neutrons is important, is not valid if one considers the exact treatment, the Watson theory gives results which closely resemble the exact theory. The similarity in the results is obtained by using a slightly larger scattering length in the case of the Watson theory. The question of whether this resemblance is accidental or not is presently under investigation. One result which is evident from Amado's work is that at the present time the theory, and probably the experimental information on this reaction, are not sufficiently precise to differentiate between n-n scattering lengths of -17 fm. and -24 fm.

B. Analysis of Coincidence Proton Spectra

The quantity which has been determined in the measurement of the coincidence proton spectra is the triple correlation distribution

$$\sigma_{lab}(T_1) = \frac{d^3\sigma}{dT_1 d\Omega_1(lab) d\Omega_2(lab)} \quad 5-13$$

If the reaction proceeds via a sequential decay through an intermediate two particle state, then the coordinate system of interest in deriving a theoretical expression for 5-13 is the recoil center of mass system of the corresponding two particles. The cross section in the rcm(i) system for a breakup involving the cluster

(j+k) is given by

$$\sigma_{\text{rcm}}^{(i)} = \frac{d^3\sigma}{dE_{jk} d\Omega_i(\text{scm}) d\Omega_j(\text{rcm})} \quad 5-14$$

where E_{jk} is the internal or excitation energy of the cluster (j+k) and i,j,k is some permutation of 1,2,3. The Jacobian required to transform this cross section to the laboratory system is

$$J_{iL} = \frac{\partial(E_{jk}; \Omega_i(\text{scm}); \Omega_j(\text{rcm}))}{\partial(T_1; \Omega_1(\text{lab}); \Omega_2(\text{lab}))} \quad 5-15$$

The Jacobian for this transformation is calculated in Appendix A.

Several approximations are made in determining the quantity $\sigma_{\text{rcm}}^{(i)}$. The first is that the breakup is isotropic in the system center of mass. For the purpose of the calculations here this is not a bad approximation as in the first series of measurements with Geometry #1 the polar angles of both detectors remain constant and in the second series with Geometry #2 the variation in polar angle is less than 15° . Since the laboratory velocities of the final state particles are in general considerably larger than the incident velocity the scm. polar angles are not too different from the lab. polar angles and therefore remain approximately the same over most of the range of measurements.

The second approximation is that the breakup of the intermediate state is isotropic in its own center of mass system. This requires that the intermediate system lasts sufficiently long that any spatial localization of its constituent particles resulting from its formation, is lost before it decays so that there is no preferred direction of breakup. The third approximation is that

the contributions from the various intermediate states are considered to be simply additive, i.e. there are no interference effects. It is realized that there are no good arguments to support these approximations. However the purpose of these calculations is mainly to provide a comparison of the predictions of the simple theories of final state interactions with the results of the experimental measurements, with the hope of indicating the directions that a more precise theory must take. With these approximations the cross section $\sigma_{\text{rcm}}^{(1)}$ for a particular two particle system depends only on the excitation energy of the system and may be evaluated using either the Watson theory or the PBG theory. In both cases the phase shifts for the corresponding elastic scattering must be known. The following section indicates how these quantities are evaluated.

p-He⁴ and n-He⁴ Phase Shifts.

The elastic scattering of nucleons by He⁴ has been extensively studied mainly on account of the use of helium targets as a polarization analyser. Therefore the phase shifts are known quite well. The P-wave phase shifts δ_1^+ and δ_1^- which correspond to the first two states of the mass 5 system may be adequately fitted using single level dispersion theory. The form of the phase shifts predicted by this theory is given in equation 5-6. From the data on p-He⁴ scattering reported by Barnard et al (30) the best fits for the states of Li⁵ are obtained with the set of parameters listed in Table 4. Similarly the experimental n-He⁴ P-wave phase shifts of Hoop and Barschall (31) are used to obtain the parameters for the first two states of He⁵. These also are

TABLE 4

Parameters obtained for states of Li^5 and He^5 from single level dispersion theory.

State J^π	Phase shift	b (fm.)	E_ℓ (MeV)	χ_ℓ^2 (MeV)
Li^5 ground state $3/2^-$	δ_1^+	3.0	4.60	8.15
Li^5 excited state $1/2^-$	δ_1^-	3.0	19.79	15.28
He^5 ground state $3/2^-$	δ_1^+	2.9	3.10	8.50
He^5 excited state $1/2^-$	δ_1^-	2.9	16.0	12.5

TABLE 5

Effective range theory parameters for nucleon-nucleon scattering phase shifts.

State	Phase shift	a (fm.)	r_0 (fm.)	P
n-p singlet state	δ_0	-23.71	2.40	.
p-p singlet state	δ_0	-7.75	2.78	0.041

listed in Table 4. It should be pointed out that the large difference between the value of E , the characteristic energy, and the resonance energy of the particular level is due to the definition of $\Delta\epsilon$, the energy shift, in this analysis.

n-p Phase Shifts

The phase shift analysis of nucleon-nucleon scattering is discussed in most textbooks on Nuclear Physics, for example Preston (32). The singlet or triplet state n-p phase shifts may separately be expanded in a Taylor's series in momentum, for which the lowest two terms are

$$k \cot \delta = -\frac{1}{a_n} + \frac{1}{2} r_{on} k^2 \quad 5-16$$

where k is the relative nucleon momentum, a_n is called the scattering length and r_{on} is called the effective range. The value of a_n and r_{on} which best fit the present experimental data are given in Table 5. The hard sphere phase shifts are obtained from the solutions of the radial wave equation. The appropriate radial functions for the n-p singlet scattering are as follows

$$\begin{aligned} F_0(\rho) &= \sin \rho \\ G_0(\rho) &= \cos \rho \end{aligned}$$

and using relation 5-9 the hard sphere phase shift is simply given by

$$\phi_0 = \rho = kb \quad 5-17$$

where b is the radius at which the hard sphere phase shift is evaluated.

p-p Phase Shifts

The phase shift analysis of p-p elastic scattering is complicated by the effect of the Coulomb field. The momentum dependence of the singlet state phase shifts is given by the following generalization of 5-16

$$C^2 k \cot \delta + \frac{1}{R} h_d(\eta) = -\frac{1}{a_p} + \frac{1}{2} r_{op} k^2 - P r_{op}^3 k^4 \quad 5-18$$

where

$$C^2 = \frac{2\pi\eta}{e^{2\pi\eta} - 1}$$

$$h_d(\eta) = \eta^2 \sum_{n=1}^{\infty} \frac{1}{n(n^2 + \eta^2)} - \ln \eta - \gamma$$

$$\gamma = 0.5772$$

$$R = 2.8815 \times 10^{-12} \text{ cm.}$$

and

$$\eta = \frac{Z_1 Z_2 e^2}{\hbar v}$$

The least squares fit to the experimental data gives the values for the parameters a_p , r_{op} and P listed in Table 5. The Coulomb functions appropriate here are given approximately as follows

$$F_0(\rho) = C^2 \rho (1 + \eta \rho)$$

$$G_0(\rho) = \frac{1}{C^2} \left[1 + 2\eta \rho (\ln 2\eta \rho + 2\gamma - 1 + h_0(\eta)) \right]$$

where $\rho = kb$ as before and ϕ_0 is obtained using relation 5-9.

It is now possible to determine the energy dependence of the cross section $\sigma_{\text{rcm}}^{(i)}$ for any of the two particle systems consisting of a proton, neutron or alpha particle. Apart from numerical constants the cross section predicted by the Watson theory has the form

$$\sigma_{rcm}^{(i)} \text{ WATSON} = \frac{\sin^2(\delta + \phi)}{P_l} \quad 5-19$$

and by the PBG theory, the form

$$\sigma_{rcm}^{(i)} \text{ PBG} = \frac{d}{dE_k} (\delta + \phi) \quad 5-20$$

The results of the evaluation of these quantities using the phase shifts described previously are shown in Figures 30 and 31 for the two states of Li^5 and He^5 and in Figures 32 and 33 for the singlet two proton state and the singlet deuteron. It is evident that there is considerable difference in the predictions of the two theories. Even in the case of the states of Li^5 and He^5 where the phase shifts follow single level dispersion theory the distributions are similar only near the ground state peak. The greatest difference is in the contribution to the excitation distribution well off resonance, with the Watson theory predicting a much larger contribution. For the nucleon-nucleon systems there is only a slight similarity in the shape of the excitation functions predicted by both theories.

The coincidence proton spectrum in the lab. system is obtained by multiplying the $\text{rcm}(i)$ cross sections for the possible intermediate state by the appropriate Jacobian and by a constant which determines its contribution relative to the other intermediate states. In the measurements with Geometry #1 the contribution from the various final state interactions to the coincidence proton spectra are separated kinematically so the analysis of these spectra is somewhat simplified. For this reason

Figure 30. Excitation functions for states of Li^5 predicted by Watson and PBG theories.

Figure 31. Excitation functions for states of He^5 predicted by Watson and PBG theories.

Figure 32. Excitation functions for singlet state of p-p system predicted by Watson and PBG theories.

Figure 33. Excitation functions for singlet state of n-p system predicted by Watson and PBG theories.

this analysis is considered first.

C. Geometry #1 Analysis of Results

For this geometry and for an angular separation of the detectors $\Delta_{12} = 180^\circ$ or 135° the contribution from the final state interaction between the protons is negligible as coincidence events observed with this configuration correspond to very high excitations in the p-p system. The lab. cross section may be written

$$\sigma_{lab}(T_1) = a^+(J_{1L}\sigma_+(E_{23}) + J_{2L}\sigma_+(E_{13})) + a^-(J_{1L}\sigma_-(E_{23}) + J_{2L}\sigma_-(E_{13})) \quad 5-21$$

where E_{23}, E_{13} both correspond to excitation energies in the Li^5 system, $p \rightarrow 1$, $p \rightarrow 2$ and $He^4 \rightarrow 3$. Since it is not known whether detector 1 or detector 2 observes the first emitted proton from the breakup through Li^5 , both contributions must be included. The quantities σ_+ and σ_- are the rcm. cross sections for the ground state (+) and first excited state (-) of Li^5 obtained using either the Watson or PBG theory and J_{1L} and J_{2L} are the Jacobians for the transformation of these cross sections to lab. coordinates. The quantities a^+ and a^- are the relative intensities of the two states and are assumed to be dependent only on the incident He^3 energy.

The procedure followed in determining the lab. cross section from expression 5-21 is to evaluate the quantities J_{1L} , J_{2L} , E_{23} and E_{13} for the appropriate detector angles and incident energy as a function of T_1 . The computer program KINJAC used to make this calculation is listed in Appendix B. The rcm. cross sections $\sigma_+(E_{23})$, $\sigma_+(E_{13})$, $\sigma_-(E_{23})$, $\sigma_-(E_{13})$ are then determined

from the excitation functions shown in Figure 30. The relative intensities were chosen to give the best fit to the observed spectrum for $\Delta_{12} = 180^\circ$. The results of these calculations and the experimental spectra are shown in Figure 34 for the incident energy $T_0 = 1.15$ MeV and in Figure 35 for the incident energy $T_0 = 5.00$ MeV. The solid curves correspond to the Watson theory and the dashed curves to the PBG theory. Also shown are the excitation energies E_{23} and E_{13} as a function of the proton energy T_1 .

In order to obtain the best fit to the experimental spectrum for $T_0 = 5.00$ MeV using the Watson theory it is necessary to set $a^- = 0$, i.e. no contribution from the excited state of Li^5 . The tail of the excitation distribution for the Li^5 ground state predicted by this theory is more than enough to account for the continuum between the ground state peaks. The Watson theory does predict the position of the ground state peak more accurately than the PBG theory.

An obvious shortcoming of this simple analysis is that it does not produce the required variation with the angle Δ_{12} as may be seen in Figure 34b for $\Delta_{12} = 135^\circ$. These curves are calculated with the same intensities a^+ and a^- used to fit the spectrum at 180° . This indicates that the assumption that the intermediate state breaks up isotropically in its own center of mass system must not be valid. The angle in the rcm. system between the recoil velocity of the Li^5 and the velocity of the breakup proton (either $\eta_2^{(1)}$ or $\eta_1^{(2)}$ in the discussion of the

Figure 34. Comparison of experimental coincidence proton spectra with Watson and PBG theories for $T_0 = 1.15 \text{ MeV}$.

Figure 35. Comparison of experimental coincidence proton spectrum with Watson and PBG theories for $T_0 = 5.00 \text{ MeV}$.

kinematics in Appendix A) may be evaluated at each proton energy T_1 . For the case of the first emitted proton entering detector 1 and $T_1 = 9.5$ MeV, the angle $\eta_2^{(1)} = 23.3^\circ$ for $\Delta_{12} = 180^\circ$ and $\eta_2^{(1)} = 70.0^\circ$ for $\Delta_{12} = 135^\circ$. The difference in $\eta_2^{(1)}$ for the two cases is fairly constant for all values of T_1 . Since fewer events are seen for the larger breakup angle it appears that the proton from the breakup of the Li^5 emerges preferentially in the recoil direction. This is not an unexpected result as in most triple angular correlations of the type (a,b γ) initiated by direct reactions the preferred axis is the recoil direction of the intermediate particle (33). An opposite effect has been observed for the Li^5 ground state by Reimann (19) in the reaction $\text{Li}^6(\text{He}^3, p)2\text{He}^4$. Here an axial asymmetry is found about the recoil direction of the Li^5 in the breakup of the ground state. In this case the emission of the proton is preferentially in the forward direction with respect to the beam axis when all kinematic effects are taken into account.

Another interesting feature of the measurements shown here is the rapid decrease in the amount of continuum between the ground state peaks as the incident energy increases. Using the PBG theory the ratio a^+/a^- required to fit the observed spectra increases by a factor of 5.4 from $T_0 = 1.15$ MeV to 5.00 MeV. The Watson theory requires an even more drastic change as $a^- = 0$ for $T_0 = 5.00$ MeV. Regardless of the theory, if it is assumed that only the Li^5 states contribute to the coincidence events in this region, the ratio of the ground state intensity to the excited state

intensity in the reaction $\text{He}^3(\text{He}^3, 2\text{p})\text{He}^4$ must increase by a factor of at least four from $T_0 = 1.15$ MeV to 5.00 MeV. This increase is uniform with incident energy as may be seen in the spectrum for $T_0 = 2.40$ MeV in Figure 18.

Attempts were made to explain this increase by considering that the ground state $J^\pi = 3/2^-$ might be populated preferentially to the excited state $J^\pi = 1/2^-$ by higher order angular momenta in the incoming beam. Although this effect produces a change in the relative intensity in the right direction it can at most account for an increase of 30%. This result is obtained by assuming the low energy beam is pure S-wave and the high energy beam is P-wave. Another possible explanation is that the effect is due to the preferential breakup of the proton from the Li^5 in the recoil direction. The breakup angle $\eta_2^{(1)}$ is found to increase slightly with increasing excitation energy in the Li^5 system but it is found that this increase is practically the same for the different incident energies. Therefore it does not seem likely that the observed change in the relative intensities is due to some kinematic effect.

Since it does not appear that the states of Li^5 can account for the observed coincidence spectra the obvious conclusion is that there is some contribution from another decay mode. The only other process which can produce coincidence events in this region is the instantaneous three particle breakup in which the final state interactions do not play a part. Figure 36 shows three curves, one is the coincidence proton spectrum for a statis-

Figure 36. Comparison of predictions of statistical breakup with Watson and PBG theories for the Li^5 excited state.

tical breakup obtained from equation A-28 and the other two are the spectra predicted by the Watson and PBG theories for the first excited state of Li^5 . These distributions are quite similar in the region $T_1 = 4 \text{ MeV}$ to 8 MeV so it would be difficult to separate the contribution of the Li^5 excited state from the statistical breakup. If it is assumed that part of the continuum between the ground state peaks for $T_0 = 1.15 \text{ MeV}$ is due to the statistical breakup and that this contribution decreases relative to the Li^5 ground state as the incident energy increases then it is possible to account for the observed spectra.

The final state interaction between the protons is important in explaining the coincidence proton spectra measured at small angular separations of the two detectors with this geometry. In the preliminary analysis of the spectra for $\Delta_{12} = 6^\circ$ to $\Delta_{12} = 22^\circ$ the contribution of the states of Li^5 and the statistical breakup is neglected. Reasons for not including these processes will be given later. The lab. cross section may then be written

$$\sigma_{\text{lab}}(T_1) = a^0 J_{3L} \sigma_0(E_{12}) \quad 5-22$$

where E_{12} is the corresponding excitation in the p-p system, J_{3L} is the Jacobian for the transformation to lab. coordinates and σ_0 is the rcm. cross section for the singlet two proton state determined from the excitation functions shown in Figure 32. The results of the evaluation of equation 5-22 for the two theories is shown in Figure 37 superimposed on the corresponding experimental spectra. As before the solid curve corresponds to the Watson

Figure 37. Comparison of experimental coincidence proton spectra at small angles with Watson and PBG theories.

theory and the dashed curve to the PBG theory. The normalization constant a^0 was chosen to fit the observed spectrum at $\Delta_{12} = 6^\circ$ for each theory and the same constant is used for the calculation of the spectrum at $\Delta_{12} = 22^\circ$. Some of the excitation energies E_{12} of the two proton system corresponding to the lab. energy T_1 are shown above each spectrum. The minimum excitation energy reached with this geometry (neglecting the width of the detectors) is 24 keV.

From these curves it is evident that the Watson theory gives the best fit to the observed coincidence spectra when the p-p final state interaction is important. However it does predict the peak in the p-p excitation function at too low an energy. The effect of changing the scattering length a_p is shown in Figure 37c. A smaller scattering length in the Watson theory gives a better fit but it still does not account for the rapid decrease at low excitation energies. This is opposite to the observation of Amado and Aaron (29) who found that in the case of the singlet deuteron the Watson theory resembles their exact theory if a larger scattering length is used. The predictions of the PBG theory fail to account for both the peak positions and the change with angle.

The effect of including the contributions from the states of Li^5 and the statistical breakup would be to add a flat structureless background. The minimum excitation energy reached in the Li^5 system for $\Delta_{12} = 6^\circ$ is about 3.0 MeV so the ground state peak is not present. The maximum amount that these processes can

contribute is determined by the dip at 4.5 MeV so that the conclusions reached in this analysis would not be affected by including this contribution.

The coincidence spectra observed for these angles at the higher incident energy are very similar as may be seen in Figures 19 and 21. However it is apparent that the formation of the singlet two proton state decreases in importance relative to the Li^5 ground state as the incident energy increases. It is for this reason that the study of this reaction at higher incident energies by Bacher (9) and Aldridge (7) did not reveal good evidence for the presence of the p-p final state interaction. The kinematic calculations show that this decrease is not a phase-space effect, as the ratio of the available phase-space for the two processes is approximately constant for incident energies from 1.15 MeV to 5.00 MeV. One of the aims of a more accurate theoretical analysis of this breakup should be to explain this effect.

D. Geometry #2 Analysis of Results

The analysis of the coincidence spectra obtained with Geometry #2 is complicated by the fact that the contribution of both the states of Li^5 or He^5 and the two nucleon system must be considered together. The procedure used in this analysis was to determine the contribution from the mass 5 systems by fitting the region of the spectrum where this process is predominant. This contribution was then subtracted from the measured spectrum, hopefully leaving the portion due only to the two nucleon singlet state. The PBG theory was used to determine the shape of the Li^5

and He^5 contributions as it gives a better estimate of the amount to be included for excitation energies lying well off the resonant part of the excitation function. The contribution from the statistical three particle breakup is included as part of the contribution from the excited state of the mass 5 nuclei. The ratio $a^+/a^- = 0.33$ for the incident energy $T_0 = 1.50$ MeV was determined for the Li^5 states by fitting the p-p coincidence spectra, which are obtained simultaneously in the reaction $\text{He}^3(\text{He}^3, 2p)\text{He}^4$.

Figure 38 shows the coincidence proton spectrum (in coincidence with alpha particles) obtained for $\Theta_1 = 62^\circ$ and $\Theta_2 = 90^\circ$ divided by the available phase-space as calculated from equation A-28. The effect of the available phase-space on the coincidence spectra measured with this geometry, is to greatly magnify the contribution at the high energy end of the spectrum. Removing this purely kinematic effect by using the ratio of the observed yield to the available phase-space gives a more accurate picture of the relative intensities of the processes contributing to the spectrum. This was not necessary in the case of the measurements made with Geometry #1 as the available phase-space calculated for this geometry does not distort the observed spectra to such an extent. The contribution from the states in Li^5 used in the background subtraction, similarly corrected for the available phase-space, is also shown in Figure 38. The curves marked E_{12} and E_{23} are the corresponding excitation energies in the Li^5 system. A similar spectrum was calculated for the case $\Theta_1 = 75^\circ$ and $\Theta_2 = 90^\circ$. The spectra obtained by subtracting these background

Figure 38. Coincidence proton spectrum from the reaction $\text{He}^3(\text{He}^3, 2p)\text{He}^4$ for $\Delta_{12} = 152^\circ$ divided by the available phase-space. Also shown is the contribution from the states of Li^5 calculated with the PBG theory.

estimates from the measured spectra are shown in Figure 39 along with some of the excitation energies in the p-p system. The solid and dashed curves represent the predictions of the Watson theory for scattering lengths $a_p = -7.75$ fm. and $a_p = -6.00$ fm. respectively. Here it is not as simple to choose between the two scattering lengths but it does appear that the smaller value gives the closest fit to the measured spectrum.

It is more difficult to determine the contribution from the mass 5 systems to the coincidence proton spectra from the reaction $T(\text{He}^3, np)\text{He}^4$ as both the Li^5 and He^5 states are present. Two quantities are required for this calculation, one being the ratio a^+/a^- for the He^5 system and the other the ratio of the cross section for the breakup through Li^5 to the cross section for the breakup through He^5 . The same ratio $a^+/a^- = 0.33$ found for the states of Li^5 is assumed to apply for He^5 as well. The cross section for the breakup through Li^5 is taken to be twice as large as the cross section for the breakup through He^5 . This estimate is obtained from the results of Barry (13).

Figure 40 shows the coincidence proton spectrum obtained for $\Theta_1 = 61.4^\circ$ and $\Theta_2 = 90^\circ$ and the contribution from the Li^5 and He^5 states determined using the PBG theory and these ratios. Again the available phase-space has been removed from both the measured and calculated spectra. The same calculation was made for the other measured spectrum at $\Theta_1 = 75^\circ$ and $\Theta_2 = 90^\circ$. The results of the background subtraction are shown in Figure 41 along with some of the excitation energies in the n-p system. The

$\text{He}^3(\text{He}^3, 2p)\text{He}^4$

$T_0 = 1.50 \text{ MeV}$

(a) $\theta_1 = 62^\circ$ $\theta_2 = 90^\circ$

— WATSON $a_p = -7.75 \text{ fm}$
 --- WATSON $a_p = -6.00 \text{ fm}$

(b) $\theta_1 = 75^\circ$ $\theta_2 = 90^\circ$

— WATSON $a_p = -7.75 \text{ fm}$
 --- WATSON $a_p = -6.00 \text{ fm}$

Figure 39. Comparison of experimental coincidence proton spectra with Watson theory for the p-p final state interaction.

Figure 40. Coincidence proton spectrum from the reaction $\text{He}^3(\text{He}^3, 2p)\text{He}^4$ for $\Delta_{12} = 151.4^\circ$ divided by the available phase-space. Also shown is the contribution from the states of Li^5 and He^5 calculated with the PBG theory.

$T(\text{He}^3, np) \text{He}^4$

$T_0 = 1.50 \text{ MeV}$

(a) $\theta_1 = 61.4^\circ$ $\theta_2 = 90^\circ$

(b) $\theta_1 = 75^\circ$ $\theta_2 = 90^\circ$

Figure 41. Comparison of experimental coincidence proton spectra with Watson theory for the $n-p$ final state interaction.

solid and dashed curves is spectrum (a) represent the predictions of the Watson theory for n-p singlet scattering lengths $a_n = -23.71$ fm. and $a_n = -30.0$ fm. Since the n-p excitation function calculated using the PBG theory increases rapidly as the excitation energy decreases the coincidence spectrum predicted by this theory would be almost a delta function at $E_{13} = 0$.

As the two curves predicted by the Watson theory for the two values of the n-p scattering length are quite similar it is apparent that the choice of the scattering length which best fits the measured spectrum will depend very critically on the amount of background contribution subtracted. In this analysis the evaluation of the contribution from the He^5 and Li^5 states has been done somewhat arbitrarily due to the lack of information about these states. A more accurate analysis would require considerable experimental effort to learn more about these states, and a more precise theory of final state interactions. The effect of the finite width of the solid state detectors, which has not been considered here, would have to be included in a more careful analysis. A discussion of some of the results of this preliminary analysis and of the improvements which can be made to the experimental technique is contained in the next chapter.

CHAPTER VI

DISCUSSION

A. Present results

One important result of this work is that it gives clear evidence that the singlet states of the p-p and n-p systems may be observed as final state interactions in the reactions $\text{He}^3(\text{He}^3, 2p)\text{He}^4$ and $\text{T}(\text{He}^3, np)\text{He}^4$. Although the singlet deuteron has been observed previously by Parker (2), Simpson (34) and more recently by Cohen et al (35), no such good evidence exists for the singlet two proton state. A combination of two factors, one purely kinematical and the other the Coulomb force accounts for this difference. Kinematics play an important role in three particle breakups in which two of the particles emerge with low relative momentum. This type of breakup is greatly enhanced by a factor in which the dominant term is inversely proportional to the relative momentum of the two particles. However only systems in which the two particles can exist together in a state of low relative momentum, or low internal energy, may take advantage of this kinematical enhancement. The Coulomb force prevents this in the case of the two proton system and for this reason the singlet deuteron is more prominent in these reactions.

The main difference between the measurements reported here and previous measurements is that a lower incident energy has been used. Although the reactions are found to proceed largely via sequential decays through the states of Li^5 or He^5 , corresponding to a nucleon-alpha particle final state inter-

action, it is possible to separate out the contribution from the singlet state of the two nucleon system in each reaction. This contribution, in the case of the p-p system, is found to decrease relative to the contribution from the Li^5 states as the incident energy increases, indicating that low incident energies allow the best information to be gained about the nucleon-nucleon interactions.

Several conclusions may be drawn from a comparison of the predictions of the Watson and PBG theories of final state interactions with the experimental results. The first is that the Watson theory describes the nucleon-nucleon interaction much better than the PBG theory. However it is apparent that neither of these formalisms can account for all the observed features of the experimental spectra. The basic assumption of both theories, that the reaction process can be separated into a primary mechanism which has no effect on the energy and angular distributions of the final particles, and a final state interaction, is obviously not valid for short-lived final state interactions. This is shown to be the case for the states of Li^5 as it is necessary to assume that the Li^5 system retains some memory of the primary mechanism which resulted in its formation, to explain the observed angular dependence of the experimental spectra.

How best to include these effects in a more accurate theory is a difficult question. One method might be to extend the direct reaction theories to predict angular correlations of the type (a, bc) where a,b,c are all particles in contrast to the $(d, n\gamma)$ type of reaction. Another possible approach is to

treat the three body effects exactly as Amado (20) has done, for the case of a final state consisting of two nucleons and an alpha particle rather than three nucleons.

B. Future work

The most useful improvement to the experimental technique used in the measurements reported here would be in particle identification. The use of thin foils for this purpose, although extremely simple and effective in the separation of protons and alpha particles, produces sufficient loss of energy resolution to make this method unsuitable for precise measurements. Two solutions are possible: one method is to place a totally depleted detector in front of one or both of the present detectors and to differentiate between the particles with dE/dx vs. E analysis; the other is to use pulse shape discrimination, relying on the response of the detector-preamplifier arrangement to be sufficiently different for the different particles to allow identification of the particles. This latter method is possible for the separation of protons and alpha particles but it is difficult to separate protons from deuterons in this way. The first method is preferable unless it is necessary to identify low energy particles, especially alpha particles, as these are completely stopped in the totally depleted detector. However for the study of the reactions of interest here, particle identification in only one of the detectors is necessary and this detector can be the proton detector. Particle identification using this technique is useful not only in separating out the contour of interest and

providing better energy resolution but also in removing the two particle final state coincidences which are a problem in these reactions. The contaminant reaction $\text{He}^3(\text{d},\text{p})\text{He}^4$ is characterized by very high proton energies and these protons can be easily removed by discriminating against very low energy pulses from the dE/dx detector. Similarly coincidence events from the reaction mode $\text{T}(\text{He}^3,\text{d})\text{He}^4$ can be removed by eliminating the deuterons with dE/dx vs. E analysis.

One of the problems found in determining the singlet deuteron contribution to the coincidence proton spectra from the reaction $\text{T}(\text{He}^3,\text{np})\text{He}^4$ is the lack of knowledge about the He^5 states and about the relative intensities of the competing reaction modes at the incident energy used. A systematic study of the coincidence spectra from this reaction, similar to that done for the reaction $\text{He}^3(\text{He}^3,2\text{p})\text{He}^4$, would be very useful in supplying this information. This information together with a more accurate theory of the breakup would improve the accuracy with which the parameters determining the n-p interaction can be obtained.

A further refinement to the experimental technique would be the detection of the neutrons from the reaction $\text{T}(\text{He}^3,\text{np})\text{He}^4$. Since one of the desired results of the study of the nucleon-nucleon interaction by means of two dimensional energy analysis of a three particle breakup is to determine the parameters describing the n-n interaction, it is apparent that some type of neutron detection is required to achieve this result. The reaction $\text{T}(\text{T},2\text{n})\text{He}^4$ which is the analog of the two reactions studied here, would be the reaction of interest in determining

these parameters. The neutron energy may be measured by the time delay between the arrival of the charged particle in a solid state detector placed close to the target and the arrival of the neutron in a neutron detector, such as an organic scintillator, placed further away. The resolution of the neutron energy measurements depends on the difference in the two flight paths. The results from the reaction $\text{He}^3(\text{He}^3, 2\text{p})\text{He}^4$ indicate that the best information about the p-p interaction is gained by studying the p-p coincidences with the two detectors at as small an angular separation as possible. This same experiment is possible in the reaction $\text{T}(\text{He}^3, \text{np})\text{He}^4$ if the neutron energy can be measured. There is no problem in this case in obtaining a small angular separation as the neutron detector can be placed directly behind the proton detector. It would be a considerably more difficult experiment to perform two dimensional energy analysis on n-n coincidences from the reaction $\text{T}(\text{T}, 2\text{n})\text{He}^4$ as two neutron time of flight systems would be required. The start pulse for the neutron time of flight could be obtained by detecting the alpha particle in a solid state detector. A simpler experiment would be the study of n- He^4 coincidences from this reaction. Although it is clear that more precise theories are required to obtain parameters such as scattering lengths from these reactions, a direct comparison of the results from the reactions $\text{T}(\text{He}^3, \text{np})\text{He}^4$ and $\text{T}(\text{T}, 2\text{n})\text{He}^4$ should give considerable information about the n-n interaction.

APPENDIX A

REACTION KINEMATICS FOR THREE FINAL PARTICLES

The symbols for the various breakup parameters are listed in Table 1 of Chapter II. For the sake of completeness some of the relations determining the breakup in the laboratory system which were derived in Chapter II will be given here as well. Using the requirement of conservation of momentum and energy

$$\vec{P}_0 = \vec{P}_1 + \vec{P}_2 + \vec{P}_3 \quad \text{A-1}$$

and $T_0 + Q = T_1 + T_2 + T_3 \quad \text{A-2}$

where $P_i^2 = 2m_i T_i$

and solving both equations for P_3 yields the quadratic equation

$$AP_2^2 - 2BP_2 + C = 0 \quad \text{A-3}$$

where $A = 1 + m_3/m_2$

$$B = P_0 \cos \Theta_2 - P_1 \cos \Delta_{12}$$

$$C = (1 + m_3/m_1) P_1^2 + (1 - m_3/m_0) P_0^2 - 2P_0 P_1 \cos \Theta_1 - 2m_3 Q$$

The solutions of A-3 are

$$P_2^\pm = (B \pm \sqrt{B^2 - AC})/A \quad \text{A-4}$$

subject to the requirement that

$$B^2 - AC \geq 0$$

By inserting the solution for $P_2(P_1)$ into equation A-3 and performing simple algebraic manipulation the quadratic equation for P_1 as a function of the angles $\Theta_1, \Theta_2, \Phi_1, \Phi_2$ is obtained.

$$DP_1^2 - 2EP_1 + F = 0 \quad \text{A-5}$$

where

$$D = (1 + m_3/m_1) A - \cos^2 \Delta_{12}$$

$$E = P_0 A \cos \Theta_1 - P_0 \cos \Theta_2 \cos \Delta_{12}$$

$$F = (1 - m_3/m_0) AP_0^2 - P_0^2 \cos^2 \Theta_2 - 2m_3 A Q$$

From these relations all the necessary parameters in the laboratory system may be evaluated.

The parameters in the scm. system, the centre of mass system of the three final state particles, are determined using the velocity triangle shown in Figure A1. The velocity of the centre of mass V_{scm} is given by

$$V_{scm} = V_0 m_0 / M \quad \text{A-6}$$

The law of cosines is used to determine the scm. velocities.

$$v_i = (V_i^2 + V_{scm}^2 - 2V_i V_{scm} \cos \Theta_i)^{1/2} \quad \text{A-7}$$

The azimuthal angles do not change from the lab. to the scm. system as polar coordinates are used with the z-axis along the beam direction.

$$\varphi_i = \Phi_i$$

The polar angles in the scm. system are obtained from Figure A1

$$\cos \theta_i = (V_i \cos \Theta_i - V_{scm}) / v_i \quad A-8$$

Using these relations the scm. kinetic energies and momenta may be calculated for the three particles. The parameters of the third particle may also be determined from those of particle 1 and 2 by utilizing momentum and energy conservation in the scm. system.

Thus

$$\sum_{i=1}^3 \vec{p}_i = 0$$

and

$$\sum_{i=1}^3 t_i = E$$

yields

$$p_3 = \sqrt{2m_3(E - t_1 - t_2)} \quad A-9$$

and

$$\cos \theta_3 = -(p_1 \cos \theta_1 + p_2 \cos \theta_2) / p_3 \quad A-10$$

The coordinate system of importance as far as the theoretical analysis of the breakup is concerned is the rcm. system, the centre of mass system of the two particle clusters. The notation used here is to specify the cluster (j+k) by the superscript (i) where i,j,k is some permutation of 1,2,3. Thus $v_{rcm}^{(i)}$ is the recoil velocity of the cluster (j+k) and is given by

$$\vec{v}_{rcm}^{(i)} = - \frac{m_i}{m_j + m_k} \vec{v}_i \quad A-11$$

The relationship between the scm. system and the rcm(i) system is shown in Figure A2. The velocity of one of the particles from the breakup of the cluster (j+k) is

$$v_j^{(i)} = (v_{rcm}^{(i)2} + v_j^2 + 2v_{rcm}^{(i)} v_j \cos \delta_{ij})^{1/2} \quad A-12$$

with the internal energy of the cluster given by

$$E_{jk} = \frac{m_j(m_j+m_k)}{2m_k} v_j^{(i)2} \quad A-13$$

The polar and azimuthal angles $\theta_j^{(i)}$, $\varphi_j^{(i)}$ which measure the orientation of $v_j^{(i)}$ with respect to the beam axis may be calculated from Figure A2.

$$\cos \theta_j^{(i)} = \frac{1}{v_j^{(i)}} (v_j \cos \theta_j + v_{rcm}^{(i)} \cos \theta_i) \quad A-14$$

$$\cos(\varphi_j^{(i)} - \varphi_i^{(i)}) = \frac{1}{v_j^{(i)} \sin \theta_j^{(i)}} (v_j \sin \theta_j \cos(\varphi_j - \varphi_i) + v_{rcm}^{(i)} \sin \theta_i) \quad A-15$$

The other angle of interest is the polar angle $\eta_j^{(i)}$ which measures the orientation of $v_j^{(i)}$, the velocity of one of the particles from the breakup of the cluster (j+k), with respect to the velocity of the cluster $v_{rcm}^{(i)}$.

$$\cos \eta_j^{(i)} = - (v_{rcm}^{(i)} + v_j \cos \delta_{ij}) / v_j^{(i)} \quad A-16$$

Jacobians

In general theoretical expressions for quantities such as energy distributions or cross sections are most easily obtained in a reference frame other than the laboratory system. This necessitates the determination of the conversion factor which transforms quantities in the reference frame used in the theoretical analysis to the laboratory system. In the three particle breakup via the sequential decay mode the important theoretical distribution is the triple correlation which is obtained with one particle in the scm. system and the other two in rcm. coordinates.

This quantity called $\sigma_{rcm}^{(i)}$ for particle i emitted first is given by

$$\sigma_{rcm}^{(i)} = \frac{d^3\sigma}{dE_{jk} d\Omega_i(scm) d\Omega_j(rcm)} \quad A-17$$

The corresponding quantities in the scm. and lab. systems are

$$\sigma_{scm} = \frac{d^3\sigma}{dt_1 d\Omega_1(scm) d\Omega_2(scm)} \quad A-18$$

remembering that t_1 is the K.E. in the scm.system of particle 1 and

$$\sigma_{lab} = \frac{d^3\sigma}{dT_1 d\Omega_1(lab) d\Omega_2(lab)} \quad A-19$$

These quantities are related by the following expressions

$$\sigma_{scm} = J_i \sigma_{rcm}^{(i)}$$

$$\sigma_{lab} = J_L \sigma_{scm}$$

where J_i and J_L , the Jacobians connecting the appropriate coordinate systems are defined as

$$J_i = \frac{\partial(E_{jk}; \cos\theta_i, \phi_i; \cos\theta_j^{(i)}, \phi_j^{(i)})}{\partial(t_1; \cos\theta_1, \phi_1; \cos\theta_2, \phi_2)} \quad A-20$$

$$J_L = \frac{\partial(t_1; \cos\theta_1, \phi_1; \cos\theta_2, \phi_2)}{\partial(T_1; \cos\theta_1, \phi_1; \cos\theta_2, \phi_2)} \quad A-21$$

$$\text{and } J_{iL} = \frac{\partial(E_{jk}; \cos\theta_i, \phi_i; \cos\theta_j^{(i)}, \phi_j^{(i)})}{\partial(T_1; \cos\theta_1, \phi_1; \cos\theta_2, \phi_2)} \quad A-22$$

The determination of these Jacobians involves the evaluation of the appropriate partial derivatives using the relations between the coordinate systems given in the preceeding

section. These calculations are given in Bronson's thesis and yield the following results

$$J_i = \frac{M P_1 P_2^2}{m_j P_i P_j^{(i)}} \frac{1}{(A P_2 + P_1 \cos \delta_{ij})} \quad A-23$$

$$J_L = \frac{P_1 P_2^2}{P_1 P_2^2} \frac{(A P_2 + P_1 \cos \delta_{ij})}{(A P_2 + P_1 \cos \Delta_{ij} - P_0 \cos \Theta_2)} \quad A-24$$

$$J_{LL} = \frac{M P_1 P_2^2}{m_j P_i P_j^{(i)}} \frac{1}{(A P_2 + P_1 \cos \Delta_{ij} - P_0 \cos \Theta_2)} \quad A-25$$

The derivation of these quantities from the partial derivatives, although straightforward, involves the manipulation of many long and complicated expressions. A simpler method of determining the Jacobians involves the evaluation of the triple correlation functions for the case of a statistical breakup in the three coordinate systems. This requires a knowledge of phase space calculations.

Phase space

The transition probability for a reaction has the form

$$\omega = \frac{2\pi}{\hbar} |H|^2 \rho(E)$$

where H is the matrix element connecting the initial and final states and $\rho(E)$ is the density of final states or phase space factor. For statistical breakups the matrix element is assumed to be constant so that the breakup probability depends only on the available phase space. The non-relativistic phase space

factor for a n-particle final state is defined as (36)

$$\rho_n(E) = \int \prod_{i=1}^n d^3 p_i \delta^3(\sum_{i=1}^n \vec{p}_i - \vec{P}) \delta(\sum_{i=1}^n \frac{p_i^2}{2m_i} - E) \quad A-26$$

where E is the total energy available to the system,

$\vec{P}$ is the total momentum of the system

$d^3 p_i$ is the momentum vector differential and can be

written as $p_i^2 dp_i d\Omega_i$ in polar coordinates.

The δ functions are used to include the requirements of momentum and energy conservation. The phase space distributions are determined by restricting the integration of equation A-26 to the appropriate variables.

To illustrate an explicit calculation of $\rho_n(E)$ consider the case of a final state consisting of two particles with momenta $\vec{p}_1$ and $\vec{p}_2$ in the center of mass. Since $\vec{p}_1 = -\vec{p}_2$ integration over p_2 gives

$$\rho_2(E) = \int p_1^2 dp_1 \int d\Omega_1 \delta(\frac{p_1^2}{2m_1} + \frac{p_1^2}{2m_2} - E)$$

If the angular distribution of particle 1 is desired the integration over the solid angle Ω_1 is omitted.

$$\frac{d\rho_2(E)}{d\Omega_1} = \int p_1^2 dp_1 \delta(p_1^2 \frac{(m_1+m_2)}{m_1 m_2} - E)$$

The rule for integration over a δ function is the following

$$\int g(x) \delta(f(x)) dx = \frac{g(x_0)}{|f'(x_0)|}$$

where

$$f(x_0) = 0$$

Integrating over p_1 using this rule gives

$$\frac{d\rho_2(E)}{d\Omega_1} = \frac{m_1 m_2}{m_1 + m_2} P_1 \quad \text{A-27}$$

with

$$P_1^2 = \frac{2 m_1 m_2}{m_1 + m_2} E$$

These results show that the statistical probability of a two particle breakup is proportional to the particle momentum and is isotropic in the center of mass system.

By extending these calculations to the case of a three particle final state it is possible to evaluate the triple correlation functions σ_{lab} , σ_{scm} , and $\sigma_{rcm}^{(i)}$ defined by equations A-17, A-18 and A-19 for a purely statistical breakup.

In the lab. system the momentum equation is

$$\vec{P}_1 + \vec{P}_2 + \vec{P}_3 = -\vec{P}_0$$

and equation A-26 for $n = 3$ is given by

$$\rho_3(E) = \int P_1^2 dP_1 \int d\Omega_1 \int P_2^2 dP_2 \int d\Omega_2 \delta\left(\underbrace{\frac{P_1^2}{2m_1} + \frac{P_2^2}{2m_2} + \frac{P_3^2(P_1, P_2)}{2m_3}}_f - E\right)$$

For constant P_1

$$\frac{dE}{dP_2} = \frac{1}{m_3} (AP_2 + P_1 \cos \Delta_{12} - P_0 \cos \Theta_2)$$

where the symbols have been defined previously. The integration of $\rho_3(E)$ over P_2 yields

$$\frac{d^3 \rho_3(E)}{dP_1 d\Omega_1 d\Omega_2} = \frac{m_3 P_1^2 P_2^2}{(AP_2 + P_1 \cos \Delta_{12} - P_0 \cos \Theta_2)}$$

$$\text{and } \sigma_{lab} \propto \frac{d^3 \rho_3(E)}{dP_1 d\Omega_1 d\Omega_2} = \frac{m_1 m_3 P_1 P_2^2}{(AP_2 + P_1 \cos \Delta_{12} - P_0 \cos \Theta_2)} \quad \text{A-28}$$

The same quantity in the scm. system is obtained by setting $P_0 = 0$.

$$\sigma_{scm} \propto \frac{d^3 p_3(E)}{dt, d\Omega_1(scm) d\Omega_2(scm)} = \frac{m_1 m_3 P_1 P_2^2}{A P_2 + P_1 \cos \delta_{12}} \quad A-29$$

so that

$$J_L = \frac{\sigma_{lab}}{\sigma_{scm}} = \frac{P_1 P_2^2 (A P_2 + P_1 \cos \delta_{12})}{P_1 P_2^2 (A P_2 + P_1 \cos \Delta_{12} - P_0 \cos \Theta_2)} \quad A-30$$

which is the same as equation A-24.

The triple correlation function rcm(i) system is obtained by considering the breakup as a two particle decay in the scm. system followed by a two particle decay in the rcm(i) system. As shown previously in equation A-27 these breakups are proportional to the particle momenta. If i is the first emitted particle in the breakup then

$$\frac{d p_2(E)}{d \Omega_i(scm)} = \frac{m_i (m_j + m_k)}{M} P_i$$

and similarly for the breakup of the cluster (j+k)

$$\frac{d p_2(E)}{d \Omega_j(rcm)} = \frac{m_j m_k}{m_j + m_k} P_j^{(i)}$$

For a statistical breakup all internal energies are equally probable so that

$$\sigma_{rcm}^{(i)} \propto \frac{d p_2(E) d p_2(E)}{d \Omega_i(scm) d \Omega_j(rcm)} = \frac{m_i m_j m_k}{M} P_i P_j^{(i)} \quad A-31$$

$$J_{iL} = \frac{M}{m_2 P_i P_j^{(i)}} \frac{P_1 P_2^2}{(A P_2 + P_1 \cos \Delta_{12} - P_0 \cos \Theta_2)} \quad A-32$$

which agrees with equation A-25. A listing of the computer program written to calculate the breakup parameters in all three systems along with the Jacobians is given in Appendix B.

APPENDIX B

COMPUTER PROGRAM LISTINGS

In this appendix two FORTRAN IV computer programs useful in calculating the quantities of interest in a three particle breakup, are listed. The FORTRAN names for the various breakup parameters given in Appendix A have been chosen, where possible, to be self-explanatory, i.e. VSCM corresponds to v_{scm} , V12 to $v_2^{(1)}$ etc. All input and output energies are in MeV, masses are in a.m.u. and angles are in degrees.

The program CONTOUR evaluates the kinematic contours $T_2(T_1)$, where T_1 and T_2 are the laboratory energies of two of the three final particles. In a two dimensional energy analysis these would be the two particles detected in coincidence. The input quantities are the following: Q , m_0 , m_1 , m_2 , m_3 , T_0 , Θ_1 , Θ_2 , Φ_1 , Φ_2 , and SF where all the quantities except SF are described in Table 1. SF is the energy increment used in determining the function $T_2(T_1)$. The form of the output is a contour plot of T_1 vs. T_2 using the Calcomp plotter, one of the output facilities of the U.B.C. Computing Center.

The program KINJAC evaluates all the breakup parameters in the three coordinate systems, lab., scm., and rcm. and the Jacobians for the transformation from rcm. coordinates to lab. coordinates. The input quantities are the same as for CONTOUR except that SF is replaced by three quantities TI, TF and DT. The calculations are carried out as a function of the laboratory energy of one of the final particles with TI the initial value

of this energy, TF the final value and DT the energy increment. The output quantities are the three corresponding excitation energies E_{ij} of the two particle systems, the three Jacobians J_{iL} and the breakup angles $\eta_j^{(i)}$. Another quantity obtained in this calculation is PSF, the phase space factor, which allows the statistical energy distribution to be determined.

SIBFIC CONTOUR

CONTOUR-1

```
C      KINEMATIC CONTOURS T2(T1) FOR A THREE PARTICLE BREAKUP
      DIMENSION T1(600), T2(600)
      CALL PLOTS
      CALL AXIS(2.0,2.0,7HT1(MEV),-7,8.0,0.0,0.0,2.0)
      CALL AXIS(2.0,2.0,7HT2(MEV),7,6.0,90.0,0.0,2.0)
50     READ(5,1) Q,AM0,AM1,AM2,AM3
      READ (5,1) T0,TH1,TH2,PH1,PH2,SF
      IF (SF) 120,45,45
45     SW1 = 1.0
      SW = 0.0
      T1SW = 0.0
      A = (AM2+AM3)/AM2
      P0 = SQRT(2.0*AM0*T0)
      CTH1 = COS(.0174532*TH1)
      CTH2 = COS(.0174532*TH2)
      STH1 = SIN(.0174532*TH1)
      STH2 = SIN(.0174532*TH2)
      CPH = COS(.0174532*(PH2-PH1))
      CD12 = CTH1*CTH2 + STH1*STH2*CPH
      D12 = ARCOS(CD12)/.01745
      D = (AM1+AM3)/AM1
      G = (AM0-AM3)/AM0
      E = P0*CTH1
      F = G*P0**2 - 2.0*AM3*Q
      P1INT = (E + SQRT(E**2-D*F))/D
      T1INT = P1INT**2/2.0/AM1
      I = 1
      T1(I) = 0.0
40     P1 = SQRT(2.0*AM1*T1(I))
      B = P0*CTH2 - P1*CD12
      C = D*P1**2 + G*P0**2 - 2.0*P0*P1*CTH1 - 2.0*AM3*Q
      DIFF = B**2 - A*C
      IF (DIFF) 20,35,35
35     IF (SW1) 28,30,30
30     P2 = (B + SQRT(DIFF))/A
29     T2(I) = P2**2/2.0/AM2
      GO TO 15
28     P2 = (B - SQRT(DIFF))/A
      GO TO 29
20     IF (SW1) 100,100,18
18     SW1 = -1.0
```

CONTOUR-2

```

T1SW = T1(I-1)
T2SW = T2(I-1)
GO TO 14
15  I = I+1
14  T1(I) = T1(I-1) + SW1*SF
    IF (T1(I).GT.T1INT.AND.D12.LE.90.1) GO TO 100
    IF (T1(I).LE.T1INT.AND.SW1.EQ.1.0) GO TO 40
    IF (T1(I).LE.T1INT.AND.SW1.EQ.(-1.0)) GO TO 100
    IF (T1(I).GT.T1INT.AND.SW.EQ.0.0) GO TO 12
    GO TO 40
12  SF = 0.1*SF
    SW = 1.0
    GO TO 40
100 KMAX = I-1
    WRITE (6,2) Q,AM0,AM1,AM2,AM3
    WRITE (6,3) TO,TH1,TH2,PH1,PH2
    DO 110 I=1,KMAX
    110 T1(I) = T1(I)/2.0 + 2.0
        T2(I) = T2(I)/2.0 + 2.0
        IF (AM1-AM2) 112,118,112
    112 IF (T1SW) 118,118,115
    115 T1SW = T1SW/2.0 + 2.0
        T2SW = T2SW/2.0 + 2.0
        GO TO 119
118  KMID = KMAX/2
        T1SW = T1(KMID)
        T2SW = T2(KMID)
119  CALL LINE (T1,T2,KMAX,1)
        GO TO 50
120  CALL PLOTND
        STOP
1  FORMAT (6F10.5)
2  FORMAT (20H Q-VALUE $ MASSES = , 5F8.3,/)
3  FORMAT (24H INC. ENERGY $ ANGLES = , 5F8.2,/)
5  FORMAT (20H      T1      T2 ,/)
6  FORMAT (1X, 2F10.3)
    END

```

\$ENTRY

SIBFTC KINJAC

KINJAC-1

```
C      THREE PARTICLE BREAKUP KINEMATICS AND JACOBIANS
5      READ (5,99) AM0,AM1,AM2,AM3,Q
      WRITE (6,100) AM0,AM1,AM2,AM3,Q
      READ (5,99) T0,SL1,SL2,CPH2
      WRITE (6,101) T0,SL1,SL2,CPH2
      A = (AM3+AM2)/AM2
      AM = AM1+AM2+AM3
      T = 2.0
      RAD = 0.0174532
      UL1 = COS(RAD*SL1)
      SL1 = SIN(RAD*SL1)
      UL2 = COS(RAD*SL2)
      SL2 = SIN(RAD*SL2)
      CPH2 = COS(CPH2*RAD)
      UL12 = UL1*UL2 + SL1*SL2*CPH2
10     READ (5,99) T1,TF,DT
      T1 = T1
      WRITE (6,102)
      WRITE (6,103)
C      LAB VALUES
15     P0 = SQRT(T*AM0*T0)
      P1 = SQRT(T*AM1*T1)
      C = (AM1+AM3)/AM1*P1**2 + (AM0-AM3)/AM0*P0**2 - T*AM3*Q
      B = P0*UL2 - P1*UL12
      IF (B) 25,20,20
20     DIFF = (B**2 - A*C)
      IF (DIFF) 5,22,22
22     P2 = (B + SQRT(DIFF))/A
      GO TO 30
25     B = -B
      DIFF = (B**2 - A*C)
      IF (DIFF) 5,27,27
27     P2 = (-B + SQRT(DIFF))/A
      IF (P2) 5,30,30
30     T2 = P2**2/T/AM2
      T3 = Q+T0-T1-T2
      P3 = SQRT(T*AM3*T3)
      UL3 = (P0 - P1*UL1 - P2*UL2)/P3
      IF (UL3) 35,40,40
35     SL3 = -UL3
```

```

GO TO 45
40 SL3 = UL3
45 SL3 = SQRT(1.0 - SL3**2)
CPH3 = -(P1*SL1 + P2*SL2*CPH2)/P3/SL3
V1 = P1/AM1
V2 = P2/AM2
V3 = P3/AM3
C SCM VALUES
VSCM = P0/AM
VSQ = VSCM**2
SV1 = SQRT(V1**2 + VSQ - T*V1*VSCM*UL1)
SV2 = SQRT(V2**2 + VSQ - T*V2*VSCM*UL2)
SV3 = SQRT(V3**2 + VSQ - T*V3*VSCM*UL3)
U1 = (V1*UL1 - VSCM)/SV1
U2 = (V2*UL2 - VSCM)/SV2
U3 = (V3*UL3 - VSCM)/SV3
U12 = U1*U2 + V1/SV1*V2/SV2*SL1*SL2*CPH2
U31 = U3*U1 + V3/SV3*V1/SV1*SL3*SL1*CPH3
C RCM VALUES
RV1 = AM1*SV1/(AM2+AM3)
RV2 = AM2*SV2/(AM1+AM3)
RV3 = AM3*SV3/(AM1+AM2)
V21 = SV2**2 + RV1**2 + T*SV2*RV1*U12
V12 = SV1**2 + RV2**2 + T*SV1*RV2*U12
V13 = SV1**2 + RV3**2 + T*SV1*RV3*U31
E23 = AM2*(AM2+AM3)/T/AM3*V21
E13 = AM1*(AM1+AM3)/T/AM3*V12
E12 = AM1*(AM1+AM2)/T/AM2*V13
V12 = SQRT(V12)
V21 = SQRT(V21)
V13 = SQRT(V13)
CN21 = -(RV1 + SV2*U12)/V21
CN12 = -(RV2 + SV1*U12)/V12
CN13 = -(RV3 + SV1*U31)/V13
C JACOBIANS
PSF = P1*P2**2/(P2*A + P1*UL12 - P0*UL2)
B = AM*PSF/AM1/AM2**2
AJ1 = B/SV1/V21
AJ2 = B/SV2/V12
AJ3 = B/SV3/V13*AM2/AM3
WRITE (6,104) T,E23,E13,E12,CN21,CN12,CN13
WRITE (6,105) AJ1,AJ2,AJ3,PSF

```

KINJAC-3

```

T1 = T1+DT
IF (T1-TF) 15,15,5
99  FORMAT (5F10.5)
100  FORMAT (1H1, 5H M0=,F6.3,4H M1=,F6.3,4H M2=,F6.3,4H M3=,F6.3,
1    3H Q=,F7.3)
101  FORMAT (/ , / , 6H T0=,F6.2,8H THETA1=,F6.2,8H THETA2=,F6.2,
1    7H PHI12=,F6.2)
102  FORMAT (/ , / , 70H      T1      E23      E13      E12      CN21
1    CN12      CN13 )
103  FORMAT (80H      J1
1    J2      J3      PSF )
104  FORMAT (/ , 7F10.3)
105  FORMAT (40H      ,4F10.3)
END
$ENTRY

```


APPENDIX C

SEMI-EMPIRICAL EXPRESSION FOR ENERGY LOSS IN THIN FOILS

As thin foils are used extensively in this experiment, both in the entrance and exit windows of the gas cell and to provide particle discrimination in the two dimensional energy analysis, it is necessary to have some means of evaluating the energy loss of the different particles in traversing the foils. A semi-empirical relation was developed for the specific energy loss for protons by fitting the available data (37) from $Z = 13$ (Al) to $Z = 79$ (Au). The usual expression for the stopping cross section is

$$-\frac{1}{N} \frac{dE}{dx} = \frac{4\pi e^4 z^2 Z}{m_0 v^2} \ln\left(\frac{2m_0 v^2}{\bar{I}} + C(E, Z)\right) \quad C-1$$

where N is the number of atoms of the stopping material per unit volume.

Z is the atomic number of the stopping material.

ze is the charge of the particle moving with velocity v .

m_0 is the electron mass.

$\bar{I}$ is the average ionization potential.

$C(E, Z)$ is a correction term which takes into account the binding energy corrections at low energies and the polarization corrections at high energies.

The functional forms of $C(E, Z)$ and $\bar{I}$ were chosen empirically to give a reasonably good fit to the data. The quantity $\bar{I}/Z$ has been found previously (38) to decrease slightly with increasing Z with the form of the decrease given approximately by

$$\bar{I}/Z = 13.5 - 0.04Z \text{ ev.}$$

C-2

$C(E,Z)$ was chosen to have the following variation with E and Z

$$C(E,Z) = a_1 + a_2/Z + (a_3 + a_4/Z + a_5/Z^2)E + a_6E^2 \quad \text{C-3}$$

There is no theoretical basis for choosing this particular form for the correction term, although it is evident that a simpler form of $C(E,Z)$ cannot give the necessary variation with E and Z .

On inserting the values of the numerical constants into expression C-1 the specific energy loss for protons becomes

$$-\frac{dE}{dx} = \frac{144.12}{A} \frac{Z}{E} \ln\left(\frac{2178.6}{I} \frac{E}{Z} + C(E,Z)\right) \frac{\text{kev-cm}^2}{\text{mg}} \quad \text{C-4}$$

The best least squares fit to the specific energy loss data for protons in Al, Cu and Au from 100 keV to 10 MeV (37) is obtained with the following parameters:

$$\begin{array}{lll} a_1 = 0.976 & a_2 = -2.435 & a_3 = 0.239 \\ a_4 = -23.40 & a_5 = -226.0 & a_6 = -0.057 \end{array}$$

The calculated energy loss with these parameters is within 10% of the tabulated data over the entire energy range with the agreement considerably better for Z about 29 and energies above 1 MeV. The specific energy loss for the other particles involved in the experiment is determined from the following relationships:

$$\frac{dE}{dx}(\text{deuteron})_E = \frac{dE}{dx}(\text{proton})_{E/2}$$

$$\frac{dE}{dx}(\text{triton})_E = \frac{dE}{dx}(\text{proton})_{E/3}$$

$$\frac{dE}{dx}(\text{He}^3 \text{ particle})_E = \bar{Z}^2 \frac{dE}{dx}(\text{proton})_{E/3}$$

$$\frac{dE}{dx}(\text{alpha particle})_E = \bar{Z}^2 \frac{dE}{dx}(\text{proton})_{E/4}$$

where $\bar{Z}^2$ is the equilibrium helium ion charge and is a function of the particle energy. The functional form for $\bar{Z}^2$ was also chosen empirically by fitting the data of Whaling (39) with the following result:

$$E > 2.0 \text{ MeV} \quad \bar{Z}^2 = 4.0$$

$$E < 2.0 \text{ MeV} \quad \bar{Z}^2 = 4.0 + b_1/E + b_2/E^2 + b_3/E^3 + b_4/E^4 \quad \text{C-5}$$

where	$b_1 = 0.757$	$b_2 = -1.83$
	$b_3 = 0.722$	$b_4 = -0.08$

The energy loss in a foil of finite thickness is determined by using an iterative calculation to take account of the change in the specific energy loss as the particle loses energy in the foil. The first approximation to the energy loss uses the specific energy loss evaluated at the entrance energy. Subsequent approximations are obtained by dividing the previously determined energy loss into four intervals and evaluating the mean specific energy loss for the corresponding five particle energies using a Simpson's rule integration. The iteration continues until successive approximations are within 1/2%. A program to do this calculation was written for the PDP-8 computer. Typical results of these calculations for the particles of interest in this experiment are shown in the Figures C1 and C2.

Figure C1. Energy loss curves for protons and deuterons in nickel foils.

Figure C2. Energy loss curves for He³ and He⁴ in nickel foils.

REFERENCES

1. Minutes of the Topical Conference on Correlations of Particles Emitted in Nuclear Reactions, Gatlinburg, Tennessee, 1964, Rev. Mod. Phys. 37 (1965).
2. R.D. Parker, R.F. Donovan, J.V. Kane and J.F. Mollenauer, Phys. Rev. Letters 14, 15 (1965).
3. D.B. Smith, N. Jarmie and A.M. Lockett, Phys. Rev. 129 785 (1963).
4. H.E. Conzett, E. Shield, R.J. Slobodrian and S. Yamabe, Phys. Rev. Letters 13, 625 (1964).
5. T.A. Tombrello and A.D. Bacher, Phys. Letters 17, 37 (1965).
6. N. Jarmie and R.C. Allen, Phys. Rev. 111, 1121 (1958).
7. J.P. Aldridge, B.H. Wildenthal and D.H. Youngblood, Rev. Mod. Phys. 37, 430 (1965).
8. G.C. Phillips, T.A. Griffy and L. C. Biedenharn, Nucl. Phys. 21, 327 (1960).
9. A.D. Bacher and T.A. Tombrello, Rev. Mod. Phys. 37, 433 (1965).
10. R.W. Zurmuhle, Nucl. Phys. 72, 225 (1965).
11. C.D. Moak, Phys. Rev. 92, 383 (1953).
12. B. Kuhn and B. Schlenk, Nucl. Phys. 48, 353 (1963).
13. J.F. Barry, R. Batchelor and B.E.F. Macefield, Proc. Rutherford Jubilee Int. Conf., Manchester, p.543 (1961).
14. E.W. Blackmore, M.Sc. Thesis, University of British Columbia (1965).
15. E.W. Blackmore and J.B. Warren, Phys. Rev. Letters 16, 520 (1966).
16. K.M. Watson, Phys. Rev. 88, 1163 (1952).
17. J.D. Bronson, Ph.D. thesis, William Marsh Rice University (1964).
18. G.G. Ohlsen, Research School of Physical Sciences, Australian National University, Canberra. Publication ANU-P/340 (1965).

References continued

19. M.A. Reimann, Ph.D. thesis, University of British Columbia (1967).
20. P. Swan, Rev. Mod. Phys. 37, 336 (1965).
21. S. Dushman, Scientific Foundations for Vacuum Technique, John Wiley and Sons, p.453 (1962).
22. R.B. Tomlinson and R.L. Brown, IEEE Transactions on Nuclear Science, Vol. NS-11, p. 28 (April, 1964).
23. B.A. Whalen, Ph.D. thesis, University of British Columbia (1965).
24. P.F. Donovan, Rev. Mod. Phys. 37, 501 (1965).
25. R.J.N. Phillips, Nucl. Phys. 53, 650 (1964).
26. F.C. Barker and P.B. Treacy, Nucl. Phys. 38, 33 (1962).
27. I. Duck, Rev. Mod. Phys. 37, 418 (1965).
28. R. Aaron, R.D. Amado and Y.Y. Yam, Phys. Rev. 140, B1291 (1965).
29. R. Aaron and R.D. Amado, Phys. Rev. 150, 859 (1966).
30. A.C.L. Barnard, C.M. Jones and J.L. Weil, Nucl. Phys. 50, 604 (1964).
31. B. Hoop and H.H. Barschall, International Conference on Polarization Phenomena of Nucleons, Karlsruhe, Ger. (1965).
32. M.A. Preston, Physics of the Nucleus, Addison-Wesley Publishing Co. (1962).
33. N. Austern, Fast Neutron Physics, John Wiley and Sons, p. 1193 (1963).
34. W.D. Simpson, J.D. Bronson, W.R. Jackson and G.C. Phillips, Rev. Mod. Phys. 37, 523 (1965).
35. B.L. Cohen, E.C. May and T.M. O'Keefe, Phys. Rev. Letters 18, 962 (1967).
36. O. Skjeggsted, Proceedings of the 1964 Easter Summer School for Physicists, CERN 64-13, Vol. II (1964).

References continued

37. J.B. Marion, Nuclear Data Tables, Part 3 (1960) National Academy of Sciences-National Research Council, Washington, D.C.
38. J. Lindhard and M. Scharff, p. 49, Penetration of Charged Particles in Matter, Nuclear Science Series Report No. 29 (1960).
39. W. Whaling, Handbuch der Physik, E. Flugge, ed., Springer-Verlag, Berlin, Vol. 34 (1958).