DEOXYRIBONUCLEASE

IN THE INTESTINAL MUCOSA OF RAT

by

C. Y. LEE

B. Sc., University of British Columbia, 1963.

A THESIS SUBMITTED IN PARTIAL FULFILMENT OF

THE REQUIREMENTS FOR THE DEGREE OF

MASTER OF SCIENCE

in the Department

of

Biochemistry.

We accept this thesis as conforming to the
required standard for the degree of
MASTER OF SCIENCE.

The University of British Columbia
April, 1965.

In presenting this thesis in partial fulfilment of the requirements for an advanced degree at the University of British Columbia, I agree that the Library shall make it freely available for reference and study. I further agree that permission for extensive copying of this thesis for scholarly purposes may be granted by the Head of my Department or by his representatives. It is understood that copying or publication of this thesis for financial gain shall not be allowed without my written permission.

Department of B iochemistry

The University of British Columbia,
Vancouver 8, Canada

Date 5th May, 1965.

ABSTRACT

The properties of the deoxyribonuclease activity in the intestinal mucosa of rat have been studied. It was found that the activity was extractable from a whole cell preparation of the tissue with either Krebs Ringer phosphate buffer (pH 7.8) or physiological saline. The former was a better extracting medium, the buffered extract being more stable to storage at -20°C .

Two active proteins were precipitated on fractionation of the Krebs Ringer phosphate buffer extract with ammonium sulfate. One fraction precipitated at 20% saturation of $(\text{NH}_4)_2\text{SO}_4$ was identified as DNase I by its optimum pH, ionic requirements and by its reaction to known DNase I inhibitors such as EDTA, citrate and arsenate. Ion-exchange chromatography on DEAE-cellulose (chloride form) established that the products of reaction ranged from mononucleotides through to oligonucleotides with a degree of polymerization larger than 7. These products were shown to carry a phosphomonoester linkage on the 5'-carbon. This fraction was designated as the 20%P enzyme and represented a 20 fold purification of the crude deoxyribonuclease extract.

The supernatant fraction obtained after the 20%P enzyme was removed by centrifugation was found to still contain

considerable DNase activity. This residual activity disappeared from the solution when the ammonium sulfate concentration was increased to 30% saturation, but all attempts to recover the DNase activity from the protein precipitated at this salt concentration were unsuccessful. Consequently, studies of this residual DNase activity were carried out using the crude extract.

This second deoxyribonuclease was shown to be qualitatively different from the 20% enzyme. It did not require activation by Mg^{++} , and was not inhibited by EDTA, citrate or arsenate. Some of the products formed by the crude enzyme extract were shown to terminate in 3'-phosphoryl groups. These products were not formed by the 20%P enzyme and therefore must be due to the residual deoxyribonuclease. In all likelihood then, this second deoxyribonuclease activity was of the DNase II type.

The intracellular distribution of the intestinal mucosal deoxyribonuclease was studied by differential centrifugation of the tissue homogenate. DNase I activity was found to occur in the mitochondria whereas DNase II was associated with some lighter subcellular particles.

ACKNOWLEDGMENTS

The author wishes to express his sincere thanks and deep appreciation to Dr. S.H. Zbarsky for his guidance and encouragement during the course of this research.

The support of the National Research Council in the form of a studentship is also gratefully acknowledged.

TABLE OF CONTENTS

	<u>Page</u>
<u>INTRODUCTION</u>	1
<u>EXPERIMENTAL</u>	
I. Preparation of Deoxyribonucleic Acids . . .	16
(i) Preparation of DNA from small intestinal mucosa of rat	16
(ii) Deproteinization of salmon sperm DNA	18
(iii) Estimation of nitrogen content of deoxyribonucleic acids	19
(iv) Determination of phosphorus content of DNA	19
(v) Determination of $\zeta(P)$ values of DNA preparations	20
II. Preparation and Purification of Deoxyribonuclease from Rat Intestinal Mucosa	21
(i) Measurement of enzyme activity . . .	21
(ii) Preparation of crude enzyme extract	27
(iii) Purification of intestinal mucosal deoxyribonuclease by ammonium sulfate fractionation	28
III. Enzymatic Degradation of Deoxyribonucleic Acids	30
(i) Digestion of deoxyribonucleic acid by DNase	30

	<u>Page</u>
(ii) Ion-exchange chromatography of DNA hydrolysates	32
(iii) Preparation of snake venom phosphodiesterase	35
(iv) Assay for phosphodiesterase activity	38
IV. Characterization of the Products of a DNA Hydrolysate	42
(i) Degree of polymerization of the oligonucleotides	42
(ii) Secondary degradation of DNA hydrolysates by venom and spleen phosphodiesterases	43
V. Effect of Heat, pH, Magnesium and Inhibitors on the Rate of Digestion of DNA by the Rat Intestinal Mucosal Extract. .	49
(i) Effect of Heat	49
(ii) Effect of pH	49
(iii) Magnesium requirements of the intestinal extract	49
(iv) Effect of inhibitors	50
VI. Intracellular Distribution of Deoxyribo- nuclease in the Rat Intestinal Mucosa . . .	52

RESULTS AND DISCUSSION

I. Preparation, Purification and Characteri- zation of Substrate Deoxyribonucleic Acids	55
II. Preparation and Purification of Deoxyribonuclease from Rat Intestinal Mucosa	59

	<u>Page</u>
(i) Effect of extracting media	59
(ii) Fractionation with ammonium sulfate	63
III. Enzymatic Degradation of Deoxyribonucleic Acids and Analysis of the Products	66
(i) Kinetics of reaction	67
(ii) Ion-exchange chromatography and characterization of the products in DNA hydrolysates	70
(iii) Secondary degradation of DNA hydrolysates by phosphodiesterases .	81
IV. Effect of Heat, pH, Mg^{++} and some Inhibitors on the Rat Intestinal Deoxyribonuclease	86
V. Intracellular Distribution of Deoxyribonucleases in the Rat Intestinal Mucosa	88
VI. General Discussion	91
<u>SUMMARY</u>	92
<u>BIBLIOGRAPHY</u>	95

LIST OF FIGURES

	<u>Page</u>
1. Change in UV-absorption Spectrum of Salmon Sperm DNA on Digestion with DNase	23
2. Rate of Increase in UV-absorption of Solution of Salmon Sperm DNA on Digestion with DNase I (Worthington Enzyme)	24
3. Nomogram for Ammonium Sulfate Solution . . .	29
4. Sodium Chloride Gradient in Ion-exchange Chromatography as Measured by the Refractive Indices of the Effluent Fractions	34
5. Time Course of the Hydrolysis of p-nitrophenyl-thymidine-5'-phosphate by Venom Phosphodiesterase	40
6. Comparison of Rate of Degradation of DNA by Crude Extracts of Rat Intestinal Mucosa	61
7. Time Course of the Degradation of DNA by Crude Intestinal DNase Extract as Measured by the Spectrophotometric Method of Kunitz .	68
8. Time Course of the Degradation of DNA by Crude Intestinal DNase Extract as Measured by Viscosimetry	69
9. Ion-exchange Chromatography of the Oligonucleotides in Rat Intestinal Mucosal DNA Hydrolysate Formed by Exposing the DNA to a Crude Intestinal DNase Preparation	71
10. Ion-exchange Chromatography of the Oligonucleotides in Salmon Sperm DNA Hydrolysate Formed by Exposing the DNA to a Crude Intestinal DNase Preparation . .	72

	<u>Page</u>
11. Ion-exchange Chromatography of the Oligonucleotides in Calf Thymus DNA Hydrolysate Formed by Exposing the DNA to a Crude Intestinal DNase Preparation	73
12. Ion-exchange Chromatography of the Oligonucleotides in Rat Intestinal Mucosal DNA Hydrolysate Formed by Exposing the DNA to the 20%P Enzyme	74
13. Ion-exchange Chromatography of the Oligonucleotides in Salmon Sperm DNA Hydrolysate Formed by Exposing the DNA to the 20%P Enzyme	75
14. Ion-exchange Chromatography of the Oligonucleotides in Calf Thymus DNA Hydrolysate Formed by Exposing the DNA to the 20%P Enzyme	76
15. Rate of Release of Mononucleotides by Snake Venom Phosphodiesterase from Salmon Sperm DNA Hydrolysate	83
16. Effect of pH on Deoxyribonuclease Activity of Rat Intestinal Mucosal Extracts	84

LIST OF TABLES

	<u>Page</u>
I. Elementary Composition and Some Other Properties of the DNA Preparations Used	58
II. Specific Activity of Deoxyribonuclease in Crude Extracts of Rat Intestinal Mucosa	62
III. Purification of Intestinal Mucosal Deoxyribonuclease Extract by Fractionation with Ammonium Sulfate	64
IV. Some Analytical Data of DNA Hydrolysates	78
V. Nucleoside Content of Salmon Sperm DNA Hydrolysate after Successive Degradation by Deoxyribonuclease, Spleen Phosphodiesterase and E. Coli Phosphomonoesterase	85
VI. Intracellular Distribution of Deoxyribonuclease in Rat Intestinal Mucosa	90

LIST OF ABBREVIATIONS

DNA	deoxyribonucleic acid
RNA	ribonucleic acid
s-RNA	soluble ribonucleic acid
DNase	deoxyribonuclease
RNase	ribonuclease
PME	phosphomonoesterase
EDTA	ethylenediamine tetraacetate
d-AMP, etc.	the 5'-phosphate of 2'-deoxyribosyl adenine, etc.
Pu	purine
Py	pyrimidine
poly d-U	a polymer of deoxyuridylic acid
g	gravitational force
gm	gram
mg	milligram
μg	microgram
M	molar
μ mole	micromole
mp	millimicron
μl	microliter

INTRODUCTION

Many, if not all, tissues have been shown to possess an enzymatic complement capable of degrading nucleic acids (1). It is more than likely that the function of this enzyme complement is not a passive one, i.e. they do not represent merely a means of degrading nucleic acids in dead or lysed cells so that the products formed may be used as substrates in nucleic acid synthesis by living members of the cell population. The possibilities are that they perform actively in many aspects of nucleic acid metabolism.

Genetic recombination has recently been shown to occur by a "breakage and reunion" of the DNA molecule (2). DNases, by analogy with pancreatic RNase (3), may under appropriate conditions act as a transoligonucleotidase transferring oligonucleotide segments from one polynucleotide to another. Studies of the replication of DNA by the polymerase enzyme isolated from E. Coli have demonstrated that DNA synthesis requires the presence of 3'-hydroxyl groups in the primer DNA (4). The rate of DNA synthesis can be greatly increased if the primer DNA is first subjected to treatment with DNase I to provide the necessary

3'-hydroxyl groups. Although indiscriminate cleavage of diester bonds by DNA endonuclease attack is inimical to biological integrity, the specific introduction of 3'-hydroxyl groups by a carefully controlled endonucleolytic fission may facilitate DNA synthesis in vivo.

Recently Setlow (64) proposed that one possible mechanism for correcting errors in the genetic code is by excision of the erratic segment by an endonuclease followed by repair of the excised portion. Thus deoxyribonucleases might represent a defence mechanism to counteract the effect of mutation.

DNases might also serve as "scavengers" within the cell, removing genetically superfluous and nonfunctional DNA segments formed, for example during replication (5). Doty et al. have used the exonuclease-I of E. coli to remove single-stranded polynucleotide ends from newly-synthesized, double-stranded DNA (5).

Despite the great variety and sometimes high concentration of nucleolytic enzymes present in cells, the nucleic acids do not undergo any precipitous breakdown except under extreme conditions of cell death and lysis. Obviously, mechanisms must exist to keep this powerful nucleolytic potency under control.

Studies of partially purified enzymes have in many instances given clues as to how they may be immobilized within the cell. For example, endonuclease-I in E. coli, which is DNA-specific, has been shown to be quantitatively inhibited by very low concentrations of RNA (6). This enzyme may thus be immobilized in vivo, by combination with the cellular RNA.

One of the two E. coli ribonucleases is found exclusively in the ribosomal fraction of cell-free extracts. Elson (7) showed that this ribonuclease activity is not detectable in carefully prepared ribosomal suspensions or ribonucleoprotein particles derived from them, but enzymatic activity appears upon disruption of the ribosomes by treatment with urea or trypsin. It is suggested that the ribosomal RNase exists only in a latent form and is therefore incapable of degrading RNA in a normally functioning cell.

Intracellular compartmentization is of course an effective way of preventing nucleases from acting on their substrates. But it is not always necessary to invoke individual peculiarities of catalytic specificity or intracellular location to explain the inactivity of these enzymes in vivo. The nucleic acids of the cell may be physically enveloped by, or otherwise bound to some structures (for

example, histone or protamines) and thus are inaccessible to the active sites of the enzymes.

Another reason which has prompted the study of various nucleolytic enzymes is the use to which these enzymes might be put as reagents in studying the structure and nucleotide sequence of nucleic acids, a branch of nucleic acid research which has received great impetus in the last few years. In the case of enzymes that attack RNA, it has been shown (8) that purified E. coli polynucleotide phosphorylase can rapidly phosphorolyse randomly coiled polyribonucleotides such as polyuridylic acids but ordered, hydrogen-bonded structures such as s-RNA are relatively resistant to attack by this enzyme. This property of polynucleotide phosphorylase suggests that the enzyme may be used in studying the secondary structure of polyribonucleotides.

Recently, Holley et al (9) were able to determine completely the nucleotide sequence in alanyl-s-RNA by exploiting the high degree of specificity of the two RNA depolymerases, pancreatic RNase and T₁-RNase.

Bacterial phosphatase, by virtue of its ability to phosphorolyse the terminal phosphomonoester linkage in oligonucleotides, and its lack of diesterase activity has

proved to be an excellent analytical tool for the determination of polynucleotide chain length and for the facilitation of the synthesis and depolymerization of polynucleotides by enzymes requiring free hydroxyl terminals for activity.

In the case of enzymes attacking DNA, depolymerases with any high degree of specificities are sadly lacking. The exonuclease-I of E. coli (10), because of its selectivity for single-stranded DNA, may be used for the identification of this conformation in a DNA sample. The enzyme is incapable of attacking the phosphodiester bond between the last two nucleotide residues at the 5'-terminus of a polydeoxyribonucleotide chain; this, coupled with its lack of endonuclease activity, permits the use of the enzyme as an endgroup reagent.

The present trend of nucleic acid research is directed chiefly towards the "cracking" of the genetic code. To this end, a direct comparison of the nucleotide sequence in DNA, the genetic material, and the primary structure of the polypeptide chain for which it codes, appears to be the ultimate aim. Such a feat cannot be accomplished until enzymes attacking DNA are discovered which show a high degree of selectivity towards the internucleotide bonds

which they hydrolyse. None of the DNases examined so far has shown any marked specificity, and the search for new enzymes capable of degrading DNA at specific internucleotide bonds must remain an imminent concern of the present-day biochemists.

The discovery of deoxyribonuclease dated as far back as 1903, when Araki observed that it was possible to liquefy gels of the β -form of thymus nucleic acid with extracts of tissues of liver, spleen and thymus (11). Iwanoff (12) demonstrated in various molds the presence of enzymes capable of liquefying thymus nucleic acid, and he applied the term nucleases to these enzymes. Iwanoff was also able to show experimentally that this nucleolytic activity did not rest with the proteases also present in the extract. This was confirmed by Sachs (13) who furthermore, found evidence that trypsin has a destructive action on nuclease. He was also able to demonstrate the presence of nucleases in fresh pancreas, in which the proteolytic enzymes are still in the zymogen state. Abderhalden and Schittenhelm (14) found that the pancreatic juice of dogs effected the liquefaction of thymus nucleic acid without liberation of inorganic phosphate or of nitrogenous bases. Feulgen (15) made the important observation that the

degradation of deoxyribonucleic acid did not progress to the mononucleotide stage, but stopped at the formation of oligonucleotides since the split products were still precipitable with mineral acids. Schmidt, Pickels and Levene (16) found in 1939 that the degradation of DNA to oligonucleotides by the pancreatic enzyme was an essential intermediary step for the enzymic cleavage of DNA to mononucleotides (and subsequently to nucleosides) by the phosphatase of intestinal mucosa. Highly polymerized deoxyribonucleic acid was not attacked by purified intestinal phosphatase (nucleotidase); but it hydrolysed all interlinkages of the mononucleotide groups in the oligonucleotide mixture obtained by the action of deoxyribonuclease (16).

The evidence for the existence of a specific polynucleotidase, both for ribonucleic and deoxyribonucleic acids, though more or less convincing, has been entirely circumstantial even as late as 1939. It was only with the isolation of crystalline ribonuclease by Kunitz (17) in 1940 that the presence of a specific polynucleotidase, at least for RNA, was definitely established. Since then, attempts to obtain purified preparations of deoxyribonuclease from pancreas preparations were made by Laskowski (18) and

McCarty (19). They were able to separate deoxyribonuclease from the very powerful ribonuclease of pancreas and to provide evidence for the strict specificity of deoxyribonuclease for DNA. The crystallization of the enzyme was achieved by Kunitz in 1950 (20).

The pH optimum of this pancreatic deoxyribonuclease lies close to 7.0, but it is somewhat influenced by the nature of the ions in the medium (20). The enzyme requires divalent cations for activity, magnesium, manganese and cobaltous ions being the most effective (20, 21). Another DNase which has an optimum pH in the range 4.5-5.5 and is inhibited by the addition of Mg^{++} was observed in spleen by Catcheside and Holmes (22) and in thymus by Maver and Greco (23). These observations were soon confirmed and extended to other tissues. The name DNase II was suggested for this enzyme in contrast to the pancreatic enzyme, which was termed DNase I.

At present, the DNases are subdivided into two classes, the DNase I enzymes and DNase II enzymes. These two classes were introduced by Laskowski (24) to compare and to contrast enzymes distributed in different tissues with their prototypes in respect to pH optimum and activating and inhibiting agents. Pancreatic DNase I of Kunitz and

thymus DNase II of Maver and Greco served as prototypes. This terminology is useful for the purpose of characterization of enzymes in different tissues. In accordance with this, the term "intestinal mucosal DNase I" implies an enzyme isolated from the intestinal mucosa and resembling pancreatic DNase in its properties.

Laskowski (25) proposed four criteria for the characterization of deoxyribonucleases: the susceptible substrates, the type of attack, the products and the preferential linkage. When DNase I is considered under these headings, it is to be seen that DNase I is specific for DNA. It is an endonuclease because it rapidly decreases the viscosity of a DNA solution and because the products of hydrolysis are predominantly large oligonucleotides. Titration data indicated that approximately 25% of all available internucleotide linkages could be hydrolysed by DNase I (20). This means that the average size of the product is that of a tetranucleotide. Ion-exchange chromatography showed that the products were oligonucleotides of a variable size, ranging from mononucleotides through di-, tri- etc., probably to octanucleotides (27). The products were identified as terminated in 5'-phosphates (26). The mononucleotides were susceptible to the action of 5'-nucleotidase and the larger

fragments yielded 5'-nucleotides exclusively on digestion with venom phosphodiesterase.

A systematic identification of products (28) established that all four of the common mononucleotides were present, with thymidylic acid being the predominant member. Sinsheimer has carried out extensive studies of the dinucleotides in DNase I digests (29). The dinucleotides with sequences pPy-pPy, pPu-pPu and pPy-pPu were found to be abundant but those with sequences pPu-pPy were very rare. This was interpreted to mean that the pPu-pPy bond was preferentially hydrolysed. Additional evidence was obtained by isolating a trinucleotide - ApApTp from a digest of DNA by DNase II, and then digesting it with DNase I (30). The trinucleotide was split into d-ApA and pTp by DNase I, as would be expected from the postulated preference. However, this preference cannot be expressed in quantitative terms.

Koerner and Sinsheimer (31) showed that DNase II from spleen was also an endonuclease. Exhaustive reaction resulted in about 25-30% of the internucleotide linkages being hydrolysed. The distribution of the fragments as evidenced by the chromatographic pattern was quite different from that of a digest by DNase I. DNase II produced more mononucleotides, considerably less dinucleotides and considerably

more of the higher oligonucleotides. All the reaction products were terminated in 3'-phosphates. DNase II did not seem to show any preference for specific purine and pyrimidine bases adjacent to the sensitive linkages.

Both DNase I and DNase II are widely distributed (1). Pancrease has long been recognized as the best source of DNase I, whereas spleen and thymus are much richer in DNase II than DNase I activities. Recently, Stewart and Zbarsky (32) reported the loss of DNA from rat intestinal mucosa during incubation in vitro of mucosal scrapings or intestinal slices and have pointed out the possible existence of a DNase in the tissue. This loss of nucleic acids from the incubated tissue was not inhibited by removal of Mg^{++} from the buffer, nor by the addition of 0.1 M arsenate. Addition of EDTA however reduced the loss of DNA to only 11% in one hour. These results appeared to indicate that both types of DNases were present in the rat intestinal mucosa and it seemed that the isolation and characterization of the intestinal enzyme(s) were desirable and might provide some insight into the nucleic acid metabolism of this actively proliferating tissue.

A method was developed for preparing crude enzyme extracts of the intestinal mucosal tissue using a variety of

extracting media. The activities of these extracts were tested against deoxyribonucleic acids from several sources. Methods used for the determination of DNase activities were based on viscosity and optical properties of nucleic acids and their components.

Employing the technique of fractionation by means of ammonium sulfate, it was found that two fractions of DNase activity were precipitable from the mucosal extract. One of these precipitated at 20% saturation of ammonium sulfate and the activity was recoverable by redissolving the protein precipitate in buffer. The other type precipitated at 30-35% saturation of ammonium sulfate, but all attempts to recover the enzymatic activity were unsuccessful. In subsequent experiments performed, both the crude extract and the partially purified enzyme preparation were therefore used separately so that a qualitative comparison between the two could be made.

The effect of pH, the ionic requirements and the effect of various inhibitors were tested and found to differ markedly for the two enzyme preparations. Thus whereas the partially purified enzyme extract showed an absolute requirement for Mg^{++} , removal of magnesium by exhaustive dialysis of

the crude extract did not completely abolish its activity. A protein inhibitor prepared from the pigeon crop gland, while completely inhibiting the purified extract, had only limited effect on the crude preparation. These findings would seem to indicate that the crude preparation contained an additional DNase different from the one precipitated at 20% saturation of ammonium sulfate.

To further clarify this point, the digests obtained by exposing DNA to the two fractions were analysed. The elution profiles from a DEAE-cellulose column were found to be different. The oligonucleotides obtained with the partially purified enzyme extract were exclusively those which contained a 5'-phosphomonoester bond, but the products obtained with the crude complete extract contained both 5'- and 3'-phosphomonoester linkages. Moreover, the corresponding oligonucleotide fractions obtained with the two extracts were found to differ both qualitatively and quantitatively. Thymidylic acid was predominant in the mononucleotide fractions; but whereas the d-cytidylic acid content in the higher oligonucleotides was low when the digest was prepared with the purified extract, no such discrimination was observed when the crude enzyme was used.

The possibility was considered that the deoxyribonuclease activity of the intestinal mucosal extract might be due to contamination of bacterial or pancreatic origin. Extra care has been taken to flush clean the intestinal lumen during the preparation of the enzyme extract; and it has also been found that after a whole cell preparation of the intestinal mucosa was extracted, further deoxyribonuclease was released into the medium when the tissue was homogenized. To further pinpoint the mucosal origin of the enzyme, it was felt that an intracellular localization and distribution of the deoxyribonuclease activity must be established. To this end, a mucosal homogenate was subjected to differential centrifugation and the various subcellular fractions assayed for DNase activity. It was found that whereas 54% of the total DNase activity was located in the heavy mitochondrial fraction, 43% of the activity was associated with some lighter particulate fraction.

On the basis of the results obtained in the experiments to be described, it has been concluded that the deoxyribonuclease activity extracted from the intestinal mucosa of the rat may contain both DNase I and DNase II types. The experiments have, however, not unequivocally established the existence of DNase II. A positive identification of this enzyme has to await its isolation from the

tissue and the characterization of its products of reaction. The problem of bacterial contamination has also not been satisfactorily solved. The use of germ-free rats would undoubtedly clarify this point.

EXPERIMENTAL

I. Preparation of Deoxyribonucleic Acids

- (1) Preparation of DNA from Small Intestinal Mucosa of Rat - Method of Colter et al.

Reagents:

- (a) 1M NaCl-0.03M EDTA - 0.02M phosphate buffer, pH 7.3.
- (b) Water-saturated Phenol: this solution contains 75 gm. of redistilled phenol and 25 ml. of dist. H₂O.
- (c) 0.15M NaCl-0.03M EDTA. The pH of the solution was adjusted to 7.0 with 0.1N NaOH.
- (d) 10% Na deoxycholate solution.

Procedure:

Male Wistar rats weighing 180-200 gm. each were starved for 24 hours. They were then killed by a blow on the head and decapitated. The small intestine was removed and cut into segments approximately 10 cm. in length. These segments were flushed entirely free of contents with chilled phosphate buffer. They were then slit open and applied to a cold glass surface with the mucosal side facing upwards. The mucosal epithelium was then scraped off with the edge of a microscope slide and immediately placed in liquid nitrogen.

Five rats afforded about 7 gms. wet weight of mucosal tissue. The pooled frozen tissue was transferred to the blender cup of a Servall Omni Mixer and 45 ml. of NaCl-EDTA-Phosphate buffer solution, 2.5 ml. of 10% sodium deoxycholate solution and 50 ml. of water-saturated phenol solution were added for each 5 gm. of tissue used. The blender cup was set in an ice bath and the mixture was homogenized at a low speed, the powerstat setting being at 20v. Following homogenization, the emulsion was centrifuged for 10 min. at 27,000g in a Servall refrigerated centrifuge. The upper aqueous layer containing the DNA was carefully removed from the lower phenol layer and denatured protein which collected at the interface. The aqueous solution was subjected to two additional extractions with equal volumes of water-saturated phenol. These extractions were carried out in the cold room on a mechanical shaker and were of 5 min. duration. After each extraction, the aqueous and phenol layers were separated by centrifugation at 4,000g in the Servall centrifuge for 10 min.

Traces of phenol were then removed from the DNA solution by 6 or 7 extractions with ether, and the residual ether removed by bubbling nitrogen through the solution. The DNA solution was then kept overnight at 4°C; the high molecular weight RNA which precipitated out of solution was

removed by centrifugation at 4,000g for 10 min. The DNA was precipitated by the addition of an equal volume of ethanol. The DNA precipitate was dissolved again in the minimum amount of neutral saline-EDTA solution and centrifuged if necessary to obtain a clear solution. DNA was precipitated with ethanol as previously, and the fibrous material was washed several times with 75% EtOH and then dried in vacuo over P_2O_5 at room temperature. The dry preparation was dissolved in the appropriate buffer solution, usually to a concentration of 1 mg. DNA/ml. The solutions were stored at $-20^{\circ}C$.

(ii) Deproteinization of Salmon Sperm DNA

A sample of crude salmon sperm DNA, a gift from Dr. Tener was found to be heavily contaminated with protein. It was felt desirable to remove the protein before the DNA was used as substrate for enzymatic degradation. Deproteinization was carried out essentially according to the method described by Zubay (34).

Procedure:

Five hundred mg. of the crude salmon sperm DNA was dissolved in a minimum amount of distilled water. The volume of the solution was measured and enough NaCl added to give a solution 1M in strength with respect to NaCl.

To approximately 400 ml. of this DNA solution, 100 ml. of chloroform-amyl alcohol mixture (3:1 v/v) were added. The mixture was emulsified by shaking it in the cold for 10 min. using a vibratory shaker, and was then centrifuged at 4,000g on the Servall refrigerated centrifuge for 10 min. The upper aqueous layer was carefully removed with a syringe and separated from the lower organic phase and the layer of denatured protein collected at the interface. The extraction was repeated until no more protein was observed at the interface. The DNA was then precipitated by adding an equal volume of ethanol and the preparation treated as described in I(i).

(iii) Estimation of Nitrogen Content of Deoxyribonucleic Acids

The nitrogen content of deoxyribonucleic acids was determined by the method of Pregl (35). The recovery of nitrogen by this procedure was very satisfactory. When recrystallized urea was used as nitrogen standard, it was found to contain, on the average of three experiments, 46.2% of nitrogen as compared to the theoretical value of 46.67%.

(iv) Determination of Phosphorus Content of DNA

The color reaction of Fiske and SubbaRow (36) was used for phosphorus determination. One mg. of dry DNA was used routinely for such determinations. The colour produced

was found to be proportional to the concentration of phosphorus up to 40 ug for 10 ml. of the reaction mixture. A calibration curve was constructed (using KH_2PO_4 as standard) from which the phosphorus content of the DNA samples was determined.

(v) The Determination of $\epsilon(\text{P})$ Values of DNA Preparations (37).

- (a) 0.1M ammonium acetate buffer - 6 gm. of glacial acetic acid and 2.03 gm $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ were dissolved in approx. 900 ml. of dist. H_2O . The pH of the solution was adjusted to 7.0 with conc. NH_4OH . The final volume of the solution was made up to 1,000 ml.

Procedure:

Accurately weighed DNA samples were dissolved in the acetate buffer so that the final concentration of DNA was 1 mg./ml. One ml. of this stock solution was made up with the same buffer to 25 ml. The ultraviolet absorption spectra of the solutions were then recorded with a Cary spectrophotometer. From the optical density value at maximum absorption (usually around 258 m μ) the $\epsilon(\text{P})$ value of the sample was calculated according to the equation (37):-

$$\epsilon(P) = \frac{\text{O.D. at max.}}{C \times d}$$

where $\epsilon(P)$ = the atomic extinction coefficient with respect to phosphorus

O.D. = optical density of the solution

C = phosphorus concentration of the solution in moles/liter

d = internal cell length in cm.

II. Preparation and Purification of Deoxyribonuclease from Rat Intestinal Mucosa.

(i) Measurement of Enzymatic Activity

(a) Spectrophotometric Method of Kunitz (20)

One of the effects of the action of deoxyribonuclease on DNA is the increase in the absorption of ultra-violet light by the digested acid. The rate of increase in the light absorption is gradual and can be readily determined. It was found (20) that the rate of increase of the optical density at 260 m μ , brought about by slow digestion of DNA in dilute solution at the appropriate pH, was constant for several minutes and was proportional to the concentration of enzyme in solution. It can thus be used as a convenient and rapid method for measuring deoxyribonuclease activity.

Reagents:

(a) Substrate DNA solutions: 40 μ g/ml in 0.1M ammonium

acetate buffer, 0.01M in MgCl_2 , pH 6.5. The solution was stable for weeks when stored in the refrigerator at about 5°C .

- (b) DNase I solution - 2 mg. DNase I (Worthington) per 10 ml. acetate buffer.

Procedure:

Three ml. of the substrate DNA was pipetted into a quartz cuvette (1 cm. light path). At zero time, 0.1 ml. of enzyme solution was added and the solution was mixed thoroughly. The increase in optical density was measured at 260 $\text{m}\mu$ with a Cary recording spectrophotometer equipped with a thermostatable cell compartment maintained at 37°C . The spectra of the DNA solution before the addition of the enzyme and after the completion of the reaction were measured.

Figure 1 shows the increased U.V. absorption at 260 $\text{m}\mu$ obtained during the course of depolymerization of DNA by DNase I. In Figure 2, it can be seen that when the increase in optical density was plotted against the incubation time, the initial portion of the graph was a straight line. The activity of the enzyme solution is expressed in terms of the initial slope of the plotted curve. This divided by the concentration of the enzyme in mg. of protein per ml. of the final digestion mixture gives the specific activity of the

FIGURE 1 : Change in UV-absorption spectrum of Salmon Sperm DNA on Digestion with DNase.

FIGURE 2 : Rate of Increase in UV-absorption of Solution of Salmon Sperm DNA on Digestion with DNase I (Worthington Enzyme).

enzyme in units of deoxyribonuclease activity per mg. of protein. One unit of activity is thus defined as the amount of enzyme capable of bringing about an increase in optical density at 260 m μ of 1 per minute under the given conditions of concentration of the substrate, pH, temperature and salt concentration.

(b) Viscosimetric Method (38)

Measurement of the progressive fall in viscosity of DNA solutions during treatment with DNase has yielded constant and reproducible results and has proved a reliable method for determining enzymatic activity.

Reagents and Apparatus:

- (a) Substrate DNA solution - 1 mg. DNA/ml. in 0.1M ammonium acetate buffer, 0.01M in MgCl₂, pH 6.5.
- (b) Commercial DNase I (Worthington) solution
- (c) Ostwald type capillary viscosimeter

Procedure:

A thoroughly cleaned and dry viscosimeter was clamped vertically in the thermostated bath (37°C) in such a position that it could be viewed easily, and 5 ml. of DNA solution was added from a pipette. The liquid was drawn up into the enlarged bulb and above the upper mark. The liquid was then permitted to run down through the capillary

and the stop watch was started when the meniscus passed the upper mark and stopped when it passed the lower mark. Two or three check determinations on the flow rate were made. The flow time for the solvent (acetate buffer) was similarly determined.

At zero time, 0.1 ml. of enzyme was added and thoroughly mixed with the substrate. The flow time was measured at 5 min. intervals over a period of about 30 minutes.

The activity of a given enzyme can be determined from the slope of a curve obtained by plotting the relative viscosity, η , of the reaction mixture against time. Relative viscosity is defined by the term T_t/T_o , where T_t is the flow time at time t . Where an understanding of the kinetics of the depolymerization reaction is desired, the velocity constant K of the reaction at any given time t can be determined from the equation

$$K = \frac{1}{t} \log \eta_o / \eta_t$$

The viscosimetric technique was used more extensively in the earlier experiments during the course of this research. But because of its requirement for large amounts of substrate DNA, it was later abandoned in favour of the spectrophotometric

method and was then used only occasionally for the qualitative demonstration of deoxyribonuclease activity, where the products of digestion could be used for other experiments.

(ii) Preparation of the Crude Enzyme Extract.

Procedure:

Five male Wistar rats, weighing about 180-200 gms. each, were sacrificed as usual. The intestinal mucosal tissue was removed as previously described and immediately placed into a 50 ml. graduated centrifuge tube standing in an ice bath. Enough of Kreb's Ringer phosphate buffer (39), pH 7.8, was then added to give a 1/10 (v/v) suspension of the tissue. The tissue clumps were broken up by drawing the suspension through a size 14 syringe needle. The suspension was then centrifuged at 105,000g for 60 minutes at 0°C in the Spinco Model L ultracentrifuge. The supernatant solution was collected and tested for deoxyribonuclease activities as described under II(i).

In some experiments, the pellets after centrifugation were washed with water several times and then centrifuged. They were then re-suspended in Krebs Ringer phosphate buffer equal in volume to that used in the first extraction. The suspension was then homogenized in the Servall Omni Mixer

set at 30v. on the rheostat for 15 minutes at 0°C. The homogenate was centrifuged at 105,000g for 60 minutes. The supernatant solution was collected and examined for deoxyribonuclease activity.

(iii) Purification of Intestinal Mucosal Deoxyribonuclease by Ammonium Sulfate Fractionation.

Procedure:

Ammonium sulfate fractionation was carried out by a modification of the method described by Kunitz (20).

A known volume of the crude enzyme extract was brought to 5% saturation with ammonium sulfate (40). The precipitate formed was removed by centrifugation at 16,000g in the Servall refrigerated centrifuge for 60 minutes. A known volume of the supernatant solution (designated 5%S) was removed and examined for nitrogen and protein contents and DNase activity. The precipitate was redissolved in a volume of buffer equal to the volume of the crude enzyme extract used and similarly examined. This solution was designated 5%P.

The rest of the supernatant solution was further charged with ammonium sulfate to a desired saturation of salt and the above procedure repeated. The amount of ammonium sulfate required to bring a solution to a degree of

FIGURE 3 : Nomogram for Ammonium Sulfate Solution.

A straight line through the initial saturation and the desired saturation gives the amount of solid $(\text{NH}_4)_2\text{SO}_4$ to be added to 1 liter of this solution. A line from this point passing through the volume of the solution gives the amount required.

saturation was determined according to Dixon (40). Figure 3 shows how this can be done with the use of a nomogram.

The fractionation was done by increasing steps of ammonium sulfate saturation of 5% intervals. The various fractions isolated were analysed for deoxyribonuclease. The specific activity of the enzyme solution was expressed in terms of enzyme activity per mg. protein.

Both the crude and the partially purified enzyme extracts were stable for several months if stored at -20°C .

III. Enzymatic Degradation of Deoxyribonucleic Acids

(i) Digestion of Deoxyribonucleic Acid by DNase.

Reagents:

(a) Substrate DNA solutions

Three types of DNA's were used: rat intestinal DNA, deproteinized salmon sperm DNA and calf thymus DNA (sodium salt, Mann Assayed product). They were dissolved separately in 0.1M acetate buffers, 0.01M in MgCl_2 , pH 6.0, 6.5 or 7.0. When used as substrates for DNase II, the DNA's were dissolved in 0.33M sodium formate buffer, pH 4.5. In each case, the concentration of DNA was 1 mg./ml.

(b) Enzyme preparations

(i) The crude enzyme extract

(ii) "20% P" enzyme preparation

(iii) DNase I (Worthington), 100 $\mu\text{g}/\text{ml}$ in acetate buffer, pH 7.0

(iv) DNase II (Worthington), 100 $\mu\text{g}/\text{ml}$. in 0.3M sodium formate buffer, pH 4.5.

Procedure:

To 20 ml. of substrate solution in a clean test-tube was added 1 ml. of the enzyme preparation and the mixture incubated at 37°C . The pH of the reaction was maintained constant by addition of 0.001N NaOH at intervals to neutralize the secondary phosphate group released during depolymerization. The reaction was followed by observing the optical density changes of the solution at 271 $\text{m}\mu$, and was found to be essentially complete after 4 hrs. However, to ensure complete degradation, the incubation was carried out for a further 20 hrs., with a drop of chloroform added as preservative. During the latter period, no attempt was made to adjust the pH of the solution.

The reaction mixture was next heated at 100°C for 5 min. to precipitate the enzyme protein. It was then cooled to room temperature and centrifuged at 16,000g for 60 min. The supernatant solution was collected and stored at -20°C for future use.

(11) Ion-exchange Chromatography of DNA Hydrolysates (41)

Reagents:

- (a) 7M urea solution
- (b) 7M urea solution 1M in NaCl
- (c) 0.1M Tris buffer, pH 7.8
- (d) 4M NaCl solution
- (e) 1M NaCl solution
- (f) 2M ammonium carbonate solution
- (g) 0.5M $(\text{NH}_4)_2 \text{CO}_3$ solution
- (h) DEAE-cellulose (Brown Co., Berlin, N.H.)

Procedure:

The DEAE-cellulose was suspended and decanted repeatedly with dist. water to get rid of fines. A thin slurry was made of the settled cellulose by adding 1M NaCl solution (or 0.5M ammonium carbonate) and poured into a chromatographic column half-filled with the same solution. (The packing of the column was done in the presence of salt to ensure a uniform bed). A pressure of 5 lbs./sq. in. was used for the packing. Excess liquid was drained from the column and fresh slurries were added until the desired bed depth was attained.

The columns used had an inner diameter of 1 cm. Those in the chloride form were packed to a depth of 30 cm., and those in the carbonate form to a depth of 10 cm.

After packing, the columns were washed with 4M NaCl or 2M $(\text{NH}_4)_2\text{CO}_3$ until the effluent showed no absorption at 260 m μ . The salts were then removed by washing with distilled water (calcium hydroxide test for $\text{CO}_3^{=}$ and silver nitrate test for chloride ions).

Five ml. of the DNA hydrolysate (approx. 450 ug. P) were added to the DEAE column (chloride form) and washed in with a small amount of water. The elution was begun with a linear gradient formed by 1 liter of 7M urea in the mixing chamber to which 30 ml. of the tris-HCl buffer had been added; the other chamber contained 1 liter of 7M urea solution, 1M in NaCl, also containing 30 ml. of the buffer. The flow rate was adjusted to approximately 1 ml. per min. Five-ml. fractions were collected by means of an automatic fraction collector (Gilson, constant volume operation) equipped with a spectrophotometric attachment for measuring the ultraviolet absorption of the eluate. The optical density of the individual fractions was also measured with a Zeiss spectrophotometer at 260 m μ .

The sodium chloride gradient was approximated in a control experiment in which no DNA digest was added to the column. The increase in NaCl concentration of the effluent

FIGURE 4 : Sodium Chloride Gradient in Ion-exchange Chromatography as Measured by the Refractive Indices of the Effluent Fractions.

was checked by measuring the refractive indices of the effluent fractions with an Abbe 60 refractometer (Bellingham and Stanley Inc.). The curve thus obtained is shown in Figure 4.

Isolation procedure:

Fractions comprising a peak were pooled and diluted with 5 volumes of distilled water to reduce the salt concentration. The solution, if necessary, was adjusted to pH 8. It was next run onto a DEAE-cellulose column in the carbonate form (10 x 1 cm.) and the column washed with water until the effluent was free of urea and inorganic ions. The optical densities of the washings were checked to ensure no leakage of nucleotides from the column occurred. The oligonucleotide was then eluted with 2M ammonium carbonate and located among the fractions with a Zeiss spectrophotometer. The UV-absorbing fractions were combined and evaporated repeatedly in a flash evaporator with a bath temperature not exceeding 30°C until all the ammonium carbonate was removed. The residue was dissolved in the appropriate solvent for characterization and other experiments.

(iii) Preparation of Snake Venom Phosphodiesterase

Materials:

Lyophilized Crotalus adamanteus venom was obtained

from the Ross Allen Reptile Inst. Carboxymethyl-cellulose was purchased from Carl Schleicher and Schnell Co., Keene, N.H. under the trade name of Selectacel, No. 77, Type 20, capacity 0.72 meq./gm.

Reagents:

- (a) 0.05M acetate buffer, pH 4
- (b) 0.05M acetate buffer, pH 6
- (c) 0.2M acetate buffer, pH 6
- (d) 0.5M acetate buffer, pH 6

Procedure:

(a) The preparation of venom phosphodiesterase was done according to the method described by Koerner and Sinsheimer (31). One gm. of crude venom was dissolved in 60 ml. of water and the solution was left standing at room temperature for one hour. It was then centrifuged at 3,000g on the Servall refrigerated centrifuge for 15 min. The supernatant solution was collected and 40 ml. of 0.05M ammonium acetate buffer, pH 4, was added to it. All subsequent procedures were carried out in the cold room at about 4°C.

The solution was transferred into polyethylene tubes and 74 ml. of acetone at -20°C was added to the buffered venom solution. The contents of the tube were mixed thoroughly and left standing for 30 min. The solution was then centri-

fuged at 3,000g for 30 min. and the supernatant solution transferred to another polyethylene tube. Fourteen ml. of acetone were added next and the solution was stored for 12 hours at cold room temperature. It was then centrifuged again and the supernate was further charged with 12 ml. acetone. The final concentration of acetone in the solution was 50%. After standing for 30 min., the mixture was centrifuged, and the precipitate was dissolved in the minimum amount of distilled water. This solution was next dialysed against 0.05M ammonium acetate buffer at pH 6 and further purified by ion-exchange chromatography.

(b) Chromatography on carboxymethyl (CM) cellulose was done according to Laskowski et al. (42). A CM-cellulose column, 2 x 18 cm., was equilibrated with 0.05M acetate buffer, pH 6. The dialysed protein solution was then added to the column. The protein was eluted stepwise using 1) 90 ml. of 0.05M acetate buffer, pH 6; 2) 90 ml. of 0.2M acetate buffer, pH 6 and 3) 90 ml. of 0.5M acetate buffer, pH 6. The flow-rate was adjusted to about 6 ml. per hour. The fractions were assayed for UV-absorbing material at 280 mμ with a Zeiss spectrophotometer, and for phosphodiesterase activity.

(iv) Assay for Phosphodiesterase Activities

- (a) Qualitative assay for venom phosphodiesterase activity in fractions collected from the CM-cellulose column.

Substrate solution: 13.6 mg. of p-nitrophenyl-thymidine-5'-phosphate in 25 ml. of 0.1M tris-0.002M magnesium acetate buffer.

Procedure:

The assay method was developed by C. Mezei (43). 0.1 ml. of the substrate solution was pipetted into a micro test-tube and incubated in a water bath at 37°C with constant shaking. One drop of solution from a given fraction was added to the test-tube and the appearance of the yellow p-nitrophenylate in one minute was assumed to indicate the presence of phosphodiesterase activity.

Two protein peaks were observed in the effluent from the CM-cellulose column, one being eluted by the 0.05M acetate buffer, the other by the 0.2M acetate buffer. Both peaks were found to contain phosphodiesterase activity. This is in accordance with observations by Khorana and Razzell (44). The peak eluted with 0.2M acetate buffer was subsequently used as the stock enzyme solution, as this was found to be free of 5'-nucleotidase activity.

(b) Quantitative assay for phosphodiesterase activity (44).

In this method, 1.5 ml. of substrate solution and 1.4 ml. of the Tris-HCl buffer were pipetted into a 1-cm. cuvette and maintained at 37°C in the thermostated cell chamber of the Cary spectrophotometer. At zero time, 0.1 ml. of the enzyme solution was added and the solution mixed thoroughly. The rate of increase in optical density at 400 mμ due to the liberation of the yellow p-nitrophenylate was a measure of enzyme activity.

In a reaction mixture volume of 3 ml., containing 1.5 μ mole of substrate, an increase in optical density of 1.2 units is effected by the hydrolysis of 0.1 μ mole of the substrate.

The pooled fractions from the CM-cellulose column had to be diluted 100 fold with the buffer before the diesterase activity contained in it could conveniently be measured. Figure 5 shows that when the optical density was plotted against time, the change in absorption varied linearly with time for at least ten minutes. During this period, the diluted enzyme preparation effected an increase of 0.039 O.D. units per minute corresponding to the hydrolysis of 1.95 μ mole of substrate per hour.

FIGURE 5 : Time Course of the Hydrolysis of p-nitrophenyl-thymidine-5'-phosphate by Venom Phosphodiesterase.

When a standard preparation of venom phosphodiesterase (Worthington, 10 ug./ml.) was used, it was found to hydrolyse 8.2 μ mole of substrate per hour. One ml. of the purified dilute phosphodiesterase solution therefore corresponds to 2.4 μ g. of the Worthington enzyme.

(c) Qualitative assay for spleen phosphodiesterase

Reagents:

- (a) a spleen phosphodiesterase solution was given by Dr. W.E. Razzell.
- (b) Substrate solution: a 0.03 solution of synthetic p-nitrophenyl thymidine-3'-phosphate was also a gift from Dr. W.E. Razzell.

Procedure:

The assay method was similar to that described for venom phosphodiesterase. The spleen phosphodiesterase solution had been in storage (-20°C) for several years, but was still found to possess considerable hydrolysing activity towards the substrate.

Both spleen and venom phosphodiesterases were used in the characterization of the products of DNA hydrolysates by the intestinal enzyme(s) as described under IV.

IV. Characterization of the Products of a DNA Hydrolysate

(i) Degree of polymerization of the oligonucleotides

Reagents:

- (a) Phosphomonoesterase preparation - the Worthington E. coli phosphomonoesterase suspension in ammonium sulfate was diluted 10 times with 0.01M tris buffer, 0.01M in $MgCl_2$, pH 8.0.
- (b) 60% $HClO_4$
- (c) 5% ammonium molybdate solution
- (d) Fiske-SubbaRow reagent (36)

Procedure:

- (a) Estimation of phosphomonoesterase-sensitive phosphorus.

This was performed by a slightly modified method used by Snider et al. (45). Each of the dry oligonucleotide fractions initially prepared as described under III(ii) was dissolved in 1 ml. of 0.01M tris buffer, 0.01M in $MgCl_2$, pH 8.0. The optical density of the solution was measured.

Duplicate samples containing 1 O.D. unit^{*} of UV-

* 1 O.D. unit of oligonucleotide is defined as the amount of material which gives rise to an optical density of 1 at 260 m μ when present in 1 ml. of solution and when measured in a cuvette with light path = 1 cm.

absorbing material from each of the oligonucleotide fractions were made up to 1 ml. with tris buffer, pH 8.0. These were transferred to 10-ml. volumetric flasks. Ten μ l. of the phosphomonoesterase preparation was added to each flask and the mixture incubated for 16 hours at 37°C. A drop of chloroform was routinely added to eliminate bacterial contaminations. After the digestion, the mixture was diluted to 6 ml. with distilled water and 0.8 ml. of 60% perchloric acid was added, followed by 1 ml. of molybdate solution and 0.4 ml. of Fiske-SubbaRow reducing reagent. The solution was well shaken after each addition. The final volume was made up to 10 ml. and the optical density of the solution was measured at 830 m μ . The phosphate liberated by the alkaline phosphatase was determined from a previously prepared standard curve.

(b) Estimation of total phosphorus.

One ml. of solution containing 1 O.D. unit of oligonucleotide in tris buffer was digested with 2 ml. of conc. perchloric acid to liberate all the phosphate, which was then determined as described under I (iv).

(ii) Secondary Degradation of DNA Hydrolysates
by Venom and Spleen Phosphodiesterases.

Materials and Reagents:

- (a) Sephadex G-25, purchased from Pharmacia,
Upsala, Sweden.

- (b) 0.1M Tris buffer, 0.002M in magnesium acetate, pH 8.9.
- (c) 0.01M Tris buffer, 0.01M in magnesium acetate, pH 8.
- (d) 0.18M ammonium acetate buffer, 0.06M in EDTA, pH 5.7.
- (e) spleen phosphodiesterase
- (f) venom phosphodiesterase
- (g) E. coli phosphomonoesterase
- (h) Standard deoxyribomononucleotides and deoxyribonucleosides

The four common deoxyribomononucleotides and the four deoxyribonucleosides in DNA were used: d-adenylic acid, d-cytidylic acid, d-guanylic acid (sodium salt), d-thymidylic acid, d-adenosine, d-cytidine, d-guanosine and thymidine. These were purchased from the Calif. Foundation for Biochem. Research, L.A., Calif. and standard solutions were prepared by dissolving 2 mg. of each substance in 1 ml. water. Where necessary, dil. HCl was added to effect complete dissolution.

- (i) Chromatographic system I - isopropanol: conc. $\text{NH}_4\text{OH}:\text{H}_2\text{O}$ in the ratio (v/v) 7:1:2.
- (j) Chromatographic system II - 100 ml. water saturated butanol plus 1 ml. conc. NH_4OH .

Procedure:

- (a) Desalting of DNA hydrolysate by gel filtration (46).

The Sephadex-G-25 powder was suspended in distilled water to swell. It was then washed several times to remove fines. A slurry (10 volumes of water to 1 vol. of gel) was prepared. The Sephadex column (10 x 1 cm.) was then packed at atmospheric pressure. The sample to be desalted was then poured onto the column and the material eluted with distilled water. The fractions were collected and assayed for ultra-violet absorbing material at 260 $m\mu$ and for chloride ions.

The fractions containing oligonucleotides were pooled, dried in vacuo and the residue redissolved in the appropriate buffer.

- (b) Action of venom phosphodiesterase on DNA hydrolysates.

The desalted DNA hydrolysate (containing 5 mg. of oligonucleotides) obtained from IV (11a) was dissolved in 5 ml. of 0.1M tris buffer (pH 8.9), 0.002M in magnesium acetate. One ml. of this solution was incubated with an excess (0.5 ml.) of venom phosphodiesterase solution at 37°C. At appropriate intervals of time, 100 μ l. of the reaction mixture were withdrawn and mixed with 10 μ l. of glacial acetic acid to stop the reaction. The mixture was then spotted onto

Whatman No. 40 filter paper. Fifty μ g. of each standard deoxyribomononucleotide were also spotted onto the same paper. The chromatogram was then developed in the isopropanol: NH_3 : H_2O system for 72 hours, employing the descending technique. The nucleotide spots were located under ultraviolet light. These were cut out and eluted with approximately 5 ml. of 0.1N HCl (43). The eluates were evaporated to dryness and the residues taken up in an exact volume of distilled water appropriate for the cells in which the extinctions were to be measured. The ultraviolet light absorption curves were recorded with a Cary spectrophotometer. Blank spots approximately equal in area to the nucleotide spots and at equal distances from the origin were cut out, eluted and read at the same wavelengths as the corresponding spots.

The amount of deoxyribonucleotides in the spots was determined from the optical densities of their solutions at the appropriate wavelengths.

Since the solvent system used did not give clean separations of d-AMP, d-CMP and TMP even after the chromatogram had been developed for 72 hours, the three nucleotides were eluted together and the solution analysed spectrophotometrically.

(c) Action of spleen phosphodiesterase on DNA hydrolysates (47)

The desalted DNA hydrolysates from III(1) were dissolved in 5 ml. 0.18M ammonium acetate buffer, 0.06M in EDTA, pH 5.7. One ml. of this solution was incubated with 0.2 ml. of spleen phosphodiesterase preparation for 16 hours at 37°C. One drop of chloroform was added as preservative. After the digestion, the reaction mixture was boiled for 5 minutes to inactivate the enzyme, and the protein precipitate removed by centrifugation at 16,000g for 60 min. at 0°C. The supernatant solution was then desalted by gel filtration (zinc uranyl acetate reagent for Na⁺). The effluent fractions from the Sephadex column which contained the oligonucleotides were pooled and evaporated to dryness. The residue was redissolved in 5 ml. 0.001M tris buffer, pH 8.0. Ten O.D. units of material were incubated with 10 μ l. of the dilute phosphomonoesterase preparation, one drop of chloroform being added as preservative. The reaction was allowed to proceed for 16 hours at 37°C, after which the enzyme protein was removed by boiling and centrifugation.

The supernatant solution was chromatographed on Whatman No. 40 filter paper. The system employed was water-saturated butanol-conc. ammonia, and the descending technique

was used. Standard solutions of nucleosides were also spotted onto the same chromatogram as markers. The chromatogram was developed for about 30 hours. The nucleoside spots were located with the aid of ultraviolet light. These were extracted with 0.1N HCl as described and quantitatively determined spectrophotometrically.

(d) Base Composition analysis

The various oligonucleotide fractions (IIIIi) isolated from a salmon sperm DNA hydrolysate prepared by exposing the DNA to the purified or the crude intestinal mucosal extracts were analysed for their base compositions. Each fraction was subjected successively to enzymatic degradation by venom phosphodiesterase, spleen phosphodiesterase and E. coli phosphomonoesterase by a combination of methods (b) and (c) above. The base composition of each fraction was determined by estimating the amount of each nucleoside present and converting these values to the corresponding bases.

Where the base composition of native deoxyribonucleic acids was desired, the complete desalted DNA hydrolysate obtained by treatment with commercial DNase I was used as substrates for venom phosphodiesterase and E. coli phosphatase.

V. Effect of Heat, pH, Mg^{++} and Inhibitors on the Rate of Digestion of DNA by the Rat Intestinal Mucosal Extract.

(1) Effect of heat

The intestinal mucosal extract was heated for 5 minutes at $100^{\circ}C$ in a boiling water bath. The solution was then centrifuged at 16,000g in the Servall refrigerated centrifuge for 60 min. to remove the protein precipitate. The supernatant solution was tested for deoxyribonuclease activity as described in II(1).

(ii) Effect of pH

Reagents:

0.1M acetic acid solutions, 0.01M in $MgCl_2$, were adjusted to the desired pH values by addition of NH_4OH . Salmon sperm DNA was dissolved in these ammonium acetate buffer solutions to a concentration of 200 $\mu g./ml.$

Procedure:

3 ml. aliquots of the buffered DNA solutions were preincubated at $37^{\circ}C$ in the thermostated cell chamber of the Cary spectrophotometer. At zero time, 0.1 ml. of enzyme solution was introduced. The initial increase in optical density at 260 $m\mu$ with time was used as a measure of enzyme activity.

(iii) Magnesium requirements of the intestinal extract

Rat intestinal mucosal DNA solution, 40 $\mu g./ml.$ in

0.1M ammonium acetate buffer, pH 6.5, was used as substrate. The crude and the purified intestinal mucosal extracts were dialysed against the acetate buffer to remove the magnesium ions. The resulting enzyme solutions were assayed for deoxyribonuclease activity by the spectrophotometric method of Kunitz as described.

(iv) Effect of Inhibitors

(a) Preparation of a specific DNase I inhibitor from the pigeon crop gland.

The existence of a specific DNase I inhibitor in the pigeon crop gland was first reported by Laskowski (48). The inhibitor was isolated as follows:-

Twelve pigeons were etherized, immediately decapitated and the blood drained from the wound. The crop gland was exposed, opened and the contents washed off. The inner lining of the gland was then peeled off and the tissue chopped up into very tiny pieces. The tissue was then homogenized with 9 times its own volume of water in the Servall Omni mixer with the rheostat setting on 30V. The homogenate was then centrifuged for 120 minutes at 16,000g in the Serval refrigerated centrifuge. The supernatant solution was collected, and since repeated thawing and freezing was found to abolish the activity of the inhibitor protein (48), the supernate was

stored at -20°C in 2 ml. fractions to be used separately.

- (b) Action of known DNase inhibitors on the activity of the intestinal enzyme.

Reagents:

- (a) EDTA solution: 0.1M in water, solubilized by the addition of a minimum amount of NaOH. When 0.1 ml. of this solution was added to 3 ml. of substrate DNA solution (40 μg DNA/ml. in 0.1M acetate buffer, pH 6.5), there was no significant rise in pH of the mixture.
- (b) 1M trisodium citrate solution
- (c) 1M disodium hydrogen arsenate solution
- (d) pigeon crop extract, as previously prepared
- (e) 1M MgCl_2

Procedure:

Three ml. of substrate DNA solution was warmed to 37°C in the cell chamber of the Cary spectrophotometer. At zero time, 0.1 ml. of the inhibitor solution was added, followed immediately by 0.1 ml. of enzyme solution. The change in optical density of the mixture with time was observed. Where inhibition was observed, the system was further charged with aliquots of enzyme preparation or Mg^{++} .

The effect of preincubating the substrate with the inhibitors was also studied. In these cases, the inhibitor

(0.1 ml.) was added to the substrate solution and the mixture incubated at 37°C for 30 minutes. 0.1 ml. of enzyme was then added and the deoxyribonuclease activity measured as before.

VI. Intracellular Distribution of Deoxyribonuclease in the Rat Intestinal Mucosa.

The tissue fractionation procedure was performed according to de Duve et al (49, 50).

Five male Wistar rats weighing 180-200 gm. each were sacrificed as usual and the small intestine removed. The lumen was flushed thoroughly with cold 0.25M sucrose and the mucosal scrapings prepared as described under I(i). The tissue was dispersed with approximately 3 volumes of chilled sucrose medium in a homogenizer of the Potter & Elvehjem type, consisting of a smooth-walled glass tube fitted with a teflon pestle. The tube, kept in cracked ice, was given an upward run against the rotating pestle (rheostat setting at 90V) until all the material had been forced above the latter. The resulting slurry was centrifuged at 1,000g for 10 min. at 0°C. The pellet was rehomogenized in about the same quantity of medium and centrifuged again at 1,000g for 10 min. This was repeated a second time and the nuclear sediment was redispersed by means of the homogenizer in 4 vols. of Krebs-Ringer phosphate buffer, pH 7.8 to give a 1:4 nuclear fraction.

The supernatant solutions from the three centrifugations were combined and made up with the sucrose medium to form a 1:10 cytoplasmic extract. This was designated the 10,000g-min. supernate in accordance with the scheme of de Duve (49). This cytoplasmic extract was next centrifuged at 3,300g for 10 minutes. The pellet was washed and resuspended in Krebs-Ringer phosphate buffer and the suspension dispersed by homogenization.

The supernatant solution from the last centrifugation was again centrifuged at 24,000g for 10 min. The sediment was collected, washed and redispersed with 2 vols. of sucrose medium by homogenization. The suspension was made to 0.3N with respect to formic acid and adjusted to pH 7.8 with NaOH for the following reason:

In the first fractionation experiment, all the subcellular fractions were redispersed with Krebs-Ringer phosphate buffer. When the enzyme activities were assayed, it was found that the 3,300g supernatant solution showed higher activities towards DNA at the more acidic pH (4.5). This seemed to indicate the presence of DNase II activity in the fraction. In the second experiment, the pellet obtained after centrifugation at 24,000g was therefore homogenized in 0.3M sodium formate to create optimum ionic conditions for

DNase II.

The three buffered homogenates together with the 24,000g-supernate were then incubated at 37°C. for three hours to release the enzyme into solution. They were next centrifuged at 105,000g for 60 minutes. The supernatant solutions were collected and assayed for enzyme activities at pH 4.5 and 6.5.

RESULTS AND DISCUSSION

I. Preparation, Purification and Characterization of Substrate Deoxyribonucleic Acids.

Integrity of substrate is essential for the study of an enzymatic reaction. Complex molecules such as deoxyribonucleic acids are especially susceptible to denaturation and degradation. Even mild mistreatments can lead to separation of the strands of the double helix and other states of degradation such as depurination and deamination of the nitrogenous bases. Therefore, in choosing the methods of preparation and purification, it is important that while the yield is to be as quantitative as possible, the degradation and impurities must be reduced to a minimum.

Several criteria are available for defining the state of integrity of deoxyribonucleic acids. The nitrogen and phosphorus contents of a DNA sample are reflections of the state of purity of the compound, while the atomic extinction coefficient with respect to phosphorus, designated $\epsilon(P)$, the viscosity and spectral data of the acid are valuable guides to its state of denaturation and degradation.

The detergent method of Colter et al (33) for the isolation of DNA from tissues, when applied to the rat intesti-

nal mucosa, yielded a white fibrous product resembling asbestos. Average of three experiments showed a yield of 34 mg. per 10 gm. wet tissue. Leslie (51) reported an average DNA content of rat intestinal mucosa as 129 mg. per 10 gm. fresh tissue. The method of Colter et al. for isolation therefore involves a very considerable loss of DNA. It has been reported that the use of stainless steel disintegrator caused degradation of DNA catalysed by traces of ferrous ions (52). However, the use of the Servall Omni mixer was necessary to ensure complete homogenization of the tissue which would minimize the contamination of DNA with cytoplasmic RNA and other materials.

Deproteinization of the salmon sperm DNA also afforded a product in the form of white, tough strands of fibres. The DNA yielded a highly viscous, slightly turbid solution when dissolved in acetate buffer, but the solution could be clarified by centrifugation.

It has been found that owing to the structural regularities in all deoxyribonucleic acids, the nitrogen and phosphorus contents are closely similar even for preparations showing large differences in base compositions. These values cannot be utilized for precise definition of composition, but they are useful as indications of the degree of purity of the samples.

In studying the action of intestinal deoxyribonuclease activity, DNA from three different tissues were used as substrates. These had been characterized and the analytical data are summarized in Table I. As can be seen, the N and P values and also the atomic N/P ratios for the rat intestinal mucosal and for the commercial calf thymus DNA are in good agreement with values reported in literature for DNA preparations from tissues of various species (53). The salmon sperm preparation showed too high a value for nitrogen and too low a value for phosphorus. This is interpreted as indicating protein contamination.

The $\epsilon(P)$ values for the rat DNA and the calf thymus DNA are also within the normal range observed with cautiously isolated DNA's from various sources (6,500-7,000). According to Chargaff (53), $\epsilon(P)$ values higher than 7,200 are indicative of denaturation. The $\epsilon(P)$ value for the deproteinized salmon sperm DNA preparation of 7,310 would be a sign of degradation. However, as pointed out earlier, the salmon sperm DNA appears to be contaminated with protein, which shows a certain amount of residual absorption at 260 m μ and consequently giving rise to a higher atomic extinction. In the light of these data, all the three DNA preparations examined were considered to be free of extensive denaturation and degradation.

TABLE I
Elementary Composition and Some Other Properties
of the DNA Preparations Used

Source	Nitrogen %	Phosphorus %	Atomic N/P ratio	$\epsilon(P)$	Relative viscosity with respect to solvent (1 mg./ml.)	Base Composition			
						A	C	G	T
rat intestinal mucosa	14.8	9.1	3.64	6770	2.51	28.6	20.3	23.3	27.8
salmon sperm	16.8	7.4	5.03	7310	3.03	27.7	22.3	23.7	26.3
calf thymus	15.2	9.2	3.67	6970	3.87	28.6	19.8	21.3	30.3

The relative viscosities of the DNA solutions were found to vary from batch to batch. Those reported in Table I are averages of values that were observed most consistently. The commercial calf thymus preparation was designated "highly polymerized" by the manufacturer and, for the purpose of comparison, was used as a standard. The relative viscosity values for the rat intestinal and the salmon sperm preparations were considered satisfactory.

While it was impossible to define an absolute standard of integrity for any DNA sample, the three preparations examined were considered to be relatively free of denaturation and degradation and were deemed satisfactory for use as substrates in enzymatic studies.

II. Preparation and Purification of Deoxyribonuclease from Rat Intestinal Mucosa.

(1) Effect of Extracting Media

It has been found that deoxyribonuclease is extremely susceptible to destruction by the action of proteolytic enzymes (54). Thus, a fresh extract of pancreatic deoxyribonuclease was found to lose its activity rapidly when the proteolytic enzymes became activated. Moreover, a DNase I inhibitor had been found in many tissues (e.g. kidney) and DNase activity was only revealed after the inhibitor was

destroyed (55). A variety of solvents was tested in an attempt to find one most suitable for extraction of the enzyme. Two of these were found to be promising: Krebs-Ringer phosphate buffer and physiological saline. Figure 6 compares the deoxyribonuclease activities in the two extracts and Table II summarizes their protein content and the specific activity.

Krebs-Ringer phosphate buffer was the first medium tested. It yielded an extract which contained considerable deoxyribonuclease activity. This activity was stable for several months when stored at -20°C and it was also not affected after repeated thawing and freezing. When physiological saline was used as the extracting medium, it was found that the specific activity of this extract was a little bit higher than that of the Krebs-Ringer phosphate buffer, indicating that the buffered enzyme preparation contained more extraneous protein. However, the deoxyribonuclease activity in the physiological saline extract diminished with storage, probably due to denaturation of the enzyme protein in the unbuffered medium. Physiological saline was therefore deemed unsuitable as an extracting solution.

When a 1M NaCl solution was used for extraction, it was found that addition of the enzyme extract brought about the precipitation of a white fibrous material out of solution.

FIGURE 6 : Comparison of Rate of Degradation of DNA by Crude Extracts of Rat Intestinal Mucosa.

TABLE II: Specific Activity of Deoxyribonuclease in Crude Extracts of Rat Intestinal Mucosa.

Extracting Medium	Optical density of protein solution (280 mμ)	Conc. of protein ^(a) (mg./ml.)	DNase activity ΔO.D./min. (260 mμ)	Specific activity of extract Units ^(b) /mg. protein
Krebs Ringer phosphate buffer, pH 7.8	11.55	9.6	0.00467	0.00048
Normal Saline	7.63	6.3	0.0075	0.0012

(a) O.D. (280 mμ) of 1.2 = 1 mg. protein per ml. (20)

(b) 1 unit of DNase activity is the amount of enzyme capable of bringing about an optical density change of 1.0 at 260 mμ in one minute under the experimental conditions as defined in the text.

This precipitate was analysed qualitatively and found to contain both phosphorus and deoxyribose (diphenylamine reaction). The rest of the reaction mixture, when analysed spectrophotometrically, showed no absorption peak in the ultraviolet region except for a slight shoulder at 280 $m\mu$. It has been found by several authors that DNase I is inhibited when the sodium chloride concentration is increased above 0.05M (20,56), and 0.5M NaCl in the medium inhibited about 65%. It was therefore considered that 1M NaCl did not constitute a good extracting solvent for deoxyribonuclease, at least for the intestinal mucosa.

In one experiment, the tissue after a preliminary extraction with Krebs-Ringer phosphate buffer, was homogenized and re-extracted with fresh buffer. This second extract was also found to contain deoxyribonuclease activity which was stable to storage. This experiment showed that at least a part of the deoxyribonuclease activity in the rat intestinal mucosa was located intracellularly.

(ii) Fractionation with Ammonium Sulfate

In Table III, the data for the purification of the intestinal mucosal DNase (Krebs-Ringer phosphate buffer extract) are presented.

In the Table, the symbol 5%S is used to designate the

TABLE III

Purification of Intestinal Mucosal
Deoxyribonuclease Extract by
Fractionation with
Ammonium Sulfate.

Protein Fraction	Optical Density of Protein Solution (280 mμ)	Concentration of protein (a) mg./ml.	Activity O.D./min. x 10 ³	Specific Activity units/mg protein x 10 ³
Total Extract	11.55	9.6	4.67	0.48
5% S (b)	4.21	3.5	4.56	1.3
10% S	1.72	1.43	4.10	2.9
15% S	0.95	0.79	4.08	5.2
20% S	0.65	0.54	1.63	3.1
25% S	0.27	0.22	1.33	6.0
30% S	0.20	0.17	0.53	3.1
5% P (c)	7.44	6.2	0	0
10% P	2.15	1.8	0	0
15% P	0.82	0.68	0	0
20% P	0.362	0.30	3.0	10
25% P	0.11	0.09	0.37	2.9
30% P	0.08	0.07	0.20	0.48

- a. Optical density of 1.2 (280 mμ) = 1 mg. protein/ml.(20)
- b. 5% S = supernatant fraction at 5% saturation of ammonium sulfate.
- c. 5% P = solution obtained by redissolving the protein precipitated at 5% saturation of ammonium sulfate.

supernatant solution obtained after centrifuging off the precipitate brought down at an ammonium sulfate saturation of 5%. The precipitate was redissolved in phosphate buffer and the resulting solution designated "5%P".

Nitrogen determination of the various protein solutions was found to give erratic results, presumably due to incomplete removal of the ammonium salt by dialysis. The protein concentration in each solution was therefore approximated spectrophotometrically at 280 m μ , the optical density being about 1.2 per mg. of protein per ml. (20).

It can be seen from Table III that the deoxyribonuclease activity was brought down separately at two concentrations of ammonium sulfate during fractionation. The first portion was precipitated at 20% saturation, the other at 30%. The 20%P fraction was found to contain 64% of the activity in the whole extract. Whereas the 25%S fraction still retained 28% of the total activity, the 30%S contained only about 10%. However, deoxyribonuclease activity was not observed in the 30%P fraction (less than 5% of the total activity in the crude extract). In a repeat experiment, similar findings were observed. The deoxyribonuclease activity brought down at 30% saturation of ammonium sulfate was lost. All attempts to recover the enzymatic activity in this fraction, such as reconstitution with the supernate and dialysis to remove excess

salts were unsuccessful. The reason for the loss of activity is not clear; it is possible that the high salt concentration might have irreversibly denatured the enzyme. Later experiments were therefore carried out using both the crude and the partially purified enzyme preparations and comparisons between the two were made.

It can be seen from the specific activity of the 20%P fraction that only a modest 20 fold purification of the enzyme was achieved by ammonium sulfate fractionation.

The observed ammonium sulfate concentration at which deoxyribonuclease activities were precipitated was different from that for pancreatic DNase I reported by Kunitz (57). Kunitz observed that the bulk of the DNase I activity from beef pancreas was precipitated at 30% saturation. The difference was not surprising since the extraction of the pancreatic enzyme was performed under extremely acidic conditions (0.25N H_2SO_4) to inhibit proteolytic activities. Moreover, the high concentration of divalent ions (which were found to be capable of binding DNA)(58) in the Krebs-Ringer phosphate buffer might have contributed towards the lesser solubility of the enzyme in the ammonium sulfate solution.

III. Enzymatic Degradation of Deoxyribonucleic Acids and Analysis of the Products.

(i) Kinetics of Reaction

The degradation of deoxyribonucleic acids by the rat intestinal enzyme preparations, whether measured spectrophotometrically or by viscosity determinations, was characterized by an initial constant reaction rate which gradually decreased after a period of time. Two typical examples were illustrated in Figures 7 and 8.

It can be seen that the rate of increase of ultraviolet light absorption or the rate of decrease in relative viscosity of the reaction mixture vary linearly with time at the start of the reaction. During this period, when the substrate concentration was high, the reaction was of zero-order. When the substrate concentration was reduced, as when the reaction had been permitted to proceed for a period of time, the reaction approximated one that was first-order in substrate concentration, as can be seen from the gradually decreasing rate of reaction.

The rate of change in viscosity of the solution is also seen to be much more rapid than the rate of change in the absorption of ultraviolet light. This is interpreted as indicating that the intestinal DNase is of the endonuclease type, chopping the DNA molecules into large fragments still containing considerable double helical structure. Similar observations had been made by Kunitz (20) for DNase I and by Jungner et al (59).

FIGURE 7 : Time Course of the Degradation of DNA by Crude Intestinal DNase Extract as Measured by the Spectrophotometric Method of Kunitz.

FIGURE 8 : Time Course of the Degradation of DNA by Crude Intestinal DNase Extract as Measured by Viscosimetry.

(ii) Ion-exchange Chromatography and Characterization
of the Products in DNA Hydrolysates.

Chromatographic separations were conducted with DEAE-cellulose columns (chloride form). To effect good resolution of the oligonucleotides, 7M urea was incorporated into the eluant to remove the secondary binding forces which existed between the ion-exchanger and the purine and pyrimidine derivatives (41). The results of these studies are shown in Figures 9 to 14.

DEAE-cellulose, being an anion exchanger, separates oligonucleotides according to the number of net negative charges they carry, ie. according to the degree of polymerization, the smaller nucleotides being eluted first. As can be seen, each elution profile contains 7 distinguishable peaks. Peaks I, II and III are usually fairly well separated from each other, but the peaks containing the higher oligonucleotides show considerable overlap.

When the digestion was performed with the crude enzyme extract (Figures 9-11) it was seen that the size of a given peak (corresponding roughly to the amount of oligonucleotides present) varies considerably from one DNA to another. Thus peak VII obtained for calf thymus DNA is definitely much smaller than the corresponding peaks for the rat intestinal or the salmon sperm DNA. On the other hand, when the partially

FIGURE 9 : Ion-exchange Chromatography of the Oligonucleotides in Rat Intestinal Mucosal DNA Hydrolysate Formed by Exposing the DNA to a Crude Intestinal DNase Preparation. The Chromatography was Done on a DEAE-cellulose Column Developed with a Linear Gradient of NaCl to 1M.

FIGURE 10 : Ion-exchange Chromatography of the Oligonucleotides in Salmon Sperm DNA Hydrolysate Formed by Exposing the DNA to a Crude Intestinal DNase Preparation. The Chromatography was Done on a DEAE-cellulose Column Developed with a Linear Gradient of NaCl to 1M.

FIGURE 11 : Ion-exchange Chromatography of the Oligonucleotides in Calf Thymus DNA Hydrolysate Formed by Exposing the DNA to a Crude Intestinal DNase Preparation. The Chromatography was Done on a DEAE-cellulose Column Developed with a Linear Gradient of NaCl to 1M.

FIGURE 12 : Ion-exchange Chromatography of the Oligonucleotides in Rat Intestinal Mucosal DNA Hydrolysate Formed by Exposing the DNA to the 20%P Enzyme. The Chromatography was Done on a DEAE-cellulose Column Developed with a Linear Gradient of NaCl to 1M.

FIGURE 13 : Ion-exchange Chromatography of the Oligonucleotides in Salmon Sperm DNA Hydrolysate Formed by Exposing the DNA to the 20%P Enzyme. The Chromatography was Done on a DEAE-cellulose Column Developed with a Linear Gradient of NaCl to 1M.

FIGURE 14 : Ion-exchange Chromatography of the Oligonucleotides in Calf Thymus DNA Hydrolysate Formed by Exposing the DNA to the 20%P Enzyme. The Chromatography was Done on a DEAE-cellulose Column Developed with a Linear Gradient of NaCl to 1M.

purified enzyme was used, the pattern of peak size and distribution is remarkably constant from one species of DNA to another. The significance of these are discussed later.

The peaks were characterized by the two criteria: (a) the ratio of phosphomonoesterase-sensitive phosphorus to total phosphorus and (b) the decreasing $\xi(P)$ ratio with increasing size. The base composition of peaks I and VII obtained from the digests of salmon sperm DNA were analysed. The amount of nucleotide present in the various peaks was also determined from its phosphorus content. Table IV summarizes these results.

The $\xi(P)$ values and the ratios of total phosphorus to phosphomonoesterase-sensitive phosphorus indicated that the degradation products of deoxyribonucleic acids were indeed eluted according to their sizes. The mononucleotides were eluted first (peak I) followed by the di-, tri-, etc. nucleotides. The overlapping and coincidence of the peaks containing the higher oligonucleotides invariably caused some contamination and the total P/PME-P ratios obtained of these peaks were considerably removed from integral values.

When the percentage of a particular oligonucleotide was examined, some interesting results were observed. The crude enzyme extract invariably gave rise to more mononucleotides and large oligonucleotides (degree of polymerization > 7),

TABLE IV

Some Analytical Data of DNA Hydrolysates

DNA	Enzyme Preparation Used	Peak	$\epsilon(P)$	Total P	Degree of Polymerization	% of oligonucleotide phosphorus	Base Composition			
				$\overline{PMEP^{(a)}}$			A	C	G	T
Salmon	Crude	I	10630	1.2	1	1.2	14.2	8.2	7.9	69.7
		II	9020	2.1	2	3.5	-	-	-	-
		III	8530	3.3	3	9.4	-	-	-	-
		IV	8200	4.2	4	10.2	-	-	-	-
		V	7850	5.2	5	7.6	-	-	-	-
		VI	7610	6.6	6	7.4	-	-	-	-
		VII	7460	7.4	7	61.7	28.2	21.9	23.2	26.7
Sperm	20% P	I	9890	1.1	1	1.04	10.5	10.4	4.9	74.2
		II	9130	2.2	2	4.1	-	-	-	-
		III	8640	3.1	3	10.9	-	-	-	-
		IV	8120	4.3	4	7.2	-	-	-	-
		V	7780	5.4	5	10.8	-	-	-	-
		VI	7510	6.2	6	8.5	-	-	-	-
		VII	7380	7.2	7	57.5	29.8	4.2	35.9	30.1
Rat Intestinal Mucosal DNA	Crude	I	9940	1.2	1	5.8				
		II	9180	2.2	2	4.1				
		III	8540	3.2	3	8.3				
		IV	7990	4.5	4	7.6				
		V	7570	5.0	5	7.0				
		VI	7160	6.5	6	7.6				
		VII	6710	7.5	7	59.6				

TABLE IV (Continued)

DNA	Enzyme Preparation Used	Peak	$\zeta(P)$	PM -P	Degree of Polymeri- zation	% of oligo- nucleotide phosphorus
Rat Intestinal Mucosal DNA	20% P	I	9950	1.0	1	2.4
		II	9230	2.1	2	5.2
		III	8630	3.2	3	11.2
		IV	8140	4.2	4	8.4
		V	7680	5.3	5	10.1
		VI	7260	6.3	6	10.0
		VII	6910	7.4	7	52.7
Calf	Crude	I	10230	1.1	1	10.55
		II	9240	2.4	2	7.0
		III	8510	2.9	3	15.6
		IV	8030	4.3	4	12.5
		V	7780	5.7	5	12.0
		VI	7510	6.3	6	11.6
		VII	7050	7.6	7	31.75
Thymus	20% P	I	9970	1.1	1	2.2
		II	9410	2.1	2	3.8
		III	8900	3.1	3	9.5
		IV	8430	4.3	4	15.4
		V	7820	5.2	5	9.6
		VI	7330	6.4	6	10.2
		VII	7110	7.4	7	48.3

(a) PME-P = Phosphorus liberated by phosphomonoesterase.

but fewer dinucleotides than the purified extract, regardless of the source of the DNA preparation. This is interpreted to mean that while the purified extract contained one DNase, the crude extract might contain more than one enzyme possessing deoxyribonuclease activity. During fractionation studies, the intestinal mucosal enzyme had been observed to precipitate out at two different salt concentrations. This together with the fact that the elution profiles of crude-enzyme digests of DNA lack any uniformity from one species of DNA to another were early clues already pointing to the existence of more than one deoxyribonuclease in the mucosal tissue.

When these percentage values were analysed in the light of the base composition of the various DNA samples (see Table I), it was found that the higher the A-T content in the DNA, the more mononucleotides were formed. Thus with salmon sperm DNA, which contains 54% A-T, the mononucleotide fraction constituted only about 1% of the total oligonucleotide. With the rat intestinal DNA, (A-T content 56.4%), the mononucleotide fraction amounted to 2.2% of the total oligonucleotide when the purified enzyme was used for digestion, and to 5.8% when the crude enzyme preparation was used. These observations seem to indicate that the intestinal deoxyribonuclease, or at least the "20%P" enzyme, shows a preference towards a phosphodiester linkage next to a thymine base.

To further clarify this point, the base composition of the mononucleotide peak and also of peak VII was analysed, and it was found that thymidylic acid constituted approximately 70% of the total mononucleotides. The high oligonucleotide fraction showed no unusual base frequency features when the crude enzyme was used, except perhaps a slight preponderance of purines. But in the "20%P" digest, the concentration of d-cytidylic acid was extremely low. This fraction was also found to be rich in purines. One possible explanation for these findings is that the presence of pyrimidines is required for hydrolysis by the "20%P" enzyme, and long chains of purines are consequently resistant to attack.

Schmidt (60) reported that in DNase I digests of calf-thymus DNA, thymidylic acid counted for 50-60% of the total mononucleotides formed. No base preference has been observed thus far for DNase II. In view of the above findings, it seems likely that the "20%P" enzyme is DNase I.

(iii) Secondary Degradation of DNA Hydrolysates
by phosphodiesterases.

(1) Venom phosphodiesterase

The rate of degradation of salmon sperm DNA hydrolysates by venom phosphodiesterase was determined by measuring (i) the deoxyguanylic acid and (ii) the composite dAMP-d-CMP-TMP

fraction formed by venom phosphodiesterase. The deoxyribonucleases used were the crude enzyme preparations, the 20%P fraction and commercial DNase I. Results of these studies are shown in Figure 15.

It has been reported (61) that oligonucleotides bearing 5'-phosphomonoester groups are hydrolysed by venom phosphodiesterase much more rapidly than the corresponding members lacking such groups. The rate of release of mononucleotides by venom phosphodiesterase from a DNA hydrolysate can therefore serve to indicate the relative abundance of 5'-phosphomonoester linkages in the products. Since DNase I gives rise exclusively to oligonucleotides bearing only 5'-PO₄ groups (62), a DNase I digest of DNA was used as a standard of comparison. As can be seen from Figure 16, the rates of release of both d-GMP and (dAMP-dCMP-TMP) from the "20%P" digest were comparable to those from a DNase I digest, while the crude enzyme digest was much more resistant to attack by venom phosphodiesterase. These findings are interpreted to mean that the 20%P digest contains much more oligonucleotides terminating in 5'-phosphate ends than the crude enzyme extract. Moreover, when the purified enzyme digest was used, the presence of undegraded oligonucleotide (presumably bearing 3'-PO₄ ends) was not detected on the chromatogram after the secondary digestion.

FIGURE 15 : Rate of Release of Mononucleotides by Snake Venom Phosphodiesterase from Salmon Sperm DNA Hydrolysate. The Optical Density of the d-GMP Solution was Measured at 245 mμ, that of (dAMP+d-CMP+TMP) at 267 mμ.

FIGURE 16 : Effect of pH on Deoxyribonuclease Activity of Rat Intestinal Mucosal Extracts.

had proceeded for over 60 minutes. These findings indicated the absence of any detectable amount of oligonucleotides bearing 3'-phosphomonoester groups in the DNA hydrolysate formed by the 20%P enzyme. This is in agreement with the previous proposition that the 20%P enzyme is DNase I, whereas the crude enzyme extract, giving rise to considerable amounts of oligonucleotides lacking a phosphate group at the 5'-position, also contains DNase II.

(2) Spleen phosphodiesterase

The amount of nucleosides obtained from the successive secondary degradations of salmon sperm DNA hydrolysate by spleen phosphodiesterase and E. Coli phosphomonoesterase are presented in Table V.

TABLE V: Nucleoside Content of mixture obtained by successive degradation of salmon sperm DNA with DNase, spleen phosphodiesterase and E. Coli phosphomonoesterase.

DNase Enzyme Used	Nucleoside* μ mole			
	adenosine	cytidine	guanosine	thymidine
DNase II	28.9	24.3	25.1	29.4
Crude intestinal prep.	9.2	10.5	7.3	8.7
20%P	0	0	0	0.4

* Ten O.D. units of the DNase-spleen phosphodiesterase digest used for degradation by phosphomonoesterase.

Spleen phosphomonoesterase degrades only those oligonucleotides which carry a 5'-hydroxyl group (47). From Table V, it can be seen that about 30% of the oligonucleotides present in the DNA hydrolysate by the crude enzyme preparation did not terminate in a 5'-phosphomonoester group and was therefore vulnerable to spleen phosphodiesterase. The presence of this type of oligonucleotides can only be explained by assuming that the crude intestinal extract contains DNase II, which is known to produce products ending in 3'-phosphate groups (31). The 20%P enzyme did not yield any detectable amount of this type of oligonucleotides and did not therefore contain any DNase activity. That some thymidine was observed when the 20%P enzyme was used for degrading the DNA is probably due to the presence of some thymidylic acid in the DNase hydrolysate.

IV. Effect of Heat, pH, Mg^{++} and some Inhibitors on the Rat Intestinal Deoxyribonuclease.

The intestinal deoxyribonuclease activity in the rat intestinal mucosa was found to be heat-labile. Heating for 5 minutes at 100°C completely abolished the activity of the extract.

Both the crude and the 20%P enzyme preparations showed pH optima at about 6. Figure 16 shows the results obtained. Activities at various pH are expressed in terms of

the slopes of the straight lines obtained in plotting increase in optical density versus time.

The 20%P enzyme was found to show an absolute requirement for Mg^{++} . When the enzyme solution was dialysed against buffer to remove the Mg^{++} originally present in the medium, it lost its activity. But activity was restored by adding Mg^{++} to the reaction mixture. Anions such as citrate, arsenate and EDTA which remove magnesium were all found to inhibit DNase activity. EDTA was particularly effective. A concentration of 1.67×10^{-4} M inhibited 70% when the Mg^{++} concentration was 1×10^{-2} M. When EDTA was preincubated with the substrate DNA solution (containing Mg^{++}), the inhibitory action was even more pronounced, 1.67×10^{-4} M EDTA completely inhibiting the enzyme. Doubling the Mg^{++} resulted in a slight recovery of enzyme activity to about 10% of that of the original extract.

The inhibitory power of citrate and arsenate was not as pronounced as that of EDTA, but still considerable. In either case, preincubation with DNA substrate did not enhance the inhibitory power. The effect of these two inhibitors was also decreased by increasing the concentration of Mg^{++} .

The pigeon crop extract was found to be an extremely potent inhibitor of the 20%P enzyme. Its effect was not reversed by increasing Mg^{++} .

When the crude enzyme extract was similarly tested, it was found that dialysis to remove Mg^{++} decreased the deoxyribonuclease activity to about 20%. This residual activity was not diminished by further dialysis, indicating that it did not require Mg^{++} for activity. Arsenate, citrate and EDTA also did not affect this activity, in accordance with the findings of Stewart and Zbarsky (32).

These studies showed that the DNase activity in the 20%P fraction resembles the pancreatic DNase I in its pH optimum and ionic requirements (38, 57). They also served to demonstrate that the crude enzyme extract contains in addition a different DNase which is independent of Mg^{++} and unaffected by citrate, arsenate etc. and resembles DNase II in these respects.

V. Intracellular Distribution of Deoxyribonucleases in the Rat Intestinal Mucosa.

Earlier evidence obtained in the course of this research favoured the idea that the rat intestinal mucosa enzyme was DNase I in nature. Since DNase I is abundant in pancreatic juice, it was feared that the enzyme activity isolated might be due to contamination, both from adsorbed pancreatic enzyme and from intestinal bacteria. It was therefore deemed desirable to establish the intracellular location and distribution of the enzyme activities in the subcellular fractions.

The intracellular location of the enzyme was indicated by the fact that after a first extraction by buffer of a whole cell preparation was made, a second extraction done on the homogenized tissue yielded a preparation only slightly less active than the whole cell extract.

When tissue fractionation was next performed on the intestinal mucosa, 3 fractions were obtained: a nuclear fraction (10,000g-min. precipitate, in accordance with the terminology of de Duve et al.)(49), the 33,000g-min. pellet and the 33,000g supernatant solution. It was found that 48% of the total DNase activity was located in the 33,000g-min. pellet (mitochondrial); 38.7% in the supernate and 3.3% in the nuclear fraction. Moreover, the enzyme present in the 33,000g-min. supernatant solution was found to be more active at pH 4.5 than at pH 6.0.

The fractionation was repeated and the 33,000g-min. supernatant solution was further separated into a fraction termed "light mitochondrial" fraction and a final supernatant solution by centrifugation at 24,000g for 10 minutes. The enzyme activities of these fractions are summarized in Table VI.

The recovery of activity during the fractionation was very satisfactory (94.6%). As in the previous experiment, the "light mitochondrial fraction" was found to be more active

TABLE VI: Intracellular distribution of deoxyribonuclease in rat intestinal mucosa.

Fraction	% Activity
Total Extract	100
Nuclear	4.3
Heavy mitochondrial	48.1
Light mitochondrial	38.4
Cytoplasmic	3.8

at pH 4.5 than 6.0. This seems to indicate DNase II.

However, the products of DNA digestion by this enzyme fraction had not been characterized.

Oth et al. (63) reported that in the normal liver cell, both DNase I and DNase II were associated mainly with mitochondria. De Duve et al. (49) found that DNase I occurred in liver cell mitochondria, but DNase II was associated with the lysosomes. Lysosomes are particles intermediate in size between mitochondria and microsomes and would be expected to be centrifuged down at 24,000g for 10 minutes. In the light of this, it seems very likely that the deoxyribonuclease activity in the "light mitochondrial" fraction of the rat intestinal mucosa is of the DNase II type.

VI. General Discussion

In summarizing the studies carried out for the rat intestinal mucosa deoxyribonuclease, it may be said that two types of enzyme activities are present. One protein could be salted out at 20% ammonium sulfate and its activity recovered by re-dissolution in buffer. It showed a pH optimum of 6.0, required Mg^{++} for activities and was inhibited by EDTA, citrate, arsenate and a protein DNase I inhibitor isolated from the pigeon crop. The products of reaction terminated in 5'-phosphomonoesters. It therefore fell into the category of enzymes termed DNase I.

The other protein was precipitable at 30% saturation ammonium sulfate. But since its activity was not recovered by redissolution in buffer, it could only be studied in the crude extract which also contained the first kind of protein. However, this protein could be shown to differ from the first in ionic requirements and nature of products formed, and to resemble DNase II. Other circumstantial evidences such as the intracellular distribution of deoxyribonuclease activity and oligonucleotide frequencies in DNA hydrolysates by the crude and 20%P enzymes also pointed to the DNase II nature of the second protein.

SUMMARY

1. The deoxyribonuclease activity in the intestinal mucosa of the rat was studied.
2. This activity could be extracted from a whole cell preparation of the mucosa with either Krebs-Ringer phosphate buffer or physiological saline. The extract obtained using the latter solvent as extracting medium was however found to be the less stable of the two.
3. The deoxyribonuclease activity was found to precipitate from solution at two different salt concentrations when the enzyme proteins were fractionated with ammonium sulfate. One active protein was precipitated at 20% saturation of $(\text{NH}_4)_2\text{SO}_4$ and its activity could be recovered by redissolving it in Krebs-Ringer phosphate buffer. This material was designated as the 20%P fraction, and represented a 20 fold purification of the crude extract.

The supernatant fraction obtained after this protein was removed by centrifugation still contained DNase activity. This residual activity precipitated from solution when the concentration of ammonium sulfate was increased to 30% saturation. However, all attempts to recover DNase activity from the precipitated protein were unsuccessful, studies of the residual deoxyribonuclease activity were

carried out indirectly in the crude extract.

4. The 20%P enzyme was found to be thermolabile. It showed an optimum pH of 6, and was found to require magnesium ions for activity. It was strongly inhibited by EDTA, citrate, arsenate and a DNase I-specific protein inhibitor isolated from the pigeon crop gland. Ion-exchange chromatography on DEAE-cellulose (chloride form) columns established that the products of reaction vary from mononucleotides through to oligonucleotides with a degree of polymerization larger than 7. These products were shown to terminate in 5'-phosphates.

The base frequencies differed in different oligonucleotides. The mononucleotide fraction contained about 70% thymidylic acid and the amount of deoxycytidylic acid in the higher oligonucleotides was shown to be very low. This was interpreted to mean that phosphodiester linkages next to a pyrimidine base were preferentially attacked by the enzyme.

5. In the crude extract, the residual deoxyribonuclease activity was shown to be qualitatively different from the 20%P activity. It did not require activation by Mg^{++} , and was not inhibited by EDTA, citrate or arsenate. Some of the products formed by the crude enzyme extract were shown to carry a monoesterified phosphoryl group at

carbon 3'. These products were not formed by the 20%P enzyme and therefore must be due to the residual deoxyribonuclease.

6. In light of the above findings, the 20%P enzyme was classified as DNase I whereas the residual deoxyribonuclease likely was of the DNase II type.
7. Studies on the intracellular distribution of deoxyribonuclease showed that the intestinal DNase I occurred in mitochondria. DNase II was associated with lighter subcellular particles.

BIBLIOGRAPHY

1. V. Allfrey and A.E. Mirsky, J. Gen. Physiol. 36, 227(1952).
2. M. Meselson and J.J. Weigle, P.N.A.S. 47, 858 (1961).
3. L.A. Heppel and P.R. Whitfield, B.J. 60, 1, 8 (1955).
4. C.C. Richardson, C.L. Schildkraut, H.V. Aposliran, A. Kornberg, W. Bodmer and J. Lederberg, in "Informational Macromolecules", (H. Vogel, ed.) p. 13, Acad. Press, New York, 1963.
5. C.L. Schildkraut, J. Marmur and P. Doty, J. Mol. Biol., 3 595 (1961).
6. I.R. Lehman, G.G. Roussos and E.A. Pratt, J.B.C. 237, 819 (1962).
7. D. Elson, B.B.A. 27, 216 (1958).
8. S. Ochoa, Arch. Biochem. Biophys. 69, 119 (1957).
9. R.W. Holley, J. Apgar, G.A. Everett, J.T. Madison, M. Marquisee, S.H. Merrill, J.R. Penswick and A. Zamir, Science 147, 1462 (1965).
10. J.B. Hall and R.L. Sinsheimer, J. Mol. Biol. 6 115 (1963).
11. T. Araki, Z. Physiol. Chem. 38, 84 (1903).
12. L. Iwanoff, Z. Physiol. Chem. 39, 31 (1903).
13. F. Sachs, Z. Physiol. Chem. 46, 337 (1905).
14. E. Abderhalden and A. Schittenhelm, Z. Physiol. Chem. 47, 452 (1906).
15. R. Feulgen, Z. Physiol. Chem., 237, 261 (1935).
16. G. Schmidt, E.G. Pickels and P.A. Levene, J.B.C. 127, 251 (1939).
17. M. Kunitz, J. Gen. Physiol. 24, 15 (1940).
18. M. Laskowski, Arch. Biochem. 11, 41 (1946).

19. M. Kunitz, J. Gen. Physiol. 32, 39 (1948).
20. M. Kunitz, J. Gen. Physiol. 33, 349 (1950).
21. J. Erkama and P. Suutarinen, Acta. Chem. Scand. 13, 323 (1959).
22. D.G. Catcheside and B. Holmes, Symp. Soc. Exptl. Biol. 1, 225 (1947).
23. M.E. Maver and A.E. Greco, J.B.C. 181, 853, 861 (1949).
24. L. Cunningham and M. Laskowski, B.B.A. 11, 590 (1953).
25. M. Laskowski, Ann. N.Y. Acad. Sci., 81, 776 (1959).
26. M. Laskowski, in Vol. 5, "The Enzymes", (P.D. Boyer, H. Lardy and K. Myrback eds.) p. 123, Acad. Press, N.Y. 1961.
27. R.O. Hurst and G.C. Becking, Can. J. of Biochem. and Physiol., 41, 469 (1963).
28. R.O. Hurst and G.C. Becking, Can. J. of Biochem. and Physiol., 41, 1433 (1963).
29. R.L. Sinsheimer, J.B.C., 215, 579 (1955).
30. J.L. Potter, U.-R. Laurila and M. Laskowski, J.B.C., 233, 915 (1958).
31. J.F. Koerner and R.L. Sinsheimer, J.B.C. 228, 1039, 1049 (1957).
32. B.E. Stewart and S.H. Zbarsky, Can. J. of Biochem. and Physiol., 41, 1483 (1963).
33. J.S. Colter, R.A. Brown and K.A.O. Ellem, B.B.A. 55, 31 (1962).
34. G. Zubay, Biochem. et Biophys. Acta 33, 244 (1959).
35. F. Pregl, in "Quantitative Organic Microanalysis", 5th edition, Julius Grant, London (1951).

36. C.H. Fiske and Y. SubbaRow, J.B.C. 66, 375 (1925).
37. E. Chargaff and S. Zamenhof, J.B.C. 173, 327 (1948).
38. M. McCarty, J. Gen. Physiol. 29, 123 (1946).
39. "Manometric Techniques", (W.W. Umbreit, R.H. Burris, and J.F. Stauffer, eds.), 4th ed., p. 132, Burgess Publ. Co., (1964).
40. M. Dixon, B.J. 54, 457 (1953).
41. R.V. Tomlinson and G.M. Tener, Biochem. 2, 697 (1963).
42. F.Felix, J.L. Potter and M. Laskowski, J.B.C. 235, 2105 (1959).
43. C. Mezei, Ph.D. thesis, U. of Brit. Col., Vancouver (1964).
44. W.E. Razzell and H.G. Khorana, J.B.C., 234, 2105 (1959).
45. E.J. King, Biochem. J. 26, 292 (1932).
46. J. Porath and P. Flodin, Nature 183, 1657 (1959).
47. W.E. Razzell and H.G. Khorana, J.B.C. 236, 1144 (1961).
48. W. Dabrowska, E.J. Cooper and M. Laskowski, J.B.C. 177, 991 (1949).
49. C. de Duve, B.C. Pressman, R. Gianetto, R. Wattiaux and F. Appelmans, Biochem. J. 60, 604 (1955).
50. F. Appelmans, R. Wattiaux and C. de Duve, Biochem. J. 59, 438 (1955).
51. I. Leslie, in Vol. II "The Nucleic Acids" (E. Chargaff and J.N. Davidson eds.) pg. 8, Acad. Press, N.Y. (1955).
52. S. Zamenhof in Vol. VI "Biochemical Preparations", John Wiley & Sons Inc., N.Y. (1958).
53. E. Chargaff, in Vol. I "The Nucleic Acids", (E. Chargaff and J.N. Davidson eds.) Acad. Press, N.Y. (1955).

54. M. Kunitz and J.H. Northrop, J. Gen. Physiol. 18, 433 (1935); 19, 991 (1936).
55. L. Cunningham and M. Laskowski, B.B.A. 11, 590 (1953).
56. J.P. Greenstein, C.E. Carter and W.H. Chalkley, Cold Spring Harbor Symp. Quant. Biol. 12, 64 (1947).
57. M. Kunitz, J. Gen. Physiol. 33, 363 (1950).
58. J. Erkama and P. Suutarinen, Acta Chem. Scand. 13, 323 (1959).
59. G. Jungner, I. Jungner and L.G. Allgen, Nature 164, 1009 (1949).
60. G. Schmidt, in Vol. I, "The Nucleic Acids", (E. Chargaff and J.N. Davidson, eds.), Acad. Press, N.Y. (1955).
61. M. Privat de Garilhe and M. Laskowski, J.B.C. 223, 661 (1956).
62. R.L. Sinsheimer and J.L. Koerner, J.B.C. 198, 293 (1952).
63. M. Goutier - Pirotte and A. Oth, B.B.A. 22, 394 (1956).
64. R. Setlow, Lecture delivered at University of British Columbia, Vancouver, March 1965.